

Ecological site group 029XESG03

Clay Uplands

Last updated: 04/13/2023
Accessed: 05/20/2024

Key Characteristics

- Uplands
- <75% bedrock outcrop
- Surface SAR <8 and Subsurface EC <8 and Surface EC <4
- EC <1.5 surface and <2 subsurface
- slope <35% or <40% surface rock
- Depth >55cm
- Rock <30% surface and <30% subsurface
- Clay >30% surface or >35% subsurface

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Physiography

This ESG is located on fan remnants.

Vegetation dynamics

The modal ecological site for this ESG is R029XY029NV Loamy 10-12 P.Z.

Major Land Resource Area

MLRA 029X
Southern Nevada Basin and Range

Subclasses

- F029XY066NV–PIMO-JUOS WSG 1R0501 12 to 16
- F029XY078NV–Shallow Ashy Loam 12-16" P.Z.
- R029XY002NV–SALINE MEADOW
- R029XY006NV–LOAMY 8-10 P.Z.
- R029XY029NV–LOAMY 10-12 P.Z.
- R029XY030NV–LOAMY 12-14 P.Z.
- R029XY104NV–SHALLOW CLAY LOAM 10-12 P.Z.
- R029XY165NV–ERODED NORTH SLOPE 12-14 P.Z.
- R029XY310UT–Upland Loam (Utah Serviceberry)

Correlated Map Unit Components

22641598, 22641321, 23237255, 23434853, 23434916, 23435162, 23151309, 23157806, 23157778, 23157782, 23157444, 23157448, 23157525, 23157527, 23237558, 23238047, 23238048, 23237561, 23237910, 22828501, 22828902, 22828904, 22828944, 22828963, 22728102

Stage

Provisional

Contributors

Curtis Talbot

Travis Nauman

State and transition model

Citations