

Ecological site F006XA001WA

Cool Frigid Xeric Ashy Slopes (Douglas-fir Cool Dry Grass)

Last updated: 9/11/2023
Accessed: 04/24/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 006X–Cascade Mountains, Eastern Slope

Major Land Resource Area (MLRA): 006X–Cascade Mountains, Eastern Slope

Stretching from northern Washington to southern Oregon, MLRA6 encompasses the mountain slopes, foothills, elevated plateaus and valleys on the eastern slopes of the Cascade mountains. This MLRA is a transitional area between the Cascade Mountains to the west and the lower lying Columbia Basalt Plateau to the east. Situated in the rain shadow of the Cascade Crest, this MLRA receives less precipitation than portions of the cascades further west and greater precipitation than the basalt plateaus to the east. Geologically, the majority of the MLRA is dominated by Miocene volcanic rocks, while the northern portion is dominated by Pre-Cretaceous metamorphic rocks and the southern portion is blanketed with a thick mantle of ash and pumice from Mount Mazama. The soils in the MLRA dominantly have a mesic, frigid, or cryic soil temperature regime, a xeric soil moisture regime, and mixed or glassy mineralogy. They generally are moderately deep to very deep, well drained, and loamy or ashy. Biologically, the MLRA is dominated by coniferous forest, large expanses of which are dominated by ponderosa pine, Douglas-fir or lodgepole pine. Areas experiencing cooler and moister conditions include grand fir, white fir, and western larch while the highest elevations include pacific silver fir, subalpine fir and whitebark pine. Economically, timber harvest and recreation are important land uses in these forests. Historically, many of these forests would have experienced relatively frequent, low and mixed severity fire favoring the development of mature forests dominated by ponderosa pine or Douglas-fir. In the southern pumice plateau forests, less frequent, higher severity fire was common and promoted the growth of large expanses of lodgepole pine forests.

LRU notes

Major Land Resource Area: 6 – Cascade Mountains, Eastern Slopes

Modal Land Resource Unit (LRU): Common Resource Area (CRA) 6.3 - Okanogan Pine / Fir Hills

This LRU occurs predominantly on slopes of hills and mountains. The soils are dominantly in the Andisols and Inceptisols taxonomic order, with some Mollisols. Soil parent materials are dominantly colluvium and residuum from igneous, sedimentary, and metamorphic rock, glacial outwash, and glacial till, with a mantle or mixture of volcanic ash in the upper part. Taxonomic soil climate is primarily a frigid temperature regime and xeric moisture regime with average annual precipitation of about 23 inches.

Other LRU'S where the site occurs:

CRA 6.2 - Pasayten / Sawtooth Highland

CRA 6.4 - Chelan Tephra Hills

CRA 6.5 - Chiwaukum Hills and Lowlands

Classification relationships

Forest Service Plant Associations:

CDG131 (WEN); CD-G1-31 (OKAN) – Douglas-fir/pinegrass (PSME/CARU)
 CDS655 (WEN) – Douglas-fir/bearberry/pinegrass (PSME/ARUV/CARU)
 CDS412 (WEN) – Douglas-fir/pachistima/pinegrass (PSME/PAMY/CARU)
 CDS675 (WEN) – Douglas-fir/bitterbrush/pinegrass (PSME/PUTR/CARU)

Associated sites

F006XA005WA	Cool Frigid Xeric Mountain Slopes (Douglas-fir Cool Moderately Dry Shrub/Herb) This site commonly has low huckleberry (VAMY) and other shrubs in the understory.
F006XD002WA	Cool Frigid Xeric Ashy Slopes (Grand fir Cool Dry Grass) Occurs on moister sites, generally associated in Chelan and northern Kittitas counties.
F006XA004WA	Cold Cryic Xeric Mountain Slopes (Subalpine fir Cold Dry Shrub) On colder sites.

Similar sites

F006XA005WA	Cool Frigid Xeric Mountain Slopes (Douglas-fir Cool Moderately Dry Shrub/Herb) This site commonly has low huckleberry (VAMY) and other shrubs in the understory.
F006XB001WA	Frigid Xeric Mountain Slopes (Douglas-fir Moderately Dry Shrub/Herb) A little dryer site.

Table 1. Dominant plant species

Tree	(1) <i>Pseudotsuga menziesii</i>
Shrub	Not specified
Herbaceous	(1) <i>Calamagrostis rubescens</i>

Physiographic features

This ecological site occurs mainly on mountain back slopes, shoulders and foot slopes, and glacial outwash terraces. It is found between 1,400 feet and 6,500 feet in elevation on all aspects. Slope gradients generally range from 20 to 40 percent, but can be found on slopes up to 90 percent.

Table 2. Representative physiographic features

Landforms	(1) Mountains > Mountain slope (2) Foothills > Hillslope (3) Outwash terrace
Flooding frequency	None
Ponding frequency	None
Elevation	1,600–4,500 ft
Slope	20–40%
Water table depth	80 in
Aspect	W, NW, N, NE, E, SE, S, SW

Table 3. Representative physiographic features (actual ranges)

Flooding frequency	Not specified
Ponding frequency	Not specified
Elevation	1,400–6,500 ft
Slope	0–90%
Water table depth	30–80 in

Climatic features

Mean Annual precipitation

Total Range: 15 – 45 inches

Central tendency: 20 – 25 inches

Mean Annual Air Temperature

Total Range: 2.5-9.7 C (36 to 49 F)

Central tendency: 5.2-7.0 C (41 to 45 F)

Frost-free period (days)

Total range: 75-135

Central tendency: 95-110

Table 4. Representative climatic features

Frost-free period (characteristic range)	95-110 days
Freeze-free period (characteristic range)	136 days
Precipitation total (characteristic range)	20-25 in
Frost-free period (actual range)	75-135 days
Freeze-free period (actual range)	136 days
Precipitation total (actual range)	15-45 in
Frost-free period (average)	
Freeze-free period (average)	136 days
Precipitation total (average)	




Figure 1. Monthly precipitation range




Figure 2. Monthly minimum temperature range




Figure 3. Monthly maximum temperature range




Figure 4. Monthly average minimum and maximum temperature




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

Climate stations used

- (1) MAZAMA [USC00455133], Mazama, WA

Influencing water features

Soil features

This ecological site is associated with several soil mapunit components. The components are dominantly Andic and Vitrandic taxonomic subgroups of Dystroxept and Haploxept great groups, and Vitrixerands in the Andisol taxonomic order. Soils are dominantly moderately deep to very deep and have average available water capacity of about 4.6 inches (11.7 cm) in the 0 to 40-inches (0-100 cm) depth range. Soil parent material is dominantly volcanic ash deposits over glacial till, glacial outwash, and colluvium and residuum from granitic, volcanic, metamorphic, and sedimentary rock.

Dominant Soil Series:

Brevco, Koepke, Louploup, Merkel, Molson, Nevine, Palmich, Radercreek, Ramparter, Wapal, Wilma

Parent Materials:

Kind – volcanic ash, glacial till, glacial outwash, colluvium, residuum

Origin – granitic, volcanic, metamorphic, and sedimentary rock




Figure 7. Map of soil mapunits with a major component linked to the Douglas-fir Cool Dry Grass Ecological Site

Table 5. Representative soil features

Surface texture	(1) Ashy fine sandy loam (2) Ashy sandy loam
Family particle size	(1) Loamy-skeletal (2) Ashy over loamy-skeletal (3) Ashy over loamy (4) Ashy-pumiceous
Drainage class	Well drained
Depth to restrictive layer	20–60 in
Surface fragment cover <=3"	0–30%
Surface fragment cover >3"	0–25%
Available water capacity (0-40in)	1.2–8.9 in
Soil reaction (1:1 water) (0-10in)	6.1–7.3
Subsurface fragment volume <=3" (Depth not specified)	5–55%
Subsurface fragment volume >3" (Depth not specified)	0–60%

Table 6. Representative soil features (actual values)

Drainage class	Moderately well drained to somewhat excessively drained
Depth to restrictive layer	Not specified
Surface fragment cover <=3"	0–30%
Surface fragment cover >3"	0–25%
Available water capacity (0-40in)	1.2–8.9 in
Soil reaction (1:1 water) (0-10in)	6.1–7.3
Subsurface fragment volume <=3" (Depth not specified)	5–55%
Subsurface fragment volume >3" (Depth not specified)	0–60%

Ecological dynamics

Open park-like stands of large diameter Douglas-fir, western larch and ponderosa pine with a pinegrass dominated understory. Frequent low severity ground fires every 10-20 years kept understory natural tree establishment low and maintained pinegrass cover. Overstory tree composition depends on seed source and moisture conditions.

Douglas-fir and western larch dominate stand composition with low percentages of ponderosa pine depending on site disturbance. On the warmer portions of this site at lower elevations ponderosa pine will be dominate with scattered Douglas-fir and western larch. This site occurs on southern exposures or ridgetops at higher elevations (4000 - 5000 feet) and on flats or outwash plains with cool air drainage at lower elevations. Most of the soils are granitic or glacial till with a volcanic ash cap. Lodgepole pine can be present in overstory and understory depending on seed source and cool air drainage conditions and fire occurrence. Fire exclusion has cause natural regeneration to advance from seedling to high density pole stands underneath the larger older trees creating ladder fuels and increased potential for stand replacing fire capable of causing mortality to the older large diameter overstory. Douglas-fir is susceptible to budworm and tussock moth defoliation during drought years. Dwarf mistletoe can be present in Douglas-fir and western larch.


This ecological site encompasses 4 Douglas-fir plant associations incorporated into the Douglas-fir Cool Dry Grass biophysical group:

Douglas-fir/pinegrass, Douglas-fir/bearberry/pinegrass, Douglas-fir/pachistima/pinegrass, and Douglas-fir/bitterbrush/pinegrass.

The most dominant is the Douglas-fir/pinegrass association. It is located north of Lake Chelan in CRA 6. This ecological site is based on this plant association. The Douglas-fir/bearberry/pinegrass association occurs in the west area of CRA 6.3 and eastern half of Lake Chelan in CRA 6.4. The Douglas-fir/pachistima/pinegrass association occurs in the southern area of CRA 6.3 just north of Lake Chelan. The Douglas-fir/bitterbrush/pinegrass association occurs in the western portion of CRA 6.3 in close proximity to the DF/bearberry/pinegrass association.

The main difference ecologically in these 4 plant associations is the occurrence and abundance of western larch and shrub species. In the DF/pinegrass and DF/bearberry/pinegrass associations western larch is common in the overstory and can regenerate after disturbance. In the DF/pachistima/pinegrass and DF/bitterbrush/pinegrass associations western larch is absent or minor. Ponderosa pine is common in all these plant associations. Another main difference is understory species abundance. In the DF/pinegrass association pinegrass dominates the coverage. The other 3 plant associations there is more abundance and coverage of shrubs with pinegrass. These shrub/pinegrass associations were included in the DF Cool Dry Grass Ecological site as they are cooler than the DF/shrub/bluebunch wheatgrass associations and react similar to the main DF/pinegrass association in regards to disturbance, with the exception of western larch absence.

State and transition model



State 1 Reference State

Reference plant community phase consists of park-like stands of large diameter Douglas-fir, western larch, and ponderosa pine. Understory is dominated by a continual sward of pinegrass. Tree regeneration is scattered. Frequent ground fires every 10-15 years cause mortality in established seedlings/saplings, mainly Douglas-fir creating sites for new seedling establishment, and perpetuate the pinegrass understory. Fire severity is low as to not cause mortality to large diameter overstory. USFS habitat type and plant association publications recognize this site as Douglas-fir/pinegrass. Douglas-fir is the dominant tree species, however, with frequent fire western larch and ponderosa pine will dominate the larger overstory. In warmer areas, western larch is absent or occasionally found in openings, and ponderosa pine becomes the major species with Douglas-fir. Also, in these warmer areas shrubs such as bearberry, pachistma, spirea, serviceberry, snowberry, bitterbrush and ceanothus spp. are abundant. Fire exclusion has caused of this site to have high density stocking of the second level overstory from regeneration growing into pole to mature stands. This ladder fuel site condition has high potential for stand replacing fire even killing the larger older overstory. Major insects that can cause mortality are the fir engraver, western spruce budworm, and Douglas-fir beetle. The Douglas-fir tussock moth can also cause concern. Drought and dense stands developing from prolonged fire intervals enhance beetle attack and tree mortality. The western pine beetle and the mountain pine beetle can cause mortality in ponderosa pine. Dwarf mistletoe is a major concern in western larch causing major growth loss and mortality. It will also be found on Douglas-fir, ponderosa pine, and lodgepole pine. Dwarf mistletoe was found in over 40% of the acreages in the Eastern Cascades where Douglas fir was a major stand component. Again, dense stands underneath infected trees from prolonged fire intervals causes mistletoe to spread easier. Fuel loads and ladder fuels increase fire intensity as a result. Other diseases include Annosum, laminated, and Armillaria root rots. These root rot diseases are enhanced through soil compaction and root damage though selective logging and road building. Forage production estimates from NRCS Range 5 Plots in Okanogan County in DF/pinegrass sites are as follows based on overstory tree canopy. Forage production figures are in

pounds/acre for all vegetation below 4.5 feet (grass, sedges, forbs, shrubs, tree regeneration): Overstory tree canopy – Forage production 0 – 20% - 800 to 1200 lbs/ac 20 – 40% - 600 – 800 lbs/ac 40 – 70% - 400 – 600 lbs/ac 70+% 0 – 400 lbs/ac The bulk of the grass production was pinegrass and Idaho fescue. Small percentages of sedge and bluebunch wheatgrass were recorded. The most prominent forbs were lupine, basalmroot, pussytoes, and hawkweed. Shrubs included currant, snowbrush ceanothus, spirea, snowberry, kinnikinnik, and rose spp. Tree regeneration included Douglas-fir and ponderosa pine.

Dominant plant species

- Douglas-fir (*Pseudotsuga menziesii*), tree
- lodgepole pine (*Pinus contorta*), tree
- western larch (*Larix occidentalis*), tree
- kinnikinnik (*Arctostaphylos uva-ursi*), shrub
- Oregon boxleaf (*Paxistima myrsinites*), shrub
- white spirea (*Spiraea betulifolia*), shrub
- russet buffaloberry (*Shepherdia canadensis*), shrub
- snowbrush ceanothus (*Ceanothus velutinus*), shrub
- Saskatoon serviceberry (*Amelanchier alnifolia*), shrub
- Cascade barberry (*Mahonia nervosa*), shrub
- common snowberry (*Symphoricarpos albus*), shrub
- antelope bitterbrush (*Purshia tridentata*), shrub
- bluejoint (*Calamagrostis canadensis*), grass
- Ross' sedge (*Carex rossii*), grass
- Geyer's sedge (*Carex geyeri*), grass
- bluebunch wheatgrass (*Pseudoroegneria spicata*), grass
- heartleaf arnica (*Arnica cordifolia*), other herbaceous
- common yarrow (*Achillea millefolium*), other herbaceous
- white hawkweed (*Hieracium albiflorum*), other herbaceous
- silky lupine (*Lupinus sericeus*), other herbaceous
- arrowleaf balsamroot (*Balsamorhiza sagittata*), other herbaceous
- Scouler's woollyweed (*Hieracium scouleri*), other herbaceous

Community 1.1 Open Park-like



Park-like widely spaced 125+ year old overstory of Douglas-fir, western larch, and ponderosa pine with understory dominated with pinegrass. Tree regeneration scattered and sporadic. Frequent ground fires every 10-20 years cause tree regeneration to be low. Overstory tree species composition dependent on seed source and moisture spectrum of site. Cooler moister locations have more Douglas-fir and western larch. Ponderosa pine dominate drier locations and have an increased understory cover of kinnikinnik. (Restoration to open overstory through tree removal and understory burning.)

Resilience management. 1.1A – Frequent low intensity ground fires maintain open stand and pinegrass understory

Community 1.2 Stand Replacement



Stand replacing fire creates a grass and potential shrub dominated site. Pinegrass will dominate the site and potentially snowbrush ceanothus shrub species can dominate in some areas where fire was severe. Severe fire damage to soil can cause ceanothus shrub fields for many years causing site to go to Alternative State 3. In low or mix severity fires natural regeneration will be sporadic depending on seed source and plant competition.

Community 1.3 Stem Exclusion



Natural regeneration growing to the pole stage can consist of Douglas-fir, western larch, and ponderosa pine. Lodgepole pine can also form dense stands depending on seed source and cool air drainage areas. Lack of sun on forest floor due to tree crown density understory is sparse with sprigs of pinegrass scattered. When tree canopy thins from competition, management, insects/disease, windthrow, and/or fire, pinegrass will rapidly increase in coverage.

Community 1.4 Understory Reinitiation



Overstory starts to thin out or gaps created by insect/disease mortality or windthrow allowing understory to develop. Snags and large woody present. Pinegrass and forbs increase and regeneration of Douglas-fir can occur.

Pathway 1.1B
Community 1.1 to 1.2



Open Park-like



Stand Replacement

Stand replacement fire caused by weather patterns prolonging fire return interval allowing tree regeneration layer to grow into sapling/pole stands creating ladder fuels.

Pathway 1.2A
Community 1.2 to 1.3



Stand Replacement



Stem Exclusion

Time, fire interval extended to allow regeneration to grow into pole stands.

Pathway 1.3B
Community 1.3 to 1.2



Stem Exclusion



Stand Replacement

Stand replacing fire

Pathway 1.3A
Community 1.3 to 1.4



Stem Exclusion



Understory Reinitiation

Time, low fire intensity, pole stands survive into mature stand, scattered mortality occurs from competition, insects, and disease.

Pathway 1.4A Community 1.4 to 1.1



Understory Reinitiation



Open Park-like

Time, with mix severity and low severity fires opening up stand.

Pathway 1.4B Community 1.4 to 1.2



Understory Reinitiation



Stand Replacement

Stand replacing fire

State 2 Fire Exclusion

Since European settlement the fire regime has changed greatly reducing frequent low intensity ground fires that kept these forests more open, with reduced tree density and more resilient to severe intensity fires. In addition, years of hygrading (harvesting largest best trees on site), overgrazing, prevalence of many roads which create fire breaks, and tree plantations has created a different forest structure. Dense multi-layered forests of Douglas-fir underneath larger old pine, fir and larch now exist. Dense stands of understory Douglas-fir can have over 1000 stems per acre. This creates extensive ladder fuel conditions which lead to the potential for stand replacing fires.

Community 2.1 Multi-Storied Stand

Scattered overstory of large old Douglas-fir, western larch and ponderosa pine with dense stands of Douglas-fir underneath. This plant community is a risk of severe stand replacing fire capable of killing the larger old overstory. Intensive fuel load reduction is needed to open up sites in this condition followed by using prescribed fire.

State 3 Dense shrubfields caused by severe fire and soil degradation

Community 3.1 Shrubfields

Severe stand replacing fire causing duff and organic matter to be destroyed negatively affecting soil fertility. Ceanothus species, especially shiny leaf ceanothus, dominates the site for 50+ years. This shrub species can grow in low nitrogen soils caused by intensive hot fires. Tree regeneration nonexistent to sporadic not capable to

reclaiming site.

State 4 Pastureland Forest converted to introduced grasses

Community 4.1 Pasture Conversion

A portion this ecological site has been converted to pasture using introduced grasses like orchard grass. Restoration would include intensive site preparation and tree planting followed by prescribed fire in future when trees more resistant to fire damage.

Transition T1A State 1 to 2

Fire exclusion allowing regeneration to develop underneath old overstory.

Transition T1B State 1 to 3

Shrubfields caused by severe fire and soil degradation.

Transition T1C State 1 to 4

Forest converted to pastureland .

Restoration pathway R2A State 2 to 1

Time, intensive tree removal of second level DF stand followed by prescribed burning

Restoration pathway R3A State 3 to 1

Careful selection of sites to determine if tree planting success is warranted.

Restoration pathway R4A State 4 to 1

Intensive tree establishment practices followed by prescribed burning when trees more fire resistant.

Additional community tables

Table 7. Representative site productivity

Common Name	Symbol	Site Index Low	Site Index High	CMAI Low	CMAI High	Age Of CMAI	Site Index Curve Code	Site Index Curve Basis	Citation
ponderosa pine	<i>PIPO</i>	65	104	50	98	40	—	—	
Rocky Mountain Douglas-fir	<i>PSMEG</i>	53	78	43	98	100	—	—	
western larch	<i>LAOC</i>	57	63	57	63	50	—	—	

Inventory data references

Relationship to Other Established Classifications:

United States National Vegetation Classification (2008) - A3395 Douglas fir- P. Pine / Herbaceous Understory

Central Rocky Mt. Forest & Woodland Alliance

Washington Natural Heritage Program. Ecosystems of Washington State, A Guide to Identification, Rocchio and Crawford, 2015 - Northern Rocky Mt. Dry-Mesic Montane Mixed Conifer Forest (D. Fir – Pine)

USDA NRCS Common Resource Areas. 6.3 Pine/fir Hills, 6.4 Chelan Tephra Hills

Level III and IV Ecoregions of WA, US EPA, June 2010 – 77e Okanogan Pine-Fir Hills, 77f Chelan Tephra Hills

This ecological site includes the following USDA Forest Service Plant Associations: PSME/CARU,

PSME/ARUV/CARU, PSME/PAMY/CARU, and PSME/PUTR/CARU. Lillibridge et. al PNW GTR – 359, Oct. 1995

Other references

Forest Plant Associations of the Okanogan National Forest, R6-Ecol-132b-1983. Williams, Lillybridge. September, 1983

Forest Plant Associations of the Wenatchee National Forest, PNW-GTR-359. October 1995. Lillybridge et al.

NRCS Soil and Site Index data for MLRA B6 in form of excel spreadsheets.

SSURGO MLRA B6 Soil Component Forest Ecoclasses (Plant Associations)

MLRA 430 Ecological Site Douglas-fir Cool Dry Grass (Douglas-fir/pinegrass)

Contributors

Gary Kuhn

Carri Gaines

Steve Campbell

Approval

Kirt Walstad, 9/11/2023

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	04/24/2024
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

-
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**
-
5. **Number of gullies and erosion associated with gullies:**
-
6. **Extent of wind scoured, blowouts and/or depositional areas:**
-
7. **Amount of litter movement (describe size and distance expected to travel):**
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-
14. **Average percent litter cover (%) and depth (in):**
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-**

production):

16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
-

17. **Perennial plant reproductive capability:**
-