

Ecological site R010XA030ID South Slope Channery 11-13 PZ ARTRX/PSSPS

Last updated: 12/13/2023
 Accessed: 04/28/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Dave Franzen and Jacy Gibbs Intermountain Range Consultants 17700 Fargo Rd. Wilder, ID 83676
Contact for lead author	Brendan Brazee, State Rangeland Management Specialist USDA-NRCS 9173 W. Barnes Drive, Suite C, Boise, ID 83709
Date	03/28/2008
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** rills can occur on this site due to steep slopes in small areas that have few surface stones. The presence of flat stones on the surface reduces erosion.

2. **Presence of water flow patterns:** surface stones usually do not allow water flow patterns to develop but if present they are short and not extensive.

3. **Number and height of erosional pedestals or terracettes:** both are rare on this site. Some terracettes can occasionally develop uphill from the large bunchgrasses and shrubs.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** no data available, but expected to range from 5-10 percent.

5. **Number of gullies and erosion associated with gullies:** none.

6. **Extent of wind scoured, blowouts and/or depositional areas:** usually does not occur.
-
7. **Amount of litter movement (describe size and distance expected to travel):** fine litter in the interspaces may move up to 5 feet or further following a significant run-off event. Rocks can trap fine litter. Coarse litter generally does not move.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** values should range from 4 to 6 but needs to be tested.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** structure ranges from weak fine granular to weak fine subangular blocky. Soil organic matter (SOM) needs to be determined. The A or A1 horizon is typically 3 inches thick. Surface soil color is very dark grayish brown moist.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** bunchgrasses, especially deep-rooted perennials, slow run-off and increase infiltration. Tall shrubs accumulate snow in the interspaces. Surface stones slow water movement and rain-drop impact and increase infiltration.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** not present.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: tall shrubs
- Sub-dominant: cool season deep-rooted perennial bunchgrasses
- Other: perennial forbs
- Additional: shallow rooted grasses
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** foothills big sagebrush and antelope bitterbrush will become decadent in the absence of fire and ungulate grazing. Grass and forb mortality will occur as tall shrubs increase.
-
14. **Average percent litter cover (%) and depth (in):** annual litter cover in the interspaces will be 20-30 percent to a depth of <0.1 inch. Under the mature shrubs litter is greater than 0.5 inches deep. Fine litter can accumulate on the terracettes and behind surface stones.
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-**

production): is 1050 lbs. per acre in a year with normal precipitation and temperatures. Perennial grasses produce 25-45 percent of the total, forbs 5-15 percent, and shrubs 45-65 percent.

16. **Potential invasive (including noxious) species (native and non-native).** List species which **BOTH** characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is **NOT** expected in the reference state for the ecological site: includes cheatgrass, medusahead, bulbous bluegrass, rush skeletonweed, scotch thistle, and spotted and diffuse knapweed.
-

17. **Perennial plant reproductive capability:** all functional groups have the potential to reproduce in most years.
-