

Ecological site R010XC041OR SR Very Shallow Rockland 12-16 PZ

Last updated: 12/13/2023

Accessed: 05/10/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	James A. Cornwell, State Rangeland Management Specialist, NRCS, Idaho (Retired) Lee Brooks, Assistant State Conservationist, NRCS, Idaho (Retired).
Contact for lead author	
Date	09/12/2009
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** None.

2. **Presence of water flow patterns:** None.

3. **Number and height of erosional pedestals or terracettes:** None.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 50-70 %.

5. **Number of gullies and erosion associated with gullies:** None.

6. **Extent of wind scoured, blowouts and/or depositional areas:** None.

7. **Amount of litter movement (describe size and distance expected to travel):** fine. Litter movement typically would be two feet.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil stability values should range from 3-5 but needs to be verified.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** The surface structure is moderate medium platy and moderately fine subangular blocky. SOM is 1 to 3 percent. The A horizon thickness is 2 to 4 inches.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Above average plant cover (30-50 % basal and crown) mediates the rainfall impact. The root mass of perennial bunchgrasses provides significant soil stability.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None. Depth to bedrock, an indurated pan or bedrock is less than 10 inches.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: perennial bunchgrasses=large bunchgrass>
- Sub-dominant: deciduous shrub = small bunchgrass >
- Other: forbs
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Normal decadence would be expected in the perennial bunchgrasses and stiff sagebrush.
-
14. **Average percent litter cover (%) and depth (in):**
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** Favorable: 600; Normal: 400; Unfavorable: 200 lbs/ac/yr.
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state**

for the ecological site: cheatgrass and medusahead

17. **Perennial plant reproductive capability:** All species should be capable of reproducing most years.
-