

Ecological site R010XC043OR SR South 9-12 PZ

Accessed: 07/27/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Jeff Repp and Bruce Franssen	
Contact for lead author	State Rangeland Management Specialist	
Date	04/15/2005	
Approved by	Bob Gillaspy	
Approval date		
Composition (Indicators 10 and 12) based on	Annual Production	

Indicators

1110	ilicators
1.	Number and extent of rills: None.
2.	Presence of water flow patterns: None to few (on steeper slopes)
3.	Number and height of erosional pedestals or terracettes: None.
4.	Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground): 5-20%
5.	Number of gullies and erosion associated with gullies: None.
6.	Extent of wind scoured, blowouts and/or depositional areas: None.
7.	Amount of litter movement (describe size and distance expected to travel): Fine - limited movement

8.	Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values): Moderate to significant resistant to erosion: aggregate stability = 3-6				
9.	Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Moderate medium granular structure to weak medium platy structure, dry color value 3 - 6, 3 - 12 inches thick; Moderate OM (2-4%)				
10.	Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Moderate ground cover (50-70%) and moderate to steep slopes (12-70%) moderately limit rainfall impact and overland flow.				
11.	Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): None.				
12.	Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):				
	Dominant: Deep rooted perennial bunchgrasses				
	Sub-dominant: Evergreen shrubs >= perennial forbs				
	Other: Shallow rooted perennial bunchgrasses				
	Additional:				
13.	Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Normal decadence and mortality expected.				
14.	Average percent litter cover (%) and depth (in):				
15.	Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production): Favorable: 1000, Normal: 800, Unfavorable: 600 lbs/acre/year				
16.	Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Western Juniper readily invades the site. Cheatgrass and Medusahead invade sites that have lost deep rooted perennial grass functional group.				

erennai piant rep	productive capability	/: All species snot	uld be capable of	reproducing annu	ually.