

Ecological site R011XY013ID Saline Silty 7-10 PZ ATNU2/ACHY

Last updated: 4/06/2020

Accessed: 05/04/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Dave Franzen and Jacy Gibbs Intermountain Range Consultants 17700 Fargo Rd. Wilder, ID 83676
Contact for lead author	Brendan Brazee, State Rangeland Management Specialist USDA-NRCS 9173 W. Barnes Drive, Suite C, Boise, ID 83709
Date	06/17/2009
Approved by	Kendra Moseley
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- Number and extent of rills:** rills rarely occur on this site due to the relatively flat slopes.
- Presence of water flow patterns:** water-flow patterns rarely occur on this site. When they do occur they are short and disrupted by cool season grasses and shrubs. They are not extensive.
- Number and height of erosional pedestals or terracettes:** rarely occur on this site.
- Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** it likely ranges from 70-80 percent but additional data is needed.
- Number of gullies and erosion associated with gullies:** none.
- Extent of wind scoured, blowouts and/or depositional areas:** usually not present in the Reference State.

7. **Amount of litter movement (describe size and distance expected to travel):** fine litter in the interspaces typically moves up to three feet primarily by wind. Coarse litter generally does not move.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** values should range from 4 to 6 but needs to be tested.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** No data.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** bunchgrasses, especially deep rooted perennials, slow runoff and increase infiltration. Medium height shrubs accumulate some snow in the interspaces.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** not present.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: medium height shrubs
- Sub-dominant: cool season perennial grasses
- Other: perennial forbs
- Additional: shallow rooted perennial bunchgrasses
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** very little mortality or decadence is expected on this site. Mortality of shallow rooted grasses may occur due to extended periods of drought.
-
14. **Average percent litter cover (%) and depth (in):** additional data is needed but is expected to be low and at a shallow depth.
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** is 400 pounds per acre (448 Kg/ha) in a year with normal precipitation and temperatures. Perennial grasses produce 30-40 percent of the total production, forbs 1-5 percent, and shrubs 60-70 percent.
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that**

become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: includes cheatgrass, kochia, Russian thistle, annual mustards, and halogeton.

17. **Perennial plant reproductive capability:** all functional groups have the potential to reproduce in normal and favorable years.
-