

Ecological site R011XY027ID Wet Meadow Carex-Juncus

Last updated: 4/06/2020
 Accessed: 04/27/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Dave Franzen and Jacy Gibbs Intermountain Range Consultants 17700 Fargo Rd. Wilder, ID 83676
Contact for lead author	Brendan Brazee, State Rangeland Management Specialist USDA-NRCS 9173 W. Barnes Drive, Suite C, Boise, ID 83709
Date	06/18/2009
Approved by	Kendra Moseley
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** rills do not occur on this site.

2. **Presence of water flow patterns:** none. Water flows over and through the plant community. Rarely are flows detrimental to the plants. The plants have adapted or evolved with this occurrence.

3. **Number and height of erosional pedestals or terracettes:** neither occurs on this site. Some plants may be hummocked due to trampling damage.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** data is not available. On sites in mid-seral status bare ground may range from 2-10 percent.

5. **Number of gullies and erosion associated with gullies:** none.

6. **Extent of wind scoured, blowouts and/or depositional areas:** does not occur.

-
7. **Amount of litter movement (describe size and distance expected to travel):** fine litter in the interspaces may move 6 feet or more due to seasonal flooding. Litter accumulates on the surface. There is little or no coarse litter developed on the site and it will be removed from the site following seasonal flooding.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** values should range from 3 to 5 but needs to be tested.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** no data.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** rhizomatous sedges and rushes slow run-off and increase infiltration. The total vegetation cover should be >60 percent to optimize infiltration. The plant community does not depend on water infiltration alone, but on the water table. The water table controls rooting depth.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** not present. Compaction layers can develop under stock trails made by livestock going to and from water or from long-term repetitive heavy grazing.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Rhizomatous sedges and rushes
- Sub-dominant: perennial forbs
- Other: shrubs
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** normal mortality of grass and grass-like is slow and occurs as aging plants. This will go unnoticed due to regeneration from roots, seeds or other new plants filling the spaces.
-
14. **Average percent litter cover (%) and depth (in):** additional litter cover data is needed but is expected to be 45-60 percent to a depth of 0.5-1.5 inches. Litter accumulates on the soil surface.
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** is 4000 pounds per acre (4480 Kg/ha) in a year with normal precipitation and temperatures. Rhizomatous sedges and rushes produce 80-90 percent of the total production, forbs 5-15 percent and shrubs < 5 percent.
-

16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: includes whitetop, Leafy spurge, Dock, Canadian thistle, and reed canarygrass. Other invasive species may include redtop and Kentucky bluegrass. At lower elevations, purple loosestrife may invade.
-

17. **Perennial plant reproductive capability:** all functional groups have the potential to reproduce in most years. Most of the plants reproduce vegetatively.
-