

Ecological site R011XY028ID Marsh TYLA-SCAC3

Last updated: 4/06/2020
Accessed: 04/30/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 011X–Snake River Plains

Major Land Resource Area (MLRA): 011X – Snake River Plains
Precipitation or Climate Zone: Additional moisture site

Classification relationships

Land Resource Region: B (Northwest Wheat and Range)
MLRA: 11 (Snake River Plains)
EPA Eco Region: Level III (Snake River Plain)

Ecological site concept

Site does receive additional moisture
Soils are:
Not saline or saline sodic
Site is poorly drained, with a surface mat of root 4" or more.
Deep, with <3% coarse fragments (by volume). Not skeletal
Not strongly or violently effervescent in the surface mineral 10"
Slope is <30%
Clay content is =<35% in surface mineral 4"
Site does not have an argillic horizon with >35% clay

Associated sites

R011XY015ID	Loamy Bottom 8-14 PZ ARTRT/LECI4
R011XY019ID	Meadow DECA18-CANE2
R011XY020ID	Dry Meadow POSE-PHAL2

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

This site generally occurs on nearly level margins of streams, ponds, lakes, springs, and reservoirs, and as

stringers paralleling stream and river channels. This site can also occur as parts of old stream courses, oxbows, and potholes. Slopes are less than 2 percent. Elevations range between 2500-6800 feet (750-2100 meters).

Table 2. Representative physiographic features

Landforms	(1) Braided stream (2) Oxbow (3) Pothole
Flooding duration	Long (7 to 30 days) to very long (more than 30 days)
Flooding frequency	Frequent to very frequent
Ponding duration	Long (7 to 30 days) to very long (more than 30 days)
Ponding frequency	Frequent
Elevation	762–2,073 m
Slope	0–2%
Ponding depth	5–30 cm
Water table depth	0–3 cm
Aspect	Aspect is not a significant factor

Climatic features

MLRA 11 is part of Idaho’s Snake River Plain. The elevation ranges from 2,077 to 7,549 feet, with a mean of 3,992 feet. Most of the precipitation falls as rain in the fall, winter and spring. Very little precipitation occurs during the summer months. In general this MLRA receives more sun than the U.S. average during the summer, but less than average during the winter.

The average annual precipitation is 10.01 inches (based on 10 long term climate stations located throughout the MLRA), with minimum and maximum values of 8.38 and 11.62 inches, respectively.

The average annual temperature ranges from 38° to 65° Fahrenheit. With a maximum average temperature of 65 degrees F. and a minimum average of 38 degrees F. The frost free interval ranges from 139 to 165 days and the freeze free interval ranges from 168 to 196 days.

Table 3. Representative climatic features

Frost-free period (average)	165 days
Freeze-free period (average)	196 days
Precipitation total (average)	305 mm




Figure 1. Monthly precipitation range




Figure 2. Monthly average minimum and maximum temperature

Influencing water features

The Marsh site is influenced by additional water from either adjacent streams through seasonal flooding, water table, seeps or springs, or from run-on from adjacent sites. The site may include the following wetlands and stream types.

Soil features

Typically these soils are erratically stratified with a surface mat of roots 4 or more inches deep over 3 to 6 inches of organic material over clays, clay loams, or silty clay loams that are over 60 inches deep. These soils are poorly drained and alluvial in origin. The soils are usually slightly acid to slightly alkaline in pH. Soils are saturated throughout the growing season and usually have standing water at or above the surface into the late summer.

Flooding can be frequent during snowmelt and just after snowmelt. Ponding is normal. The plant community is dependent on saturated soils and standing water during the growing season.

Soil Series Correlated to this Ecological Site -

No data

Ecological dynamics

The dominant visual aspect of this site is broadleaf cattail and/or hardstem bulrush. Hardstem bulrush is more dominant where water is 6 to 8 inches deep or deeper throughout the growing season. Broadleaf cattail is often more dominant where the water is at the soil surface through most of the growing season and may dry out at the surface later in the summer. When the two species occur together, broadleaf cattail is in shallower water than the hardstem bulrush. This is common on margins of ponds, lakes, and reservoirs.

The site often occurs within a complex of wetland sites when adjacent to streams or rivers. In these situations, the soil surface of the area where the site is found is typically slightly undulating causing small depressions and high spots with variable soil moisture regimes. The plant communities found on these areas are sites within the complex. Characteristics of these sites are as follows:

1. Marsh site. Deeper depressions with the water table at or near the surface or slightly above the surface for the entire growing season. This site is dominated by broadleaf cattail, hardstem bulrush, and common threesquare.
2. Wet Meadow site. Shallow to depression areas with the water table at or near the surface for the entire growing season. This site is dominated by *Carex* spp. and *Junus* spp.
3. Meadow site. Slightly higher areas that are drier during the growing season. The water table is down to a depth of 20 to 40 inches by the end of the growing season. This site is dominated by tufted hairgrass and Nebraska sedge.
4. Dry Meadow site. Even higher areas that are even drier during the growing season and the water table is > 40 inches deep by the end of the growing season. The plant community is dominated by Nevada bluegrass and alpine

timothy.

Either hardstem bulrush or broadleaf cattail can form impenetrable monotypic stands. Composition by weight is 95 percent grass or grass-like and 5 percent forbs.

During the last few thousand years, this site has evolved in semi-arid and montane climates characterized by dry summers and cold, moist or wet winters. The site has evolved on deep alluvial soils that are saturated at the soil surface or standing water throughout most of the growing season. Herbivory has historically occurred on this site at very low levels of utilization. Herbivory by some small mammals such as muskrats does occur.

Fire has had little influence on the development of the site. Rare wildfires can occur following consecutive drought years, particularly broadleaf cattail stands that go dry late in the summer.

The conditions for the plant community of this site are variable due to differences in the duration and depth of standing water. Dominant species are mostly rhizomatous and can form monotypic stands.

An infinite number of combinations of factors that influence the ecology of potential plant communities exist. For practical purposes, four plant communities where the depth to the water table drives the vegetative composition have been described. They are:

- Dry meadow Water table at >40" at end of growing season
- Meadow Water table at 20-40" at end of growing season
- Wet meadow Water table at 10-20" at end of growing season
- Marsh Water at surface to <10" at end of growing season

Most wetland species have a wide range of tolerance for variations in soil moisture. Most species occur in more than one site, although most are dominant on just one site.

The following table shows the amplitude of wetland species that occur on the four sites.

- Ecological Amplitude of Meadow/Marsh Plants.

Grass and Grass-like Species

Scientific name Dry Meadow Meadow Wet Meadow Marsh

Leymus cinereus

Danthonia californica

Carex filifolia

Pascopyron smithii

Poa nevadensis

Juncus dudleyi

Muhlenbergia richardsonis

Hordeum brachyantherum

Phleum alpinum

Scientific name Dry Meadow Meadow Wet Meadow Marsh

Juncus balticus

Juncus torreyi

Alopecurus aequalis

Carex athrostachya

Calamagrostis canadensis

Deschampsia caespitosa

Carex nebrascensis

Glyceria striata

Carex lasiocarpa

Carex utriculata

Carex aquatilis

Eleocharis palustris

Carex rostrata

Carex hoodii

Carex exsiccata

Scirpus microcarpus

Juncus effusus

Beckmannia syzigachne

Typha latifolia

Schoenoplectus acutus

Schoenoplectus pungens

Sparganium erectum

Schoenoplectus tabernaemontani

Forb Species

Scientific name Dry Meadow Meadow Wet Meadow Marsh

Arnica fulgens

Pyrocoma lanceolata

Arenaria congesta

Artemisia ludoviciana

Achillea millefolium

Wyethia amplexicaulis

Pyrocoma uniflora

Ranunculus spp.

Trifolium spp.

Potentilla gracilis

Senecio integerrimus

Aster spp.

Cirsium scariosum

Symphyotrichum ascendens

Iris missouriensis

Senecio serra

Helianthus nuttallii

Camassia quamash

Epilobium ciliatum

Montia chamissoi

Plantago major

Alisma triviale

Cicuta douglassii

Argentina anserina

Scientific name Dry Meadow Meadow Wet Meadow Marsh

Veronica anagallis-aquatica

Symphyotrichum frondosum

Polygonum bistortoides

Triglochin maritimum

Polygonum amphibium

Symphyotrichum foliaceum

Potamogeton natans

Lemna minor

The Historic Climax Plant Community (HCPC), the Reference State (State 1), moves through many phases depending on the natural and man-made forces that impact the community over time. State 1, described later,

indicates some of these phases. The Reference Plant Community Phase is Phase A. This plant community is dominated by broadleaf cattail and/or hardstem bulrush. There are a few forbs that may occur in minor amounts including water knotweed, leafybract aster, floating pondweed, and common duckweed. There may also be other grass-like species occurring in minor amounts. The plant species composition of Phase A is listed later under "Reference Plant Community Phase Plant Species Composition". The Reference Plant Community is intentionally written broadly, but these species should occur in the plant community in variable amounts.

The total annual production is 4500 pounds per acre (5040 kilograms per hectare) in a normal year. Production in a favorable year is 5500 pounds per acre (6160 kilograms per hectare). Production in an unfavorable year is 3500 pounds per acre (3920 kilograms per hectare). Most of this variation occurs from temperature differences, not moisture. Structurally, rhizomatous grass and grass-like species are very dominant, followed by perennial forbs.

FUNCTION:

This site is suitable for waterfowl, some shore birds, and small mammals. The site is not suitable for livestock grazing or big game, although some deer may use the edges for cover. Standing water limits grazing opportunities. In some drought years, some grazing can occur with livestock although wet soils should be avoided.

This site can be used for waterfowl hunting and may be adjacent to open water that offer fishing opportunities.

The site is very resistant to degradation due to standing water, low value unpalatable forage, and limited grazing opportunities. Site degradation is usually the result of permanently lowering of standing water. This can occur with down cutting of adjacent stream channels. This can result from off-site conditions adjacent to the site or in the upper watershed. Once adjacent streams are down-cut, concentrated flows can lower the water table and standing water.

Impacts on the Plant Community:

Influence of fire:

This site usually does not burn from wildfire. A rare fire may occur following prolonged drought. If a fire occurs, it usually does not adversely affect the plant community due to rhizomatous species. Most plants sprout back.

Influence of improper grazing management:

Due to standing water and low value forage, grazing during the growing season is rare. Grazing may occur during the dormant period. If dry soil conditions occur into the spring and livestock remove the tops of either broadleaf cattail or hardstem bulrush, the plants can die if water then covers them. Care must be exercised in grazing adjacent sites to avoid downcutting of stream channels that can influence standing water in this site. The potential of the site can be lost.

Weather influences:

Annual precipitation has little direct influence on this site. Prolonged drought can affect amounts and duration of standing water and can change the composition of this site.

Carex and Juncus species can increase and broadleaf cattails and hardstem bulrush can decrease. Lower production can occur with below normal spring temperatures that affect both air and water. An early, hard freeze can occasionally kill some plants.

Influence of Insects and disease:

Periodic disease and insect outbreaks can affect vegetation health. An outbreak of a particular insect is usually influenced by weather but no specific data for this site is available.

Influence of noxious and invasive plants:

Purple loosestrife can become invasive, but most plants on this site are very competitive against most potentially

invasive species due to monotypic stands of plants with strong rhizomes.

Influence of wildlife:

This site is important for many species of waterfowl. Some small mammals and shore birds also use the site. Big game animals use the edges for cover. Total numbers are seldom high enough to adversely affect the plant community.

Watershed:

The largest threat to degradation of this site is the lowering of permanent standing water during the growing season. Off-site conditions can affect the gradient of adjacent stream channels that can affect the water table. If down-cutting of adjacent stream occurs, this can reduce the depth and duration of the standing water. High run-off events from the adjacent uplands can severely damage or change the normal stream channel on the site. As the standing water is permanently reduced, either in depth or duration, productive potential is lost. Eventually the standing water is no longer the driving factor for the plant community. The site may regress to a meadow or even a low seral upland site.

Plant Community and Sequence:


Transition pathways between common vegetation states and phases:

State 1 to State 2. Develops through permanently reducing the standing water during the growing season to the point that dry soil is at the surface during the growing season. This can occur with down-cutting of adjacent stream channel. The site has crossed the threshold. This state cannot be returned to State 1 without returning the standing water regime.



Practice Limitations:

There are severe limitations to facilitating or accelerated practices due to standing water.

State and transition model



Permanently reducing the standing water during the growing season.



PLANT LEGEND STATES 1 & 2

TYLA- Broadleaf cattail
 SCAC3- Hardstem bulrush

State 1
State 1 Phase A

Community 1.1
State 1 Phase A

Reference Plant Community Phase. This plant community is dominated by broadleaf cattail and/or hardstem bulrush. There are a few forbs that may occur in minor amounts including water knotweed, leafybract aster, floating pondweed, and common duckweed. There may also be other grass-like species occurring in minor amounts.




Figure 3. Plant community growth curve (percent production by month). ID0315, DRY MEADOW. State 1.

State 2
State 2

Community 2.1
State 2

This plant community may be similar to Wet Meadow, Meadow, Dry Meadow or upland sites in low to mid-seral status. Initially, *Carex* and *Juncus* species may increase or become dominant. These species can be lost as degradation continues. Forbs may increase as degradation becomes severe. This state developed due to permanent reduction of standing water. The site has crossed the threshold. This state cannot be returned to State 1 without returning the standing water regime.

Additional community tables

Animal community

Wildlife Interpretations.

Animal Community – Wildlife Interpretations

This ecological site provides habitat for unique wetland wildlife species. The duration of hydrology results in abundant herbaceous cover and invertebrate production which provides food and cover for waterfowl. Wetland dependant species utilizing the site include marsh wren, great blue heron, ibis, teal, mallard, yellow-head blackbird, red-winged blackbird, northern leopard frog, western toad, muskrat, and beaver. Large herbivores infrequently use the site due to the extreme hydrologic conditions and low nutritional value of the forage. Loss of site hydrology significantly reduces the habitat value of the adjacent ecological sites. Open water is seasonal being provided by seasonal runoff, ponding, flooding, seasonal high water table, and natural springs.

State 1 Phase 1.1 – Broadleaf Cattail/ Hardstem Bulrush Reference Plant Community (RPC): The RPC and associated hydrology exhibit deep ponded water, providing a diverse population of invertebrates (benthic and terrestrial) unique to wetlands and beneficial to water loving birds. Arthropods such as dragonflies, damselflies, midges, backswimmers, scuds, and waterfleas make up a large portion of potential invertebrates in these wetlands. They are a basic food source for many higher trophic level organisms. The reptile and amphibian community is represented by western skink, rubber boa, western rattlesnake, western toad, boreal chorus frog, long-toed salamander, and northern leopard frog. A diverse amphibian population is a key indicator of good ecological health on this site. Loss of hydrology will limit or exclude amphibians from this ecological site. The RPC provides nesting, rearing, roosting, and forage for bird species such as Canada goose, American wigeon, mallard, pied billed grebe, great blue heron, sora, whimbrel, Wilson’s snipe, marsh wren, and long-billed curlew. Small mammal populations are extremely limited due to hydrologic conditions. Deer mouse, montane vole, and western jumping mouse would utilize the site for water throughout the year or during the winter when the water is frozen.

State 2 *Carex* sp./ *Juncus* Sp./ Forbs/ Grasses Plant Community: The animal community may be similar to Wet Meadow, Meadow, Dry Meadow, or upland sites in low to mid-seral status depending on the severity of hydrology manipulation of State 1 Phase 1.1. The management of the resulting plant communities will be a major factor in determining the diversity and populations of the animal community. Refer to the above mentioned ecological sites

for MLRA 11 for descriptions of the potential wildlife community.

Grazing Interpretations.

Due to standing water and low value unpalatable forage, grazing during the growing season is rare. Grazing may occur during the dormant period. If dry soil conditions occur into the spring and livestock remove the tops of either broadleaf cattail or hardstem bulrush, the plants can die if water then covers them. Care must be exercised in grazing adjacent sites to avoid downcutting of stream channels that can influence standing water in this site. The potential of the site can be lost.

Estimated initial stocking rate will be determined with the landowner or decision-maker. They will be based on the inventory, past use history and type, condition of vegetation, production, season of use, and seasonal preference. Calculations used to determine estimated initial stocking rate will be based on forage preference ratings.

Hydrological functions

No data.

Recreational uses

This site presents an aesthetically pleasing view of lush vegetation consisting primarily of grass-like plants and standing water. Hikers and fisherman often traverse the edges of this site. Fishing opportunities often exist on open waters adjacent to this site. Waterfowl hunting can occur on or adjacent to the site.

Wood products

None.

Other products

None.

Other information

Field Offices

Marsing, ID

Payette, ID

Emmett, ID

Meridian, ID

Caldwell, ID

Mountain Home, ID

Weiser, ID

Inventory data references

Information presented here has been derived from NRCS clipping and other inventory data. Also, field knowledge of range-trained personnel was used. Those involved in developing this site description include:

Dave Franzen, co-owner, Intermountain Rangeland Consultants, LLC

Jacy Gibbs, co-owner, Intermountain Rangeland Consultants, LLC

Jim Cornwell, Range Management Specialist, IASCD

Dan Ogle, Plant Materials Specialist, NRCS, Idaho

Chris Hoag, Wetland Plant Ecologist, NRCS, Idaho

Brendan Brazee, State Rangeland Management Specialist, NRCS, Idaho

Leah Juarros, Resource Soil Scientist, NRCS, Idaho
Lee Brooks, Range Management Specialist, IASCD

Other references

USDA, NRCS. 2001. The PLANTS Database, Version 3.1 (<http://plants.usda.gov>.) National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

USDA NRCS. 1992. Major Land Resource Area, Owyhee High Plateau, Range Site Descriptions. Reno, Nevada.

USDA NRCS. Major Land Resource Area, Owyhee High Plateau, Range Site Descriptions. Portland, Oregon.

USDA, Forest Service. 2004. (www.fs.fed.us/database/feis/plants).

USDI, BLM and Idaho Conservation Data Center. 2001. Riparian and Wetland Plant Associations of Southwestern Idaho.

USDI Bureau of Land Management, US Geological Survey; USDA Natural Resources Conservation Service, Agricultural Research Service; Interpreting Indicators of Rangeland Health. Technical Reference 1734-6; Version 4-2005.

Approval

Kendra Moseley, 4/06/2020

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Dave Franzen and Jacy Gibbs Intermountain Range Consultants 17700 Fargo Rd. Wilder, ID 83676
Contact for lead author	Brendan Brazee, State Rangeland Management Specialist USDA-NRCS 9173 W. Barnes Drive, Suite C, Boise, ID 83709
Date	06/18/2009
Approved by	Kendra Moseley
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** rills do not occur on this site.

2. **Presence of water flow patterns:** water-flow patterns do not occur.

3. **Number and height of erosional pedestals or terracettes:** neither occurs on this site.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** none.

-
5. **Number of gullies and erosion associated with gullies:** none.
-
6. **Extent of wind scoured, blowouts and/or depositional areas:** blowouts and depositional areas do not occur.
-
7. **Amount of litter movement (describe size and distance expected to travel):** litter may move in the standing water.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** values should range from 4 to 6 but needs to be tested.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** No data.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** site has standing water.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** is not present.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: rhizomatous species
- Sub-dominant: perennial forbs
- Other:
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** little mortality occurs on this site.
-
14. **Average percent litter cover (%) and depth (in):** not applicable.
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** is 4500 pounds per acre (5040 Kg/ha) in a year with normal amounts of standing water and temperatures. Rhizomatous species produce 90-95 percent of the total production and forbs less than 10 percent.
-

16. **Potential invasive (including noxious) species (native and non-native).** List species which **BOTH** characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is **NOT** expected in the reference state for the ecological site: includes purple loosestrife and common reed.
-

17. **Perennial plant reproductive capability:** all functional groups have the potential to reproduce in most years. Most of the plants can reproduce vegetatively.
-