

Ecological site R013XY002ID Stony Loam 13-16 PZ ARTRV/PSSPS

Last updated: 9/23/2020
Accessed: 05/18/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 013X–Eastern Idaho Plateaus

013X-Eastern Idaho Plateaus

Precipitation or Climate Zone: 13-16" P.Z.

<https://soils.usda.gov/survey/geography/mlra/index.html>

Classification relationships

Land Resource Unit: B (Northwestern Wheat and Range)

MLRA: 13 (Eastern Idaho Plateaus)

EPA EcoRegion: Level III (Middle Rockies)

Artemisia vaseyana/*Agropyron spicatum* HT in "Hironaka, M., M.A. Fosberg, A. H. Winward. 1983. Sagebrush-Grass Habitat Types of Southern Idaho. University of Idaho, Moscow, Idaho. Bulletin Number "35."

Ecological site concept

Site does not receive any additional water.

Soils are:

not saline or saline-sodic.

moderately deep, deep, with >35% stone (10-25") and boulder (>25") cover. skeletal within 20" of soil surface.

not strongly or violently effervescent in surface mineral 10".

textures usually range from very fine sandy loam to clay loam in surface mineral 4".

Slope is < 30%.

Clay content is = <32% in surface mineral 4".

Site does not have an argillic horizon with > 35% clay.

Associated sites

R013XY001ID	Loamy 12-16 PZ
R013XY005ID	Loamy 16-22 PZ ARTRV/FEID-PSSPS
R013XY013ID	Stony 12-16 PZ ARTRV/FEID

Similar sites

R013XY002ID	Stony Loam 13-16 PZ ARTRV/PSSPS
-------------	--

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Artemisia tridentata</i> ssp. <i>vaseyana</i> (2) <i>Purshia tridentata</i>
Herbaceous	(1) <i>Pseudoroegneria spicata</i> (2) <i>Purshia tridentata</i>

Physiographic features

This site occurs on fan remnants, hillslopes, lava plains, and on some mountain slopes. Occasional rock outcrops are associated with this site. Slopes predominantly range from 5 to 30 percent and occur on all aspects. Elevations range from 4500 to 7500 feet (1375 to 2300 meters).

Table 2. Representative physiographic features

Landforms	(1) Hill (2) Fan remnant (3) Lava plain
Flooding frequency	None
Ponding frequency	None
Elevation	1,372–2,286 m
Slope	5–30%
Water table depth	152 cm
Aspect	Aspect is not a significant factor

Climatic features

MLRA 13, the Eastern Idaho Plateaus, is part of the Northwestern Wheat and Range Region. Its elevation ranges from 4209 to 9331 feet above sea level, with an average elevation of 5787 feet. The average annual precipitation is 16.41 inches, with a range of 13.56 to 18.75 inches, based on ten long term climate stations located throughout the MLRA. A spike in precipitation amount often occurs in late spring, usually in May.

Temperatures vary widely in the MLRA throughout the year. A maximum temperature of 103° Fahrenheit occurred at the McCammon climate station (# 105716; elevation 4770 feet), while a minimum of -41° was recorded at the Kilgore station (#104908). At all stations temperatures throughout the year are usually below the national average. Kilgore also recorded the greatest annual snowfall amount of 217 inches. The average temperature is 41.4 degrees F. with an average high of 55.3 degrees and an average low of 27.5 degrees.

The frost-free period ranges from 64 to 90 days, while the freeze-free period can be 98 to 123 days.

Table 3. Representative climatic features

Frost-free period (average)	90 days
Freeze-free period (average)	123 days
Precipitation total (average)	483 mm




Figure 1. Monthly precipitation range




Figure 2. Monthly average minimum and maximum temperature

Influencing water features

This site is not influenced by adjacent wetlands, streams, or run on.

Soil features

The soils in this site have gravelly, stony, and very stony loam and silty loam surface textures. Coarse fragments are present in the soil profile generally increasing with depth, and can exceed 35 percent by volume. The soils are moderate deep to very deep to bedrock. They are formed primarily in alluvium, loess and colluvium derived from sedimentary deposits, limestone, rhyolite and quartzite. Permeability is moderately slow to moderate and the soils are well to somewhat excessively drained. The available water holding capacity (AWC) is very low to low. These soils are characterized by a xeric soil moisture regime and either a frigid or cryic soil temperature regime.

Soil Series Correlated to this Ecological Site

Acord
 Araveton
 Firading
 Jacanyon
 Lizzant
 Mike
 Poisonhol
 Rammel
 Ridgecrest
 Rexburg
 Tahquats
 Vipont
 Wahtigup

Table 4. Representative soil features

Surface texture	(1) Stony loam (2) Very stony silt loam (3) Gravelly
Drainage class	Well drained to somewhat excessively drained
Permeability class	Moderately slow to moderate
Soil depth	51–152 cm
Surface fragment cover ≤3"	0–25%
Surface fragment cover >3"	0–35%
Available water capacity (0-101.6cm)	4.32–15.24 cm
Calcium carbonate equivalent (0-101.6cm)	0–40%
Electrical conductivity (0-101.6cm)	0–2 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0–2
Soil reaction (1:1 water) (0-101.6cm)	5.6–8.4
Subsurface fragment volume ≤3" (Depth not specified)	15–45%
Subsurface fragment volume >3" (Depth not specified)	0–35%

Ecological dynamics

Ecological Dynamics of the Site:

The dominant visual aspect of this site is mountain big sagebrush in the overstory with bluebunch wheatgrass in the understory. Composition by weight is approximately 60 to 70 percent grasses, 5 to 15 percent forbs, and 20 to 30 percent shrubs.

During the last few thousand years, this site has evolved in a semi-arid climate characterized by dry summers and cold, wet winters. Herbivory has historically occurred on this site at low levels of utilization. Herbivores include mule deer, Rocky Mountain elk, some pronghorn antelope, and lagomorphs.

Fire has historically occurred on the site at intervals of 20-50 years.

The Historic Climax Plant Community (HCPC), the Reference State (State 1), moves through many phases depending on the natural and man-made forces that impact the community over time. State 1, described later, indicates some of these phases. The Reference Plant Community Phase is Phase A. This plant community is dominated by bluebunch wheatgrass and mountain big sagebrush. Antelope bitterbrush is usually present. Other subdominant species can include Idaho fescue, Sandberg bluegrass, arrowleaf balsamroot, and tapertip hawksbeard. A wide variety of other grasses, forbs and shrubs occur in small amounts. The plant species composition of Phase A is listed later under “Reference Plant Community Phase Plant Species Composition”.

Total annual production is 1000 pounds per acre (1120 kilograms per hectare) in a normal year. Production in a favorable year is 1400 pounds per acre (1568 kilograms per hectare). Production in an unfavorable year is 600 pounds per acre (672 kilograms per hectare). Structurally, cool season deep rooted perennial bunchgrasses are very dominant, followed by tall shrubs being more dominant than perennial forbs while shallow rooted perennial bunchgrasses are subdominant.

FUNCTION:

This site is well suited for big game in the late spring and fall. It is also well suited for livestock use in the late spring, early summer, and fall. Recreation opportunities are limited due to the surface stones.

Due to the relatively high rainfall, elevation, and surface stones on this site, it is fairly resistant to disturbances that can potentially degrade the site.

Due to the gentle topography, infiltration is normally high and runoff low. Runoff, when it does occur is non-erosive except during high intensity convection storms. Snow accumulates on the site due to high elevation and presence of tall shrubs

Impacts on the Plant Community.

Influence of fire:

In the absence of normal fire frequency, mountain big sagebrush and antelope bitterbrush can gradually increase on the site. Utah juniper can invade the site if a seed source is in the proximity. Grasses and forbs decrease as shrubs increase. With the continued absence of fire, mountain big sagebrush and/or juniper can displace most of the primary understory species. See "Influence of juniper invasion" below.

When fires become more frequent than historic levels (20-50 years), mountain big sagebrush and antelope bitterbrush are reduced significantly. Rabbitbrush and threetip sagebrush can be present due to sprouting from the root crown and may increase. With continued short fire frequency, mountain big sagebrush, antelope bitterbrush, and many of the other shrubs can be completely eliminated along with many of the desirable understory species such as bluebunch wheatgrass and Idaho fescue. These species may be replaced by Sandberg bluegrass along with a variety of annual and perennial forbs including noxious and invasive plant species. Cheatgrass will invade the site at lower elevations. These fine fuels will increase the fire frequency.

Influence of improper grazing management:

Season-long grazing and/or excessive utilization can be very detrimental to this site. This type of management leads to reduced vigor of the bunchgrasses. With reduced vigor, recruitment of these species declines. As these species decline, an increase in mountain big sagebrush and noxious and invasive plants will occur. The plant community also becomes susceptible to Utah juniper invasion.

Continued improper grazing management influences fire frequency by increasing fine fuels. As cheatgrass increases and becomes co-dominant with Sandberg bluegrass and other annuals, fires become more frequent, particularly at lower elevations.

Proper grazing management that addresses frequency, duration, and intensity of grazing can also keep fine fuels from developing, thereby reducing fire frequency. This can lead to gradual increases in mountain big sagebrush and juniper. A planned grazing system can be developed to intentionally accumulate fine fuels in preparation for a prescribed burn. Any brush management should be carefully planned as a reduction in shrubs without a suitable understory of perennial grasses can lead to an increase in fine fuels which can lead to more frequent fire intervals. There can be shrub species present that are extremely important to wildlife and need to be protected if control measures are implemented. Loss of the shrub component on this site can have very negative impacts on wildlife.

Weather influences:

Above normal precipitation in April, May, and June can dramatically increase total annual production of the plant community. These weather patterns can also increase viable seed production of desirable species to provide for recruitment. Likewise, below normal precipitation during these spring months can significantly reduce total annual production and be detrimental to viable seed production. Overall plant composition is normally not affected when perennials have good vigor. Below normal temperatures in the spring can have an adverse impact on total production regardless of the precipitation. An early, hard freeze can occasionally kill some plants.

Prolonged drought adversely affects this plant community in several ways. Vigor, recruitment, and production are usually reduced. Mortality can occur. Prolonged drought can lead to a reduction in fire frequency.

Influence of Insects and disease:

Outbreaks can affect vegetation health. Bitterbrush can be severely affected by the western tent caterpillar (*Malacosoma fragilis*). Two consecutive years of defoliation by the tent caterpillar can cause mortality in bitterbrush. Mormon cricket and grasshopper outbreaks occur periodically. Outbreaks seldom cause plant mortality since defoliation of the plant occurs only once during the year of the outbreak. Snow mold can reduce the vigor of mountain big sagebrush.

Influence of noxious and invasive plants:

Many of these species add to the fine-fuel component and lead to increased fire frequency. Perennial and annual invasive species compete with desirable plants for moisture and nutrients. The result is reduced production and change in composition of the understory.

Influence of wildlife:

Big game animals use this site in the spring and fall. Their numbers are seldom high enough to adversely affect the plant community. Herbivory can be detrimental to bitterbrush when livestock grazing and browsing by big game occurs at the same time and season. This will occur when both kinds of animal are using the plant in the late summer or fall. The adverse impact is excessive use of the current year's leader growth. The deer mouse is beneficial to this site as it is the principal vector for planting bitterbrush seed.

Watershed:

Decreased infiltration and increased runoff occur with an increase in mountain big sagebrush. Desired understory species can be reduced. This composition change can affect nutrient and water cycles. Increased runoff also causes sheet and rill erosion. Abnormally short fire frequency also gives the same results, but to a lesser degree. The long-term effect is a transition to a different state.

Influence of Utah juniper invasion:

In plant communities that are invaded by juniper, the species has a competitive advantage for the following reasons:

- Juniper is very drought tolerant.
- It has the ability to extract soil moisture from a wide range of soil depths.
- Juniper has high evapo-transpiration rates.
- The species intercepts rain and snow before it reaches the soil surface.
- It has the ability to grow as long as there is soil moisture and the temperature is above freezing.
- Juniper has a relatively rapid growth rate and is long-lived. It can readily over-top shade intolerant species which leads to mortality.
- Nutrient cycling is reduced.
- As the canopy closes, juniper gains control of energy capture.

As juniper extracts water, other plants are unable to acquire sufficient water and nutrients to sustain growth and reproduction, thus reducing cover and biomass in the interspaces. After the canopy closes, there is sufficient soil moisture available for shallow-rooted, shade tolerant species to persist directly under the tree.

The following hydrologic impacts occur on sites invaded by juniper:

- Infiltration in the interspaces is reduced.
- Run-off increases resulting in increased sheet and rill erosion with elevated sediment loads.
- Soil temperatures increase in the interspaces which results in accelerated drying of the soil surface.
- Increased bare ground in the interspaces.
- Soil moisture storage is reduced.

As bare ground and interconnectiveness of bare ground increases, flow rates are accelerated (reduction of flow

sinuosity) and run-off out of the area increases.

Degradation of these systems can result in the formation of a feedback cycle in which greater juniper cover and density results in greater plant and soil disturbance between the canopies.

In summary, a closed juniper community takes control of the following ecological processes: (1) hydrology, (2) energy capture, and (3) nutrient cycling. The changes are primarily driven by the hydrologic processes. The development of a closed juniper canopy always results in a transition across the threshold to a different state. Generally, when juniper canopy cover nears 20%, the plant community is approaching the threshold.

Plant Community and Sequence:

Transition pathways between common vegetation states and phases:

State 1.

Phase A to B. Develops in the absence of fire and improper grazing management. There is a Utah juniper seed source present in the vicinity near the site.

Phase A to C. Results from a fire or prescribed burning.

Phase A to D. Results from improper grazing management and no fire. There is no Utah juniper seed source present in the vicinity.

Phase B to C. Results from a wildfire, prescribed burning, or brush management.

Phase B to A. Occurs with prescribed grazing and brush management or prescribed burning.

Phase C to A. Results from prescribed grazing and no fire.

Phase D to A. Occurs with prescribed grazing.

Phase D to C. Occurs with fire or prescribed burning and prescribed grazing.

Phase C to D. Occurs with no fire and improper grazing management.

State 1 Phase C & D to State 2. Develops through improper grazing management and frequent fire. This site has crossed the threshold. It is economically impractical to return this state to State 1 with accelerated practices.

State 1, Phase B to 3. Develops with no fire and improper grazing management from a juniper invaded phase of State 1. This site has crossed the threshold. It is economically impractical to return this state to State 1 with accelerated practices.


State 2 to unknown site. Excessive soil loss and changes in the hydrologic cycle caused by improper grazing management and no fire or frequent fire cause this state to cross the threshold and retrogress to a new site with reduced potential. It is economically impractical to return this state to State 1 with accelerated practices.

State 3 to unknown site. Continued improper grazing management or the lack of fire cause this state to cross the threshold and retrogress to a new site with reduced potential due to significant soil loss and changes in hydrology. It is economically impractical to return this state to State 1 with accelerated practices.

Practice Limitations:

Slight to moderate limitations exist on this site for implementing vegetative management practices. Moderate to severe limitations exist on this site for implementing facilitating practices. Stoniness and occasional slopes over 30 percent require certain precautions be followed in placement of water developments, salting, and fencing. Severe limitations exist on this site for implementing accelerating practices. Stoniness and occasional slopes over 20 percent also require certain precautions be followed in seeding, brush management, prescribed burning, and certain mechanical means of treating the soil.

State and transition model



State 1

State 1 Phase A

State 1, Phase A, Reference Plant Community Phase. This plant community has mountain big sagebrush in the overstory with bluebunch wheatgrass in the understory. Antelope bitterbrush is usually present. Other subdominant

species can include Idaho fescue, Sandberg bluegrass, arrowleaf balsamroot, and tapertip hawksbeard. A wide variety of other grasses, forbs, and shrubs occur in small amounts. Natural fire frequency is 20-50 years.

Community 1.1

State 1 Phase A

Reference Plant Community Phase. This plant community has mountain big sagebrush in the overstory with bluebunch wheatgrass in the understory. Antelope bitterbrush is usually present. Other subdominant species can include Idaho fescue, Sandberg bluegrass, arrowleaf balsamroot, and tapertip hawksbeard. A wide variety of other grasses, forbs, and shrubs occur in small amounts. Natural fire frequency is 20-50 years.

Table 5. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	0%
Grass/grasslike foliar cover	0%
Forb foliar cover	0%
Non-vascular plants	0%
Biological crusts	0%
Litter	60-70%
Surface fragments >0.25" and ≤3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%




Figure 3. Plant community growth curve (percent production by month).
ID0805, B13 ARTRV . State 1.

Community 1.2

State 1, Phase B

This plant community is dominated in the overstory by mountain big sagebrush with some Utah juniper seedlings and saplings invading. Sandberg bluegrass is the dominant understory species. Bluebunch wheatgrass and other bunchgrasses have been reduced in amounts and are in low vigor. A wide variety of other grasses, forbs, and shrubs still may occur but in very small amounts and may be in low vigor. Antelope bitterbrush may still be present but in very small amounts and may be hedged. A Utah juniper seed source is present in nearby sites. Some cheatgrass may be present particularly at lower elevations. This state has developed due to fire frequency being much longer than normal and improper grazing management

Community 1.3

State 1, Phase C

This plant community is dominated by bluebunch wheatgrass. Sub-dominant species include Sandberg bluegrass, bottlebrush squirreltail, and thickspike wheatgrass along with other perennial grasses and forbs. Idaho fescue and Letterman's and Columbia needlegrasses have decreased and may have died out due to fire. Most shrubs are absent from the site due to recent fire, except for some rabbitbrush and threetip sagebrush (if present before the fire) may be present due to root sprouting. Some cheatgrass may have invaded the site at lower elevations. The community is a result of recent wildfire or prescribed burning.

Community 1.4

State 1, Phase D

This plant community is dominated by mountain big sagebrush in the overstory with significantly reduced amounts of bluebunch wheatgrass and in reduced vigor. Sandberg bluegrass, thickspike wheatgrass and bottlebrush squirreltail have increased. Other bunchgrasses have been reduced and are in low vigor. Antelope bitterbrush has reduced vigor and maybe hedged. There is no Utah juniper seed source in the proximity. Some cheatgrass may be present. This plant community has developed due to improper grazing management and no fire.

Pathway A to B

Community 1.1 to 1.2

Develops in the absence of fire and improper grazing management. There is a Utah juniper seed source present in the vicinity near the site.

Pathway A to C

Community 1.1 to 1.3

Results from a fire or prescribed burning

Pathway A to D

Community 1.1 to 1.4

Results from improper grazing management and no fire. There is no Utah juniper seed source present in the vicinity

Pathway B to A

Community 1.2 to 1.1

Occurs with prescribed grazing and brush management or prescribed burning

Pathway B to C

Community 1.2 to 1.3

Results from a wildfire, prescribed burning, or brush management

Pathway C to A

Community 1.3 to 1.1

Results from prescribed grazing and no fire

Pathway C to D

Community 1.3 to 1.4

Occurs with no fire and improper grazing management

Pathway D to A

Community 1.4 to 1.1

Occurs with prescribed grazing

Pathway D to C
Community 1.4 to 1.3

Occurs with fire or prescribed burning and prescribed grazing

State 2
State 2

This plant community is dominated by Sandberg bluegrass and annuals in the understory. Cheatgrass is a dominant annual in the community, particularly at lower elevations. There may be a variety of invasive forbs and some noxious plant species may have invaded the site. Some soil loss has occurred. The community has developed due to continued improper grazing management and frequent fire. This site has crossed the threshold. It is economically impractical to return this state to State 1 with accelerated practices.

Community 2.1
State 1 Phase B

This plant community is dominated in the overstory by mountain big sagebrush with some Utah juniper seedlings and saplings invading. Sandberg bluegrass is the dominant understory species. Bluebunch wheatgrass and other bunchgrasses have been reduced in amounts and are in low vigor. A wide variety of other grasses, forbs, and shrubs still may occur but in very small amounts and may be in low vigor. Antelope bitterbrush may still be present but in very small amounts and may be hedged. A Utah juniper seed source is present in nearby sites. Some cheatgrass may be present particularly at lower elevations. This state has developed due to fire frequency being much longer than normal and improper grazing management.

Table 6. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	0%
Grass/grasslike foliar cover	0%
Forb foliar cover	0%
Non-vascular plants	0%
Biological crusts	0%
Litter	60-70%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%




Figure 4. Plant community growth curve (percent production by month). ID0805, B13 ARTRV . State 1.

State 3
State 3

This plant community is dominated by Utah juniper with Sandberg bluegrass and annuals in the understory. Cheatgrass is a dominant annual at lower elevations. There are few shrubs present due to competition from junipers. Some deep-rooted perennials may be present under the junipers. . When shrub cover is below 10-15%, bare ground is above 25-30% and juniper cover is greater than 20%, the plant community has crossed the threshold. Some soil loss has occurred. This plant community has developed due to the lack of fire and continued improper grazing management. It is economically impractical to return this community to State 1 with accelerated practices.

Community 3.1
State 1 Phase C

This plant community is dominated by bluebunch wheatgrass. Sub-dominant species include Sandberg bluegrass, bottlebrush squirreltail, and thickspike wheatgrass along with other perennial grasses and forbs. Idaho fescue and Letterman's and Columbia needlegrasses have decreased and may have died out due to fire. Most shrubs are absent from the site due to recent fire, except for some rabbitbrush and threetip sagebrush (if present before the fire) may be present due to root sprouting. Some cheatgrass may have invaded the site at lower elevations. The community is a result of recent wildfire or prescribed burning.

Table 7. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	0%
Grass/grasslike foliar cover	0%
Forb foliar cover	0%
Non-vascular plants	0%
Biological crusts	0%
Litter	60-70%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%




Figure 5. Plant community growth curve (percent production by month). ID0805, B13 ARTRV . State 1.

State 4
State 4

Unknown new site. This plant community has gone over the threshold to a new site. Site potential has been

reduced. Significant soil loss has occurred. Infiltration has been reduced and run-off has become more rapid. This state has developed due to continued improper grazing management, no fire, or frequent fires from State 2 or with continued improper grazing management and the absence of fire from the juniper dominated phase of State 3. It is economically impractical to return this community to State 1 with accelerated practices

Community 4.1
State 1 Phase D

This plant community is dominated by mountain big sagebrush in the overstory with significantly reduced amounts of bluebunch wheatgrass and in reduced vigor. Sandberg bluegrass, thickspike wheatgrass and bottlebrush squirreltail have increased. Other bunchgrasses have been reduced and are in low vigor. Antelope bitterbrush has reduced vigor and maybe hedged. There is no Utah juniper seed source in the proximity. Some cheatgrass may be present. This plant community has developed due to improper grazing management and no fire.

Table 8. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	0%
Grass/grasslike foliar cover	0%
Forb foliar cover	0%
Non-vascular plants	0%
Biological crusts	0%
Litter	60-70%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%




Figure 6. Plant community growth curve (percent production by month). ID0805, B13 ARTRV . State 1.

Transition T1A
State 1 to 2

State 1 Phase C & D to State 2. Develops through improper grazing management and frequent fire. This site has crossed the threshold. It is economically impractical to return this state to State 1 with accelerated practices.

Transition T1B
State 1 to 3

State 1, Phase B to 3. Develops with no fire and improper grazing management from a juniper invaded phase of State 1. This site has crossed the threshold. It is economically impractical to return this state to State 1 with

accelerated practices.

Transition T2A

State 2 to 4

State 2 to unknown site. Excessive soil loss and changes in the hydrologic cycle caused by improper grazing management and no fire or frequent fire cause this state to cross the threshold and retrogress to a new site with reduced potential. It is economically impractical to return this state to State 1 with accelerated practices.

Transition T3A

State 3 to 4

State 3 to unknown site. Continued improper grazing management or the lack of fire cause this state to cross the threshold and retrogress to a new site with reduced potential due to significant soil loss and changes in hydrology. It is economically impractical to return this state to State 1 with accelerated practices

Additional community tables

Animal community

Wildlife Interpretations.

Animal Community – Wildlife Interpretations

This rangeland ecological site provides diverse habitat for many native wildlife species. The plant community exhibits a diverse mixture of forbs throughout the growing season offering excellent habitat for invertebrates. Mule deer, pronghorn antelope, and elk may utilize the site at different times of the year. The site provides seasonal habitat for resident and migratory animals including western toad, shrews, bats, ground squirrels, mice, coyote, red fox, badger, Ferruginous hawk, and prairie falcon. Area sensitive bird species include Brewer's sparrow, sage thrasher, sage sparrow, and sage-grouse. Water features are sparse provided by seasonal runoff, artificial water catchments, and springs.

State 1 Phase 1.1 – Mountain Big Sagebrush/ Idaho Fescue/ Bluebunch Wheatgrass/ Sandberg Bluegrass Reference Plant Community (RPC): This plant community provides a diversity of grasses, forbs, and shrubs used by native insect communities that assist in pollination. An extensive array of forbs is represented throughout the growing season leading to a diverse insect community. Many avian and mammal species utilize this habitat based on the availability of invertebrate prey species. The reptile and amphibian community is represented by common sagebrush lizard, western rattlesnake, Great Basin spadefoot toad, western toad, and northern leopard frog. Amphibians are associated with springs and isolated water bodies adjacent to this plant community. Development of spring sites that collect all available water would exclude amphibian use on these sites. Native shrub-steppe obligate bird species utilizing the habitat include the Brewer's sparrow, sage sparrow, sage grouse, and sage thrasher. Sage-grouse habitat (leks, nesting, brood-rearing, and winter) is provided by this plant community. The plant community provides seasonal food and cover for large mammals including mule deer, pronghorn, and elk. Antelope bitterbrush may be present in this plant community along with Idaho fescue and bluebunch wheatgrass which are all important forage species for these animals. A diverse small mammal population including golden-mantled ground squirrels, jackrabbits, deer mice, and Great Basin Kangaroo rat may utilize this plant community. The deer mouse is beneficial to this site as it is the principal vector for planting bitterbrush seed.

State 1 Phase 1.2- Mountain Big Sagebrush/ Utah Juniper/ Sandberg Bluegrass Plant Community: This plant community is the result of improper grazing management and no fire. An increase in canopy cover of sagebrush and juniper contributes to a sparse herbaceous understory. A reduced herbaceous understory results in lower diversity and numbers of insects. The reptile and amphibian community is represented by common sagebrush lizard, western rattlesnake, Great Basin spadefoot toad, western toad, and northern leopard frog. The reduced diversity of insects and understory cover may reduce quality of food and cover for reptile populations. As juniper increases, habitat for Brewer's sparrow, sage thrasher, and sage sparrow may increase. Remaining sagebrush provides brood-rearing, winter cover, and winter food habitat for sage-grouse but as juniper encroaches the quality of this habitat is severely reduced or eliminated. The plant community supports limited seasonal habitat for mule deer, elk, and pronghorn. As juniper encroaches the site will provide additional thermal cover for large mammals. A

small mammal population including golden-mantled ground squirrels, chipmunks, and yellow-bellied marmots may utilize this site.

State 1 Phase 1.3 – Bluebunch Wheatgrass/ Sandberg Bluegrass/ Bottlebrush Squirreltail/ Thickspike Wheatgrass Plant Community: This plant community is a result of recent wildfire, prescribed burning, or brush management. The plant community, dominated by herbaceous vegetation with little to no sagebrush or antelope bitterbrush would provide less vertical structure for animals. Patches of root sprouting shrubs (threetip sagebrush and rabbitbrushes) may be present to provide limited vertical structure for wildlife. Insect diversity would be reduced but a native forb plant community similar to that in State 1 Phase 1.1 would still support select pollinators. Habitat for common sagebrush lizard and western rattlesnake would be limited due to the loss of sagebrush. Amphibian habitat would be tied to permanent spring sites in the area. Development of spring sites that collected all available water would exclude the use of amphibians on these sites. The dominance of herbaceous vegetation with little sagebrush or antelope bitterbrush canopy cover would limit use of these areas as nesting habitat by Brewer's sparrow, sage sparrow, sage grouse, and sage thrasher. The dominant herbaceous vegetation improves habitat for grassland avian species (horned lark, savannah sparrow, vesper sparrow, and western meadowlark). Mule deer and elk use would be seasonal but the site would offer little thermal or young of year cover with the loss of shrubs. The populations of small mammals would be dominated by open grassland species. Large blocks of this plant community would fragment the reference plant community and reduce the quality of habitat for shrub-steppe obligate animal species.

State 1 Phase 1.4 – Mountain Big Sagebrush / Sandberg Bluegrass/ Thickspike Wheatgrass Plant Community: This plant community is the result of improper grazing management and no fire. An increase in canopy cover of sagebrush contributes to a sparse herbaceous understory. Grasses, forbs, and shrubs are used by native insects that assist in pollination but the reduced herbaceous understory results in lower diversity of insects. The reptile and amphibian community is represented by common sagebrush lizard, western rattlesnake, Great Basin spadefoot toad, western toad, and northern leopard frog. The reduced diversity of insects may reduce reptile diversity and populations. Reduced herbaceous understory is a key factor in limiting the use of this plant community by avian species. Key shrub-steppe obligate birds include Brewer's sparrow, sage sparrow, sage thrasher, and sage-grouse. Habitat (brood-rearing and nesting cover) quality for sage-grouse is reduced due to a less diverse herbaceous plant community. Winter habitat (cover and food) for sage-grouse is provided. The reduced vigor of understory vegetation provides a shorter forage season for mule deer, elk, and pronghorn. Young of year cover would be provided for deer. Small mammal diversity and populations would be similar to that in State 1 Phase 1.1.

State 2 – Sandberg/ Annuals Plant Community:

This community has developed due to continued improper grazing management and frequent fire. The plant community does not support a diverse insect community. The reduced forb and shrub component in the plant community would support a very limited population of pollinators. Most reptilian species are not supported with food, water, or cover. This plant community does not support the habitat requirements for sage thrasher, Brewer's sparrow, sage-grouse, or sage sparrow. Diversity of grassland avian species is reduced due to poor cover and available food. Birds of prey including hawks and falcons may range throughout these areas looking for prey species. Large mammals may utilize the herbaceous vegetation in spring and summer when herbaceous vegetation is more palatable. The diversity and populations of small mammals would be dominated by open grassland species. Large blocks of this plant community would fragment the reference plant community and reduce the quality of habitat for shrub-steppe obligate animal species.

State 3 - Utah Juniper/ Sandberg Bluegrass/ Annuals Plant Community: This site has developed due to improper grazing management and no fire. The loss of native understory vegetation will reduce insect diversity on the site. Habitat for sagebrush lizard, Great Basin spadefoot toad, and western toad would be limited due to the loss of brush cover. This plant community does not support the habitat requirements for sage-grouse. Birds using this site as resident or migratory habitat include Juniper titmouse, western bluebird, and Virginia's warbler. The Juniper titmouse relies heavily on juniper seeds for winter food. Hunting success by raptors may decrease due to a heavy overstory of juniper. The plant community supports limited seasonal habitat for mule deer and elk in spring and fall. As juniper encroaches the site will provide additional thermal cover for large mammals.

Grazing Interpretations.

This site is most suitable for livestock grazing in the spring, early summer, and fall.

Estimated initial stocking rate will be determined with the landowner or decision-maker. They will be based on the inventory which includes species composition, similarity index, production, past use history, season of use, and seasonal preference. Calculations used to determine estimated initial stocking rate will be based on forage preference ratings.

Hydrological functions

The soils in this site are in hydrologic group B. When the hydrologic conditions of the vegetative cover are good, natural erosion hazard is slight.

Recreational uses

This site has very little recreational value. It does have spring and early summer blooming flowers which offer some aesthetic values. The site has some value for hunting of small game, big game and upland game birds.

Wood products

none.

Other products

none.

Other information

Field Offices

American Falls, ID
Blackfoot, ID
Burley, ID
Driggs, ID
Idaho Falls, ID
Malad, ID
Pocatello, ID
Rexburg, ID
Soda Springs, ID
St. Anthony, ID

Revision Notes: "Previously Approved" Provisional
This Provisional ecological site concept has passed Quality Control (QC) and Quality Assurance (QA) to ensure that the site meets the 2014 NESH standards for a Provisional ecological site description. This is an updated "Previously Approved" ESD that represents a first-generation tier of documentation that, prior to the release of the 2014 National Ecological Site Handbook (NESH), met all requirements as an "Approved" ESD as laid out in the 1997 (rev.1, 2003) National Range and Pasture Handbook (NRPH). The document fully described the Reference State and Community Phase in the State-and-Transition model. All other alternative states are at least described in narrative form. The "Previously Approved" ESD has been field-tested for a minimum of five years and is a proven functional document for conservation planning. The "Previously Approved" ESD does not contain all tabular and narrative entries as required in the current "Approved" level of documentation, but it is expected that the "Previously Approved" ESD will continue refinement toward an "Approved" status.

Site Development and Testing Plan:

Future work, as described in a Project Plan, is necessary to validate the information in this Provisional Ecological Site Description. This will include field activities to collect low-, medium-, and high-intensity sampling, soil correlations, and analysis of that data. Annual field reviews should be done by soil scientists and vegetation specialists. The final field review, peer review, quality control, and quality assurance reviews of the ESD will be required to produce the final document.

Inventory data references

Information presented here has been derived from NRCS clipping and other inventory data. Also, field knowledge of range-trained personnel was used. Those involved in developing this site description include:

Dave Franzen, co-owner, Intermountain Rangeland Consultants, LLC
Jacy Gibbs, co-owner, Intermountain Rangeland Consultants, LLC
Jim Cornwell, Range Management Specialist, IASCD
Brendan Brazee, State Rangeland Management Specialist, NRCS, Idaho
Lee Brooks, Range Management Specialist, IASCD
Kristen May, Resource Soil Scientist, NRCS, Idaho

Other references

Hironaka, M., M.A. Fosberg, A. H. Winward. 1983. Sagebrush-Grass Habitat Types of Southern Idaho. University of Idaho, Moscow, Idaho. Bulletin Number "35".
Petersen, S.L., 2004. A Landscape-Scale Assessment of Plant Communities, Hydrologic Processes, and State-and-Transition Theory in a Western Juniper Dominated Ecosystem. PhD Dissertation. Oregon State University, Corvallis, Oregon.

USDA Forest Service, Rocky Mountain Research Station. 2004. Restoring Western Ranges and Wildlands. General Technical Report RMRS-GTR-136-vols. 1-3.
USDA, NRCS.2001. The PLANTS Database, Version 3.1 (<http://plants.usda.gov>). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.
USDA, Forest Service, Fire Effects Information Database. 2004. www.fs.fed.us/database/feis
USDI Bureau of Land Management, US Geological Survey; USDA Natural Resources Conservation Service, Agricultural Research Service; Interpreting Indicators of Rangeland Health. Technical Reference 1734-6; Version 4-2005.

Contributors

D. Edgerton

Approval

Kendra Moseley, 9/23/2020

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Dave Franzen and Jacy Gibbs Intermountain Range Consultants 17700 Fargo Rd. Wilder, ID 83676
--------------------------	--

Contact for lead author	Brendan Brazee, State Rangeland Management Specialist USDA-NRCS 9173 W. Barnes Drive, Suite C, Boise, ID 83709
Date	05/08/2008
Approved by	Kendra Moseley
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- Number and extent of rills:** rills rarely occur on this site. If rills are present they are likely to occur on slopes greater than 15 percent and immediately following wildfire. Rills are most likely to occur on soils with surface textures of silt loam and clay loam. Surface stones reduce rill formation by breaking up water flow patterns.

- Presence of water flow patterns:** water-flow patterns rarely occur on this site. If rills are present they are likely to occur on slopes greater than 15 percent and immediately following wildfire. Rills are most likely to occur on soils with surface textures of silt loam and clay loam. Surface stones reduce rill formation by breaking up water flow patterns.

- Number and height of erosional pedestals or terracettes:** both are rare on this site. In areas where slopes approach 15 percent and where flow patterns and/or rills are present, a few pedestals may be expected.

- Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** data is not available. On sites in mid-seral status bare ground may range from 30-50 percent.

- Number of gullies and erosion associated with gullies:** none.

- Extent of wind scoured, blowouts and/or depositional areas:** usually not present. Immediately following wildfire some soil movement may occur on lighter textured soils and on small areas without surface stones. Surface stones effectively stops wind erosion.

- Amount of litter movement (describe size and distance expected to travel):** fine litter in the interspaces may move up to 2 feet following a significant run-off event. Coarse litter generally does not move. Surface stones catch litter.

- Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** values should range from 4 to 6 but needs to be tested.

- Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** structure ranges from weak and moderate very fine, fine and medium granular to weak very thin, thin and very thick platy. Soil organic matter (SOM) ranges from 1 to 6 percent. Surface color can be very dark brown too very dark grayish brown to dark brown. The A or A1 horizon is typically 2 to 7 inches thick.

-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** bunchgrasses, especially deep-rooted perennials, slow run-off and increase infiltration. Tall shrubs accumulate snow in the interspaces. Surface stones slow water run-off and increase infiltration in the interspaces.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** not present.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: cool season deep-rooted perennial bunchgrasses
- Sub-dominant: tall shrubs
- Other: perennial forbs
- Additional: shallow rooted bunchgrasses
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** mountain big sagebrush and antelope bitterbrush will become decadent in the absence of fire and ungulate grazing. Grass and forb mortality will occur as tall shrubs increase.
-
14. **Average percent litter cover (%) and depth (in):** additional litter cover data is needed but is expected to be 20-25 percent to a depth of 0.1 inches. Under mature shrubs litter is >0.5 inches deep and is 90-100 percent ground cover.
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** is 1000 pounds per acre (1120 kilograms per hectare) in a year with normal temperatures and precipitation. Perennial grasses produce 60-70 percent of the total production, forbs 5-15 percent and shrubs 20-30 percent.
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** includes whitetop, rush skeletonweed, musk and scotch thistle, leafy spurge, Utah juniper, and diffuse and spotted knapweed. Cheatgrass may invade at lower elevations on the site.
-
17. **Perennial plant reproductive capability:** all functional groups have the potential to reproduce in most years.
-

