

## **Ecological site R013XY024ID Loamy 22+ PZ ARTRV/FEID-BRMA4**

Last updated: 9/23/2020  
 Accessed: 04/20/2024

### **Rangeland health reference sheet**

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Dave Franzen and Jacy Gibbs Intermountain Range Consultants 17700 Fargo Rd. Wilder, ID 83676
Contact for lead author	Brendan Brazee, State Rangeland Management Specialist USDA-NRCS 9173 W. Barnes Drive, Suite C, Boise, ID 83709
Date	05/02/2008
Approved by	Kendra Moseley
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### **Indicators**

1. **Number and extent of rills:** rills rarely occur on this site. If rills are present they are most likely to occur immediately following wildfire. Rills are most likely to occur on soils with surface textures of silt loam.

---

2. **Presence of water flow patterns:** water-flow patterns rarely occur on this site. When they occur they are short, disrupted by cool season perennial grasses and tall shrubs and are not extensive.

---

3. **Number and height of erosional pedestals or terracettes:** both are rare on this site. In areas where flow patterns and /or rills are present, a few pedestals may be expected.

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** data not available. On sites in mid-seral status, bare ground may range from 20-40 percent.

---

5. **Number of gullies and erosion associated with gullies:** none.

---

6. **Extent of wind scoured, blowouts and/or depositional areas:** usually does not occur.

---

7. **Amount of litter movement (describe size and distance expected to travel):** fine litter in the interspaces may move up to 2 feet following a significant run-off event. Terracettes and bunchgrass can trap fine litter. Coarse litter generally does not move.

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** values should range from 4 to 6 but needs to be tested.

---

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** structure ranges from weak fine granular to weak and moderate fine subangular blocky. Soil organic matter (SOM) generally ranges from 2 to 5 percent, but in some cases have a range of 60 to 95 percent. Surface color is generally very dark brown to very dark grayish brown. The A or A1 horizon is typically 4 to 12 inches thick.

---

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** bunchgrasses, especially deep-rooted perennials, slow run-off and increase infiltration. Shrubs accumulate snow in the interspaces.

---

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** not present.

---

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: cool season deep rooted perennial bunchgrasses

Sub-dominant: perennial forbs

Other: tall shrubs

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** mountain big sagebrush will become decadent in the absence of fire and ungulate grazing. Grass and forb mortality will occur as tall shrubs increase.

---

14. **Average percent litter cover (%) and depth ( in):** annual litter cover in the interspaces will be 5-10 percent to a depth of <0.2. Under the mature shrubs litter is greater than 0.5 inches. Fine litter can accumulate on the terracettes.

---

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-**

**production):** is 2600 lbs. per acre in a year with normal precipitation and temperatures. Perennial grasses produce 70-80 percent of the total, forbs 10-20 percent, and shrubs 5-15 percent.

---

16. **Potential invasive (including noxious) species (native and non-native).** List species which **BOTH** characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is **NOT** expected in the reference state for the ecological site: includes cheatgrass at lower elevations. Medusahead, bulbous bluegrass, rush skeletonweed, scotch thistle, spotted and diffuse knapweed, and leafy spurge. Other thistles.
- 

17. **Perennial plant reproductive capability:** all functional groups have the potential to reproduce in most years.
-