

Ecological site R013XY028ID Shallow Sand 12-16 PZ ARTRV/PSSPS

Last updated: 9/23/2020
Accessed: 05/05/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 013X–Eastern Idaho Plateaus

013X-Eastern Idaho Plateaus

Precipitation or Climate Zone: 12-16" P.Z.

<https://soils.usda.gov/survey/geography/mlra/index.html>

Classification relationships

Artemisia tridentata/*Agropyron spicatum* HT in "Hironaka, M., M.A. Fosberg, A. H. Winward. 1983. Sagebrush-Grass Habitat Types of Southern Idaho. University of Idaho, Moscow, Idaho. Bulletin Number "35".

Land Resource Unit: B (Northwestern Wheat and Range)

MLRA: 13 (Eastern Idaho Plateaus)

EPA EcoRegion: Level III (Middle Rockies)

Ecological site concept

Site does not receive any additional water.

Soils are:

not saline or saline-sodic.

shallow, very shallow with < 3% stone (10-25") and boulder (>25") cover. Soils skeletal within 20" of soil surface with gravel and cobble sized rock.

not strongly or violently effervescent in surface mineral 10".

textures usually range are sand, and fine sandy loam in surface mineral 4".

Slope is < 30%.

Clay content is = <10% in surface mineral 4".

Site does not have an argillic horizon with > 35% clay

Associated sites

R013XY001ID	Loamy 12-16 PZ
R013XY002ID	Stony Loam 13-16 PZ ARTRV/PSSPS
R013XY004ID	Shallow Gravelly 12-16 PZ ARTRV/PSSPS
R013XY008ID	Steep South Slopes 12-16 PZ ARTRV/PSSPS
R013XY012ID	Gravelly South Slope 12-16 PZ ARTRV/PSSPS

Similar sites

R013XY027ID	Sand 12-16 PZ PUTR2/HECOC8
-------------	-----------------------------------

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Artemisia tridentata ssp. vaseyana</i>
Herbaceous	(1) <i>Pseudoroegneria spicata</i>

Physiographic features

This site occurs on basalt plains in areas with sand dunes. Associated within the site are areas of rock outcrop and pockets of deep soils. Slopes range from 1 to 6 percent on all aspects. Elevations range from 4900 to 5500 feet (1500 to 1675 meters).

Table 2. Representative physiographic features

Landforms	(1) Lava plain
Flooding frequency	None
Elevation	1,494–1,676 m
Slope	1–6%
Water table depth	152 cm
Aspect	Aspect is not a significant factor

Climatic features

MLRA 13, the Eastern Idaho Plateaus, is part of the Northwestern Wheat and Range Region. Its elevation ranges from 4209 to 9331 feet above sea level, with an average elevation of 5787 feet. The average annual precipitation is 16.41 inches, with a range of 13.56 to 18.75 inches, based on ten long term climate stations located throughout the MLRA. A spike in precipitation amount often occurs in late spring, usually in May.

Temperatures vary widely in the MLRA throughout the year. A maximum temperature of 103° Fahrenheit occurred at the McCammon climate station (# 105716; elevation 4770 feet), while a minimum of -41° was recorded at the Kilgore station (#104908). At all stations temperatures throughout the year are usually below the national average. Kilgore also recorded the greatest annual snowfall amount of 217 inches. The average temperature is 41.4 degrees F. with an average high of 55.3 degrees and an average low of 27.5 degrees.

The frost-free period ranges from 64 to 90 days, while the freeze-free period can be 98 to 123 days.

Table 3. Representative climatic features

Frost-free period (average)	90 days
Freeze-free period (average)	123 days
Precipitation total (average)	483 mm




Figure 1. Monthly precipitation range




Figure 2. Monthly average minimum and maximum temperature

Influencing water features

This site is not influenced by adjacent wetlands, streams, or run on.

Soil features

This site contains well to excessively drained soils that are very shallow to shallow to bedrock. They are on basalt plains that formed from sandy eolian deposits. Textures are dominantly sand and very cobbly fine sandy loam. Coarse fragments are present and can exceed 35 percent by volume in the soil profile. Permeability is moderately rapid to very rapid and a very low to low runoff. The available water holding capacity is very low, however moisture does penetrate the fractures in the bedrock. . The soils are characterized by a xeric soil moisture regime and a frigid soil temperature regime.

Soil Series Correlated to this Ecological Site

Nayrib
Sandcreek

Table 4. Representative soil features

Surface texture	(1) Very cobbly sand (2) Fine sandy loam
Drainage class	Well drained to excessively drained
Permeability class	Moderately rapid to very rapid
Soil depth	15–51 cm

Surface fragment cover <=3"	0–20%
Surface fragment cover >3"	0–25%
Available water capacity (0-101.6cm)	1.52–3.56 cm
Soil reaction (1:1 water) (0-101.6cm)	6.6–7.8
Subsurface fragment volume <=3" (Depth not specified)	5–40%
Subsurface fragment volume >3" (Depth not specified)	0–5%

Ecological dynamics

The dominant visual aspect is mountain big sagebrush and antelope bitterbrush in the overstory with bluebunch wheatgrass in the understory. Composition by weight is approximately 40 to 50 percent grasses, 10 to 20 percent forbs, and 40 to 50 percent shrubs.

During the last few thousand years, this site has evolved in a semi-arid climate characterized by dry summers and cold, wet winters. Herbivory has historically occurred on this site at low levels of utilization. Herbivores include mule deer, Rocky Mountain elk, lagomorphs, and small rodents.

Fire has historically occurred on the site at intervals of 80 to 100 years.

The Historic Climax Plant Community (HCPC), the Reference State (State 1), moves through many phases depending on the natural and man-made forces that impact the community over time. State 1, described later, indicates some of these phases. The Reference Plant Community Phase is Phase A. This plant community is dominated by bluebunch wheatgrass in the understory and mountain big sagebrush and antelope bitterbrush in the overstory. Subdominant species include Idaho fescue, Sandberg bluegrass, prairie junegrass, arrowleaf balsamroot, lupine and buckwheat. There are a large variety of other grasses, forbs and shrubs in the plant community in minor amounts. The plant species composition of Phase A is listed later under "Reference Plant Community Phase Plant Species Composition".

Total annual production is 400 pounds per acre (450 kilograms per hectare) in a normal year. Production in a favorable year is 600 pounds per acre (670 kilograms per hectare). Production in an unfavorable year is 200 pounds per acre (224 kilograms per hectare). Structurally, cool season deep rooted perennial bunchgrasses and shrubs are about equal in the plant community followed by perennial forbs while shallow rooted bunchgrasses are subdominant.

FUNCTION:

This site is suited for livestock grazing in late spring, early summer, and fall. The site also provides big game habitat in the spring, summer, and fall. The site can be winter range for big game in moderate winters. This site offers minimal recreational or aesthetic values.

This site is susceptible to degradation due to the gentle slopes, ease of access, and instability of sandy surface textures. Infiltration is good where the community is in mid to late seral status. The site has moderately low runoff potential. Snow accumulates on the site due to elevation and presence of tall shrubs.

Impacts on the Plant Community.

Influence of fire:

In the absence of normal fire frequency, mountain big sagebrush and antelope bitterbrush can gradually increase on the site. Utah juniper can invade the site if a seed source is in the proximity. Grasses and forbs decrease as

shrubs increase. With the continued absence of fire, mountain big sagebrush or juniper can displace most of the primary understory species. See "Influence of juniper invasion" below.

When fires become more frequent than historic levels (80-100 years), mountain big sagebrush and bitterbrush are reduced significantly. Rabbitbrush can increase slightly. With continued short fire frequency, mountain big sagebrush and bitterbrush can be completely eliminated along with many of the desirable understory species such as bluebunch wheatgrass and Idaho fescue. These species may be replaced by Sandberg bluegrass along with a variety of annual and perennial forbs including noxious and invasive plants. Cheatgrass will invade the site. These fine fuels will increase the fire frequency.

Influence of improper grazing management:

Season-long grazing and/or excessive utilization can be very detrimental to this site. This type of management leads to reduced vigor of the bunchgrasses. With reduced vigor, recruitment of these species declines. As these species decline, the plant community becomes susceptible to juniper invasion, an increase in mountain big sagebrush, and noxious and invasive plants.

Continued improper grazing management influences fire frequency by increasing fine fuels. As cheatgrass increases and becomes co-dominant with other annuals, fires become more frequent, particularly at lower elevations.

Proper grazing management that addresses frequency, duration, and intensity of grazing can also keep fine fuels from developing, thereby reducing fire frequency. This can lead to gradual increases in mountain big sagebrush. An increase in tall shrubs generally leads to an increase in juniper by providing bird perches and "nursery" sites for juniper establishment. A planned grazing system can be developed to intentionally accumulate fine fuels in preparation for a prescribed burn. Due to the shrub species on the site, any brush management efforts need to be very carefully evaluated. Antelope bitterbrush is very important as a browse species for wildlife and needs to be protected with any brush control practices applied. A reduction of shrubs in areas without a suitable understory of perennial grasses can lead to an increase in fine fuels which will lead to a more frequent fire regime. Loss of shrub species on this site can have very negative impacts on wildlife.

Weather influences:

Above normal precipitation in April, May, and June can dramatically increase total annual production of the plant community. These weather patterns can also increase viable seed production of desirable species to provide for recruitment. Likewise, below normal precipitation during these spring months can significantly reduce total annual production and be detrimental to viable seed production. Overall plant composition is normally not affected when perennials have good vigor.

Below normal temperatures in the spring can have an adverse impact on total production regardless of the precipitation. An early, hard freeze can occasionally kill some plants.

Prolonged drought adversely affects this plant community in several ways. Vigor, recruitment, and production are usually reduced. Mortality can occur. Prolonged drought can lead to a reduction in fire frequency.

Influence of Insects and disease:

Outbreaks can affect vegetation health. Bitterbrush can be severely affected by the western tent caterpillar (*Malacosoma fragilis*). Two consecutive years of defoliation by the tent caterpillar can cause mortality in bitterbrush. Mormon cricket and grasshopper outbreaks occur periodically. Outbreaks seldom cause plant mortality since defoliation of the plant occurs only once during the year of the outbreak. Snow mold can occasionally cause reduced vigor in mountain big sagebrush, but it is a short-term effect.

Influence of noxious and invasive plants:

Many of these species add to the fine-fuel component and lead to increased fire frequency. Many of the annual and perennial invasive species compete with desirable plants for moisture and nutrients. The result is reduced production and change in composition of the understory.

Influence of wildlife:

Big game animals use this site in the spring, early summer, and fall and in moderate winters. Their numbers are seldom high enough to adversely affect the plant community. Herbivory can be detrimental to bitterbrush when livestock grazing and browsing by big game occurs at the same time and season. This will occur when both kinds of animal are using the plant in the late summer or fall. The adverse impact is excessive use of the current year's leader growth. The deer mouse is beneficial to this site as it is the principal vector for planting bitterbrush seed.

Watershed:

Decreased infiltration and increased runoff occur with the invasion of Utah juniper or an increase in mountain big sagebrush. Juniper invasion can be triggered by lack of fire, improper grazing management, and prolonged drought. Desired understory species can be reduced. This composition change can affect nutrient and water cycles. Increased runoff also causes sheet and rill erosion. Abnormally short fire frequency also gives the same results, but to a lesser degree. The long term effect is a transition to a different state.

Influence of juniper invasion:

The following discussion deals with both western juniper and Utah juniper.

In plant communities that are invaded by juniper, the species has a competitive advantage for the following reasons:

- Juniper is very drought tolerant.
- It has the ability to extract soil moisture from a wide range of soil depths.
- Juniper has high evapo-transpiration rates.
- The species intercepts rain and snow before it reaches the soil surface.
- It has the ability to grow as long as there is soil moisture and the temperature is above freezing.
- Juniper has a relatively rapid growth rate and is long-lived. It can readily over-top shade intolerant species which leads to mortality.
- Nutrient cycling is reduced.
- As the canopy closes, juniper gains control of energy capture.

As juniper extracts water, other plants are unable to acquire sufficient water and nutrients to sustain growth and reproduction, thus reducing cover and biomass in the interspaces. After the canopy closes, there is sufficient soil moisture available for shallow-rooted, shade tolerant species to persist directly under the tree.

The following hydrologic impacts occur on sites invaded by juniper:

- Infiltration in the interspaces is reduced.
- Run-off increases resulting in increased sheet and rill erosion with elevated sediment loads.
- Soil temperatures increase in the interspaces which results in accelerated drying of the soil surface.
- Increased bare ground in the interspaces.
- Soil moisture storage is reduced.

As bare ground and interconnectiveness of bare ground increases, flow rates are accelerated (reduction of flow sinuosity) and run-off out of the area increases.

Degradation of these systems can result in the formation of a feedback cycle in which greater juniper cover and density results in greater plant and soil disturbance between the canopies.

In summary, a closed juniper community takes control of the following ecological processes: (1) hydrology, (2) energy capture, and (3) nutrient cycling. The changes are primarily driven by the hydrologic processes. The development of a closed juniper canopy always results in a transition across the threshold to a different state. Generally, when juniper canopy cover nears 20%, the plant community is approaching the threshold.

Plant Community and Sequence:

Transition pathways between common vegetation states and phases:

State 1.

Phase A to B. Develops in the absence of fire. No Utah juniper seed source in the proximity.

Phase A to C. Usually results from improper grazing management and absence of fire. A juniper seed source is present.

Phase A to D. Results from one or more fires.

Phase A to E. Develops in the absence of fire. A juniper seed source is present.

Phase A to F. Results from improper grazing management and absence of fire. No juniper seed source is present.

Phase B to A. Results from prescribed grazing management.

Phase C to A. Develops with prescribed grazing management and prescribed burning or fire.

Phase D to A. Usually results from prescribed grazing management and no fire.

Phase E to A. Develops from prescribed grazing management and prescribed burning or brush management.

Phase F to A. Results from prescribed grazing management, no fire or brush management.

Phase B to D. This develops from prescribed burning or fire.

Phase C to D. This develops from prescribed burning or fire.

Phase E to D. Results from prescribed burning or fire.

Phase F to D. Results from prescribed burning or fire.

State 1 Phase D to State 2, Phase B. Develops through improper grazing management and frequent fire. This site has crossed the threshold. It is usually uneconomical to return this community to State 1 through accelerated practices.

State 1 Phase F to State 2 Phase A. Develops through improper grazing management with no fire. It has crossed the threshold. This site has crossed the threshold. It is usually uneconomical to return this community to State 1 through accelerated practices.

State 1 Phase C or E to State 3. Results from improper grazing management and lack of fire. It has crossed the threshold. It is usually uneconomical to return this community to State 1 through accelerated practices.

State 2 Phase A to State 2 Phase B. Results from improper grazing management and fire. This site has crossed the threshold. It is usually uneconomical to return this community to State 1 through accelerated practices.

State 2 Phase B to State 2 Phase A. Results from no fire. This site has crossed the threshold. It is usually uneconomical to return this community to State 1 through accelerated practices.


State 2 to unknown site. Excessive soil loss and changes in the hydrologic cycle caused by improper grazing management and/or frequent fire cause this state to cross the threshold and to retrogress to a new site with reduced potential. It is usually uneconomical to return this community to State 1 through accelerated practices.

State 3 to unknown site. Continued lack of fire and improper grazing management cause this state to cross the threshold and to retrogress to a new site with reduced potential due to significant soil loss and changes in hydrology. It is usually uneconomical to return this community to State 1 through accelerated practices.

Practice Limitations:

Slight limitations exist on this site for implementing vegetative management practices. Opportunities for rangeland seeding are severely limited due to shallow soil and associated rock outcrop areas. Moderate to severe limitations exist for brush management due to surrounding rock outcrops and the species involved.

State and transition model



State 1
State 1

Community 1.1
State 1 Phase A

Reference Plant Community Phase. This plant community is dominated by bluebunch wheatgrass in the understory and mountain big sagebrush and antelope bitterbrush in the overstory. Subdominant species include Idaho fescue, Sandberg bluegrass, prairie junegrass, arrowleaf balsamroot, lupine, and buckwheat. There are a large variety of other grasses, forbs and shrubs in the plant community in minor amounts. Natural fire frequency is 20-50 years.

Table 5. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	0%
Grass/grasslike foliar cover	0%
Forb foliar cover	0%
Non-vascular plants	0%
Biological crusts	0%
Litter	15-30%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%




Figure 3. Plant community growth curve (percent production by month). ID0805, B13 ARTRV . State 1.

Community 1.2 State 1, Phase B

This plant community is dominated by bitterbrush and mountain big sagebrush. Bluebunch wheatgrass is the dominant species in the understory with Idaho fescue and Sandberg bluegrass subdominant. Antelope bitterbrush has increased. A wide variety of other grasses and forbs are present in reduced amounts. No Utah juniper seed source is present. This state has developed due to fire frequency being much longer than normal.

Community 1.3 State 1, Phase C

This plant community is dominated mountain big sagebrush in the overstory with small Utah juniper trees or saplings invading. Bluebunch wheatgrass has decreased in the community but is still dominant. Sandberg bluegrass, thickspike wheatgrass and bottlebrush squirreltail have increased along with forbs. All bunchgrass species are reduced in amounts and have low vigor. Antelope bitterbrush is present but in reduced vigor and may be hedged. Cheatgrass and Kentucky bluegrass may have invaded the site. This state has developed due to improper grazing management and lack of fire. A Utah juniper seed source is in the proximity.

Community 1.4 State 1, Phase D

This plant community is dominated by bluebunch wheatgrass. Idaho fescue may have been killed due to fire. Sandberg bluegrass, bottlebrush squirreltail, sand dropseed, thickspike wheatgrass, and western wheatgrass have increased. Forbs are about the same as in Phase A. Mountain big sagebrush and antelope bitterbrush have been significantly reduced, but root sprouting shrubs such as rabbitbrush and horsebrush may be present. This plant community is a result of prescribed burning or wildfire.

Community 1.5

State 1, Phase E

This plant community is similar to the Phase A except that juniper seedlings and saplings are invading the site due to a lack of fire. A Utah juniper seed source is in the proximity. This state has developed due to the absence of fire.

Community 1.6

State 1, Phase F

This plant community is dominated by mountain big sagebrush in the overstory. Sandberg bluegrass has increased. Bluebunch wheatgrass and other bunchgrasses are present but in reduced amounts and typically in low vigor. Antelope bitterbrush is present but in reduced amounts and low vigor and it may be hedged. This state has developed due to improper grazing management and a lack of fire. No Utah juniper seed source is in the proximity.

Pathway A to B

Community 1.1 to 1.2

Develops in the absence of fire. No Utah juniper seed source in the proximity.

Pathway A to C

Community 1.1 to 1.3

Usually results from improper grazing management and absence of fire. A juniper seed source is present.

Pathway A to D

Community 1.1 to 1.4

Results from one or more fires.

Pathway A to E

Community 1.1 to 1.5

Develops in the absence of fire. A juniper seed source is present.

Pathway A to F

Community 1.1 to 1.6

Results from improper grazing management and absence of fire. No juniper seed source is present.

Pathway B to A

Community 1.2 to 1.1

Results from prescribed grazing management.

Pathway B to D

Community 1.2 to 1.4

This develops from prescribed burning or fire.

Pathway C to A

Community 1.3 to 1.1

Develops with prescribed grazing management and prescribed burning or fire.

Pathway C to D

Community 1.3 to 1.4

This develops from prescribed burning or fire.

Pathway D to A Community 1.4 to 1.1

Usually results from prescribed grazing management and no fire.

Pathway E to A Community 1.5 to 1.1

Develops from prescribed grazing management and prescribed burning or brush management.

Pathway E to D Community 1.5 to 1.4

Results from prescribed burning or fire.

Pathway F to A Community 1.6 to 1.1

Results from prescribed grazing management, no fire or brush management.

Pathway F to D Community 1.6 to 1.4

Results from prescribed burning or fire.

State 2 State 2

Community 2.1 State 2, Phase A

This plant community is dominated by mountain big sagebrush with Sandberg bluegrass and annuals in the interspaces. This state has developed due to improper grazing management and the absence of fire from phase F, State 1 or with no fire from phase B, State 2. Some soil loss has occurred. This site has crossed the threshold. It is usually uneconomical to return this community to State 1 through accelerated practices.

Community 2.2 State 2 Phase B

This plant community is dominated by Sandberg bluegrass, annuals and forbs. Root sprouting shrubs such as rabbitbrush and horsebrush can be present in small amounts. This state has developed due to improper grazing management and frequent fires from phase D, State 1 or with improper grazing management and fire from phase A, State 2. Some soil loss has occurred. This site has crossed the threshold. It is usually uneconomical to return this community to State 1 through accelerated practices.

Table 6. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	0%
Grass/grasslike foliar cover	0%
Forb foliar cover	0%
Non-vascular plants	0%

Biological crusts	0%
Litter	15-30%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%




Figure 4. Plant community growth curve (percent production by month). ID0805, B13 ARTRV . State 1.

Pathway A to B Community 2.1 to 2.2

Results from improper grazing management and fire. This site has crossed the threshold. It is usually uneconomical to return this community to State 1 through accelerated practices.

Pathway B to A Community 2.2 to 2.1

Results from no fire. This site has crossed the threshold. It is usually uneconomical to return this community to State 1 through accelerated practices.

State 3 State 3

This plant community is dominated by Utah juniper. Remnants of Bluebunch wheatgrass and some other bunchgrasses may be present, often under trees. Shallow-rooted grasses, such as Sandberg bluegrass and annuals can be found in the interspaces. Few shrubs are present. When shrub cover is below 10-15%, bare ground is above 25-30%, and juniper cover is greater than 20%, the site has crossed the threshold. Some soil loss has occurred. This state has developed with continued improper grazing management and in the absence of fire. It is usually uneconomical to return this community to State 1 through accelerated practices.

State 4 State 4

This plant community has gone over the threshold to a new site. Site potential has been reduced. Significant soil loss has occurred. Infiltration has been reduced and run-off has become more rapid. This state has developed due to continued improper grazing management and/or frequent fires from State 2 or the continued absence of fire and improper grazing management from State 3 where the site is dominated by juniper. This site will not return to State 1 or 2 because of significant soil loss.

Transition T1A

State 1 to 2

State 1 Phase D to State 2, Phase B. Develops through improper grazing management and frequent fire. This site has crossed the threshold. It is usually uneconomical to return this community to State 1 through accelerated practices. State 1 Phase F to State 2 Phase A. Develops through improper grazing management with no fire. It has crossed the threshold. This site has crossed the threshold. It is usually uneconomical to return this community to State 1 through accelerated practices.

Transition T1B

State 1 to 3

State 1 Phase C or E to State 3. Results from improper grazing management and lack of fire. It has crossed the threshold. It is usually uneconomical to return this community to State 1 through accelerated practices.

Transition T2A

State 2 to 4

Excessive soil loss and changes in the hydrologic cycle caused by improper grazing management and/or frequent fire cause this state to cross the threshold and to retrogress to a new site with reduced potential. It is usually uneconomical to return this community to State 1 through accelerated practices.

Transition T3A

State 3 to 4

Continued lack of fire and improper grazing management cause this state to cross the threshold and to retrogress to a new site with reduced potential due to significant soil loss and changes in hydrology. It is usually uneconomical to return this community to State 1 through accelerated practices.

Additional community tables

Animal community

Wildlife Interpretations.

Animal Community – Wildlife Interpretations

This rangeland ecological site provides diverse habitat for many native wildlife species. Forb and grass production is low but in combination with a diverse shrub community, provides excellent cover and browse for insects, reptiles, birds, and small to large mammals. Mule deer, elk, and sometimes pronghorn antelope may utilize the site at different times of the year. The rangeland provides seasonal habitat for resident and migratory animals including western toad, shrews, bats, ground squirrels, mice, coyote, red fox, badger, Ferruginous hawk, and prairie falcon. Area sensitive bird species include Brewer's sparrow, sage thrasher, sage sparrow, sharp-tailed grouse, and sage-grouse. Water features are sparse provided by seasonal runoff, artificial water catchments, and springs.

State 1 Phase 1.1 – Mountain Big Sagebrush/ Antelope Bitterbrush Bluebunch Wheatgrass/ Idaho Fescue/ Sandberg Bluegrass/ Prairie Junegrass Reference Plant Community (RPC): This plant community provides a diversity of grasses, forbs, and shrubs used by native insect communities that assist in pollination. A diverse array of forbs is represented throughout the growing season leading to a diverse insect community. Many avian and mammal species utilize this habitat based on the availability of invertebrate prey species. The reptile and amphibian community is represented by common sagebrush lizard, western rattlesnake, Great Basin spadefoot toad, western toad, and northern leopard frog. Amphibians are associated with springs and isolated water bodies adjacent to this plant community. Development of spring sites that collect all available water would exclude amphibian use on these sites. Native shrub-steppe obligate avian species utilizing the habitat include the Brewer's sparrow, sage sparrow, sage grouse, and sage thrasher. Sage-grouse habitat (leks, nesting, brood-rearing, and winter) is provided by this plant community. The plant community provides seasonal food and cover for large mammals including mule deer, pronghorn antelope, and elk. Antelope bitterbrush, bluebunch wheatgrass, and Idaho fescue are important forage species for these animals. Some areas (south and west facing slopes) may provide winter food for mule deer and elk. A diverse small mammal population including golden-mantled ground squirrels, jackrabbits, deer mice, and

Great Basin pocket mice may utilize this plant community. The deer mouse is beneficial to this site as it is the principal vector for planting bitterbrush seed.

State 1 Phase 1.2- Antelope Bitterbrush/ Mountain Big Sagebrush/ Bluebunch Wheatgrass/ Idaho Fescue/ Sandberg Bluegrass Plant Community: This phase has developed due to fire return intervals being much longer than normal. An increase in canopy cover of sagebrush and antelope bitterbrush contributes to a declining herbaceous understory. The reptile and amphibian community will be similar to the State 1 Phase 1.1 community. Sagebrush provides nesting, brood-rearing, winter cover, and winter food habitat for sage-grouse but as understory vegetation declines the quality of this habitat is reduced. The plant community supports spring and fall forage for mule deer, elk, and pronghorn. Quality of winter habitat for mule deer, elk, and pronghorn is similar to that in State 1 Phase 1.1. A diverse small mammal population including golden-mantled ground squirrels, jackrabbits, deer mice, and Great Basin pocket mice may utilize this plant community.

State 1 Phase 1.3 – Mountain Big Sagebrush/ Utah Juniper/ Bluebunch Wheatgrass/ Sandberg Bluegrass/ Thickspike Wheatgrass Plant Community: This plant community is the result of improper grazing management and no fire. An increase in canopy cover of sagebrush and juniper contributes to a sparse herbaceous understory. A reduced herbaceous understory results in lower diversity and numbers of insects. The reptile and amphibian community will be similar to the State 1 Phase 1.1 community, represented by common sagebrush lizard, western rattlesnake, Great Basin spadefoot toad, western toad, and northern leopard frog. The reduced diversity of insects and understory cover may reduce the quality of food and cover for reptile populations. As juniper increases, habitat for Brewer's sparrow, sage thrasher, and sage sparrow may increase. The remaining sagebrush provides brood-rearing, winter cover, and winter food habitat for sage-grouse but as juniper encroaches the quality of this habitat is severely reduced or eliminated. The plant community provides limited spring and fall forage for mule deer, elk, and pronghorn antelope due to the loss of understory vegetation. Juniper can provide winter habitat (cover and food) for mule deer. As juniper encroaches the site will provide additional thermal cover for large mammals. A diverse small mammal population including golden-mantled ground squirrels, jackrabbits, deer mice, and Great Basin pocket mice may utilize this plant community.

State 1 Phase 1.4 – Bluebunch Wheatgrass/ Sandberg Bluegrass/ Bottlebrush Squirreltail/ Sand Dropseed Plant Community: This plant community is a result of recent wildfire or prescribed burning. The plant community, dominated by herbaceous vegetation with little to no sagebrush or antelope bitterbrush would provide less vertical structure for animals. Patches of root sprouting shrubs (horsebrush and rabbitbrushes) may be present to provide limited vertical structure for wildlife in the future. Insect diversity would be reduced, but a native forb plant community similar to that in State 1 Phase 1.1 would still support select pollinators. The quality of food and cover habitat for reptiles would decline due to the loss of sagebrush and antelope bitterbrush. Amphibian habitat would be tied to permanent spring sites in the area. Development of spring sites that collect all available water would exclude the use of amphibians on these sites. The dominance of herbaceous vegetation with little sagebrush or antelope bitterbrush canopy cover would limit the use of these areas as nesting habitat by Brewer's sparrow, sage sparrow, sage grouse, and sage thrasher. The dominant herbaceous vegetation improves habitat for grassland avian species (horned lark, savannah sparrow, vesper sparrow, and western meadowlark). Sharp-tailed grouse may prefer this plant community over sites with more sagebrush canopy cover. Mule deer, pronghorn, and elk use would be seasonal (spring, summer, and fall) but the site would offer little thermal or young of year cover due to the loss of shrub cover. The diversity and populations of small mammals would be dominated by open grassland species. Large blocks of this plant community would fragment the reference plant community and reduce the quality of habitat for shrub-steppe obligate animal species.

State 1 Phase 1.5 – Mountain Big Sagebrush/ Bluebunch Wheatgrass/ Idaho Fescue/ Columbia Needlegrass/ Big Bluegrass Plant Community: This phase has developed due to the absence of fire. An increase in canopy cover of sagebrush contributes to a sparse herbaceous understory. The reptile and amphibian community is represented by common sagebrush lizard, western rattlesnake, Great Basin spadefoot toad, western toad, and northern leopard frog. Shrub-steppe obligate avian species include Brewer's sparrow, sage sparrow, sage thrasher, and sage-grouse. Quality of habitat (brood-rearing and nesting cover) for sage-grouse is reduced due to a less diverse herbaceous plant community and increase in junipers. Winter habitat (cover and food) for sage-grouse is provided but the quality is reduced or eliminated when junipers become dominant. The reduced vigor of understory vegetation provides a shorter forage season for mule deer, elk, and pronghorn antelope. Young of year cover would be provided for deer and pronghorn antelope. As juniper is established additional thermal cover is provided for mule deer and elk. Small mammal diversity and populations would be similar to those in State 1 Phase 1.1.

State 1 Phase 1.6 – Mountain Big Sagebrush/ Sandberg Bluegrass Plant Community: This phase has developed due to improper grazing management and a lack of fire. An increase in canopy cover of sagebrush and improper grazing results in a reduction of Idaho fescue and bluebunch wheatgrass. The reptile and amphibian community is represented by common sagebrush lizard, western rattlesnake, Great Basin spadefoot toad, western toad, and northern leopard frog. Shrub-steppe obligate avian species include Brewer's sparrow, sage sparrow, sage thrasher, and sage-grouse. Quality of habitat (brood-rearing and nesting cover) for sage-grouse is reduced due to poor vigor and a less diverse herbaceous plant community. Winter habitat (cover and food) for sage-grouse is provided. Young of year cover would be provided for deer and pronghorn antelope. Small mammal diversity and populations would be similar to those in State 1 Phase 1.1.

State 2 Phase 2.1 – Mountain Big Sagebrush/ Sandberg Bluegrass/ Annuals/ Plant Community: This plant community is the result of improper grazing management and no fire. An increase in canopy cover of sagebrush and improper grazing management results in a reduction of Idaho fescue and bluebunch wheatgrass. The reduced diversity of native herbaceous understory plants and increase in invasive plants results in a lower diversity of insects. The reptile and amphibian community is similar to that in Phase 1.6, State 1. The reduced diversity of insects may reduce reptile diversity and populations. Reduced herbaceous understory will lower quality of habitat for bird species. Shrub-steppe obligate avian species include Brewer's sparrow, sage sparrow, sage thrasher, and sage-grouse. Quality of habitat (brood-rearing and nesting cover) for sage-grouse is reduced due to poor vigor and a less diverse herbaceous plant community. Winter habitat (cover and food) for sage-grouse is provided. Young of year cover would be provided for deer and pronghorn antelope. Small mammal diversity and populations would be similar to those in Phase 1.1, State 1.

State 2 Phase 2.2 – Sandberg/ Annuals/ Forbs/ Green Rabbitbrush/ Mountain Snowberry Plant Community: This phase has developed due to improper grazing management and frequent fires from Phase 1.4, State 1 or with improper grazing management and fire from Phase 2.1, State 2. The reduced forb and shrub components in the plant community would support a very limited population of pollinators. Habitat for common sagebrush lizard, Great Basin spadefoot toad, and western toad would be limited due to the loss of sagebrush. The loss of sagebrush and antelope bitterbrush would severely reduce the quality of the habitat for sage thrasher, Brewer's sparrow, sage-grouse, and sage sparrow. Birds of prey including hawks and falcons may range throughout these areas looking for prey species. If managed properly, Kentucky bluegrass will provide excellent forage for elk and deer. The populations of small mammals would be dominated by open grassland species. Predator hunting success would increase due to the reduced quality of cover provided to small mammals. Large blocks of this plant community would fragment the reference plant community and reduce the quality of habitat for shrub-steppe obligate animal species.

State 3 – Utah Juniper/ Sandberg Bluegrass/ Annuals Plant Community: This state has developed due to improper grazing management and no fire. The loss of native forbs and understory vegetation will reduce insect diversity on the site. The lack of flowering plants reduces use by pollinators like butterflies and moths. Habitat for common sagebrush lizard, Great Basin spadefoot toad, and western toad would be limited due to the loss of sagebrush. This plant community does not support the habitat requirements for sage-grouse. Birds using this site as resident or migratory habitat include Juniper titmouse, western bluebird, waxwings, chipping sparrow, and Virginia's warbler. The Juniper titmouse relies heavily on juniper seeds for winter food. Hunting success by raptors may decrease due to a heavy overstory of juniper. As juniper encroaches the site will provide additional thermal cover for large mammals. The plant community provides winter food and cover for mule deer.

Grazing Interpretations.

This site is best suited for grazing by livestock in spring, early summer, and fall. Estimated initial stocking rate will be determined with the landowner or decision-maker. They will be based on the inventory which includes species composition, similarity index, production, past use history, season of use, and seasonal preference. Calculations used to determine estimated initial stocking rate will be based on forage preference ratings.

Hydrological functions

The soils in this site are in hydrologic group A. They have low runoff potential.

Recreational uses

This site offers minimal recreational or aesthetic values.

Wood products

none.

Other products

none.

Other information

Field Offices

American Falls

Blackfoot

Burley

Driggs

Idaho Falls

Malad

Pocatello

Rexburg

Soda Springs

St. Anthony

Revision Notes: "Previously Approved" Provisional

This Provisional ecological site concept has passed Quality Control (QC) and Quality Assurance (QA) to ensure that the site meets the 2014 NESH standards for a Provisional ecological site description. This is an updated "Previously Approved" ESD that represents a first-generation tier of documentation that, prior to the release of the 2014 National Ecological Site Handbook (NESH), met all requirements as an "Approved" ESD as laid out in the 1997 (rev.1, 2003) National Range and Pasture Handbook (NRPH). The document fully described the Reference State and Community Phase in the State-and-Transition model. All other alternative states are at least described in narrative form. The "Previously Approved" ESD has been field-tested for a minimum of five years and is a proven functional document for conservation planning. The "Previously Approved" ESD does not contain all tabular and narrative entries as required in the current "Approved" level of documentation, but it is expected that the "Previously Approved" ESD will continue refinement toward an "Approved" status.

Site Development and Testing Plan:

Future work, as described in a Project Plan, is necessary to validate the information in this Provisional Ecological Site Description. This will include field activities to collect low-, medium-, and high-intensity sampling, soil correlations, and analysis of that data. Annual field reviews should be done by soil scientists and vegetation specialists. The final field review, peer review, quality control, and quality assurance reviews of the ESD will be required to produce the final document.

Inventory data references

Information presented here has been derived from NRCS clipping and other inventory data. Also, field knowledge of range-trained personnel was used. Those involved in developing this site description include:

Dave Franzen, co-owner, Intermountain Rangeland Consultants, LLC

Jacy Gibbs, co-owner, Intermountain Rangeland Consultants, LLC

Jim Cornwell, Range Management Specialist, IASCD

Brendan Brazee, State Rangeland Management Specialist, NRCS, Idaho

Lee Brooks, Range Management Specialist, IASCD

Kristen May, Resource Soil Scientist, NRCS, Idaho

Other references

Hironaka, M., M.A. Fosberg, A. H. Winward. 1983. Sagebrush-Grass Habitat Types of Southern Idaho. University of Idaho, Moscow, Idaho. Bulletin Number "35".

Petersen, S.L., 2004. A Landscape-Scale Assessment of Plant Communities, Hydrologic Processes, and State-and-Transition Theory in a Western Juniper Dominated Ecosystem. PhD Dissertation. Oregon State University, Corvallis, Oregon.

USDA Forest Service, Rocky Mountain Research Station. 2004. Restoring Western Ranges and Wildlands. General Technical Report RMRS-GTR-136-vols. 1-3.

USDA, NRCS.2001. The PLANTS Database, Version 3.1 (<http://plants.usda.gov>). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

USDA, Forest Service, Fire Effects Information Database. 2004. www.fs.fed.us/database/feis

USDI Bureau of Land Management, US Geological Survey; USDA Natural Resources Conservation Service, Agricultural Research Service; Interpreting Indicators of Rangeland Health. Technical Reference 1734-6; Version 4-2005.

Approval

Kendra Moseley, 9/23/2020

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Dave Franzen and Jacy Gibbs Intermountain Range Consultants 17700 Fargo Rd. Wilder, ID 83676
Contact for lead author	Brendan Brazee, State Rangeland Management Specialist USDA-NRCS 9173 W. Barnes Drive, Suite C, Boise, ID 83709
Date	05/18/2008
Approved by	Kendra Moseley
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** rills are rare on this site. If rills are present, they are likely to occur immediately following wildfire and a high intensity convection storm. Sandy surface texture will limit rill development.

-
2. **Presence of water flow patterns:** water-flow patterns are rare on this site. They may occur immediately following a high intensity convention storm. If they occur, they are short and disrupted by cool season grasses and tall shrubs and are not extensive. Water infiltration is generally rapid for the site.
-
3. **Number and height of erosional pedestals or terracettes:** both are rare on this site.
-
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** data is not available for this site. On sites in mid-seral status, bare ground is expected to be about 50-60 percent. This site is naturally unstable due to sandy surface textures particularly following a wildfire.
-
5. **Number of gullies and erosion associated with gullies:** none.
-
6. **Extent of wind scoured, blowouts and/or depositional areas:** wind-scour and deposition areas can occur on this site particularly following a wildfire. Old depositions will be noticeable in the crowns of bunchgrasses and at the base of shrubs.
-
7. **Amount of litter movement (describe size and distance expected to travel):** fine litter in the interspaces may move up to 2 feet following a significant run-off event or further with wind. Coarse litter generally does not move.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** values average 1 to 2 but needs to be tested.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** structure is weak fine granular to single grain and massive. Soil organic matter (SOM) ranges from 2 to 4 percent. Surface color is usually very dark grayish brown. The A or A1 horizon is typically 4 to 8 inches thick.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** bunchgrasses, especially deep-rooted perennials, slow run-off and increase infiltration. Tall shrubs accumulate snow in the interspaces.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** not present.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: cool season deep-rooted perennial bunchgrasses

Sub-dominant: tall shrubs

Other: perennial forbs

Additional: shallow rooted bunchgrasses

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** mountain big sagebrush and antelope bitterbrush will become decadent in the absence of fire and ungulate grazing. Grass and forb mortality will occur as tall shrubs increase.
-

14. **Average percent litter cover (%) and depth (in):** additional litter cover data is needed but is expected to be 5-10 percent to a depth of <0.1 inches. Under mature shrubs litter is >0.5 inches deep and is 90-100 percent ground cover.
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** is 400 pounds per acre (450 kilograms per hectare) in a year with normal temperatures and precipitation. Perennial grasses produce 40-50 percent of the total production, forbs 10-20 percent and shrubs 40-50 percent.
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** includes cheatgrass, leafy spurge, dalmatian toadflax, rush skeletonweed, musk and scotch thistle, and diffuse, Russian, and spotted knapweed, Russian thistle, and mustard.
-

17. **Perennial plant reproductive capability:** all functional groups have the potential to reproduce in most years.
-