

Ecological site R022AX101CA Frigid Anastomosed System

Accessed: 04/19/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 022A–Sierra Nevada and Tehachapi Mountains

Major Land Resource Area 22A, Sierra Nevada Mountains, is located predominantly in California and a small section of western Nevada. The area lies completely within the Sierra Nevada Section of the Cascade-Sierra Mountains Province. The Sierra Nevada range has a gentle western slope, and a very abrupt eastern slope. The Sierra Nevada consists of hilly to steep mountains and occasional flatter mountain valleys. Elevation ranges between 1,500 and 9,000 ft throughout most of the range, but peaks often exceed 12,000 ft. The highest point in the continental US occurs in this MLRA (Mount Whitney, 14,494 ft). Most of the Sierra Nevada is dominated by granitic rock of the Mesozoic age, known as the Sierra Nevada Batholith. The northern half is flanked on the west by a metamorphic belt, which consists of highly metamorphosed sedimentary and volcanic rocks. Additionally, glacial activity of the Pleistocene has played a major role in shaping Sierra Nevada features, including cirques, arêtes, and glacial deposits and moraines. Average annual precipitation ranges from 20 to 80 inches in most of the area, with increases along elevational and south-north gradients. Soil temperature regime ranges from mesic, frigid, and cryic. Due to the extreme elevational range found within this MLRA, Land Resource Units (LRUs) were designated to group the MLRA into similar land units.

LRU “X” represents ecological sites driven by abiotic features that override the typical soils or climatic features that drive most of the other LRU zones. In the Sierra Nevada these sites are typically driven by water features associated with lotic or lentic riparian systems. Other features maybe shallow bedrock, or unique chemical development which affects the growth of typical vegetation.

Ecological site concept

This riparian complex is associated with lacustrine (X) or delta valley (XI) types at elevations of 6,200 to 7,500 feet, and has an anastomosed Da channel type (Rosgen 1996). The channels are typically narrow and deep, but can be highly variable in sinuosity and width to depth ratio. The floodplains are saturated for most of the year, and the stream banks and floodplains have a dense cover of sedges (*Carex* spp.) and other grasslike species (*Scirpus* spp. and *Juncus* spp.). Willows (*Salix* spp.) and thinleaf alder (*Alnus incana* ssp. *tenuifolia*) are also common. The deep channels and access to deeper waters of the adjacent lakes provide ideal habitat for the North American beaver (*Castor canadensis*) (Beier and Barrett 1987). Beavers heavily influence the channel morphology with construction of dams, which pond and divert flow. The dominant soils have a histic epipedon, with sandy soils or layers of buried horizons below.

Associated sites

F022AE013CA	Frigid, Loamy, Volcanic Mountain Slopes This is a Jeffrey pine (<i>Pinus jeffreyi</i>)- white fir (<i>Abies concolor</i>) mixed conifer forest found on volcanic soils which occasionally surround this site.
F022AF002CA	Frigid, Sandy, Or Loamy Outwash This is a dry Jeffrey pine (<i>Pinus jeffreyi</i>) forest found on adjacent outwash and occasionally on the barrier beaches.
F022AF004CA	Frigid, Shallow To Deep, Sandy Mountain Slopes This is an open Jeffrey pine (<i>Pinus jeffreyi</i>) woodland with montane shrubs. It occurs on shallow to moderately deep soils on south facing slopes.
F022AF005CA	Frigid, Deep To Very Deep, Sandy-Loamy Mountain Slopes This is a Jeffrey pine (<i>Pinus jeffreyi</i>)- white fir (<i>Abies concolor</i>) forest, which occurs on deep to very deep granitic soils, on north facing mountain slopes.
F022AX100CA	Frigid, Sandy, Moist, Outwash Fan This is a Sierra lodgepole pine (<i>Pinus contorta</i> var. <i>murrayana</i>) forest, similar to community component 7, and occurs on outwash terraces.
R022AX003CA	Steep subalpine, valley bottom riverine system This ecological site has a single thread E-C type channel and has a cryic soil temperature regime.
R022AX004CA	Cryic basin peatland This fen complex occurs in closed basins, and soils are composed of sphagnum moss throughout.
R022AX102CA	Frigid E-C Meadow System This riparian complex has a single thread E-C type channel. It may occur just upstream of this ecological site.

Similar sites

R022AX101CA	Frigid Anastomosed System This is a single thread E-C type channel system. It may occur upstream from this site.
R022AX103CA	Cryic E Meadow System This is a single thread E-C channel system that occurs at higher elevations and has a cryic soil temperature regime.

Table 1. Dominant plant species

Tree	(1) <i>Populus tremuloides</i>
Shrub	(1) <i>Salix lemmonii</i> (2) <i>Alnus incana</i> subsp. <i>tenuifolia</i>
Herbaceous	(1) <i>Carex utriculata</i> (2) <i>Scirpus microcarpus</i>

Physiographic features

This site occurs on broad valley bottoms, with a lacustrine (X) or delta (XI) valley type. Glacio-lacustrine deposits are typically deeply buried below more recent river alluvium and organic material. Associated fluvial surfaces include ponds, fens, floodplains, floodplain steps, and outwash terraces. The fens and floodplains are seasonally flooded and ponded, for brief to very long durations from March through June. Ponding depth ranges from 0 to 12 inches. Slopes range from 0 to 5 percent, but the channel gradients are typically less than 0.5 percent. Elevations range from 6,220 feet to 9,420 feet, but are typically between 6,220 and 7,500 feet.

Table 2. Representative physiographic features

Landforms	(1) Fen (2) Flood plain (3) Outwash terrace
Flooding duration	Brief (2 to 7 days) to very long (more than 30 days)
Flooding frequency	None to frequent
Ponding duration	Brief (2 to 7 days) to very long (more than 30 days)
Ponding frequency	None to frequent
Elevation	6,220–9,420 ft
Slope	0–5%
Ponding depth	0–12 in
Water table depth	0–30 in
Aspect	Aspect is not a significant factor

Climatic features

The average annual precipitation ranges from 23 to 55 inches, and falls mostly in the form of snow from November to April. The mean annual air temperature ranges from 40 to 46 degrees Fahrenheit. The frost-free (>32F) season is 20 to 60 days, and the freeze-free (>28F) season is 40 to 90 days.

Climate stations: (1) 048762, Tahoe Valley FFA AP, California. Period of record 1968-2008

Table 3. Representative climatic features

Frost-free period (average)	65 days
Freeze-free period (average)	40 days
Precipitation total (average)	39 in

Influencing water features

This ecological site occurs in wetlands and meadows associated with a Da type channel system. Most of the area is saturated for long durations, but some areas have deeper water tables and are only saturated for a brief duration after snow melt.

Soil features

There are several soils associated with this ecological site. They are associated with different fluvial surfaces, and vary in wetness and organic matter accumulation. These soils are all very deep, and formed in organic matter over alluvium or alluvium derived from predominately granitic parent material. In some areas volcanic and metamorphic alluvium is also intermixed. Soils are very poorly drained with moderate permeability. The soil moisture regime is aquic and the soil temperature regime is frigid. Surface rock cover and subsurface rock fragments of all sizes are typically absent, but gravels may be up to 30 percent by volume in the lower horizons of the Watah soil.

Soils by fluvial surface:

Fens, active channel and floodplain:

The Watah (coarse-loamy, mixed, superactive, acid, frigid Histic Humaquepts) soil is the dominant component associated with this ecological site. The Watah soils develop in the wettest locations, in broad low lying areas among braided channels. The Watah soils have 20 to 40 cm of peat or mucky peat, developed primarily from decomposed sedge roots. Below the organic horizons the soil is sandier with gravelly mucky coarse sandy loam and gravelly loamy coarse sand textures. Gleyed soil colors occur below 73 cm. Community components 2 and 3 (CC2 and CC3) are associated with the Watah soils. CC2 occurs along the active channel where more flooding occurs, and CC3 occurs in areas where ponding is more common.

Floodplain step:

The Tahoe (coarse-loamy, mixed, superactive, acid, frigid Cumulic Humaquepts) soil occurs on floodplains typically on the outside of the braided channel network. The surface texture is mucky silt loam and subsurface textures are mucky silt loam and gravelly coarse sand. CC4 is associated with this soil component.

Floodplain step, dry:

The Tahoe gravelly phase (coarse-loamy, mixed, superactive, acid, frigid Cumulic Humaquepts) soil component occurs on floodplain steps, or old gravelly deposition areas. The surface texture is mucky gravelly silt loam. Subsurface textures are gravelly silt loam, gravelly sand, gravelly coarse sand, and gravelly fine sand. CC5 is associated with this soil component.

Outwash terrace:

The Marla (sandy, mixed, frigid Aquic Dystrochrepts) soils occur on outwash terraces that border this site, and are influenced by the ground water hydrology of the meadow. The surface textures are loamy coarse sand and subsurface textures are loamy coarse sand and clay loam. The Marla soils are associated with CC6.

This ecological site has been correlated with the following mapunits and soil components in the Tahoe Basin soil survey area (CA693):

Mapunit, Mapunit name, Component, Phase, percent

7011 Beaches Watah, 7

7051 Oxyaquic Xerorthents-Water association, 0 to 5 percent slopes, Watah, 1

7071 Watah peat, 0 to 2 percent slopes, Watah, 75

7071 Watah peat, 0 to 2 percent slopes, Tahoe, gravelly wet, 9

7071 Watah peat, 0 to 2 percent slopes, Tahoe, silt loam wet, 8

7071 Watah peat, 0 to 2 percent slopes, Marla, 3

9001 Bidart complex, 0 to 2 percent slopes, Watah, 5

Table 4. Representative soil features

Surface texture	(1) Mucky silt loam (2) Loamy coarse sand
Family particle size	(1) Loamy
Drainage class	Very poorly drained
Permeability class	Moderate
Soil depth	160 in
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-40in)	4-6.2 in
Calcium carbonate equivalent (0-40in)	0%
Electrical conductivity (0-40in)	0-20 mmhos/cm

Soil reaction (1:1 water) (0-40in)	5.1–7.3
Subsurface fragment volume <=3" (Depth not specified)	0–30%
Subsurface fragment volume >3" (Depth not specified)	0%

Ecological dynamics

Abiotic Factors:

This ecological site is a riparian complex associated with lake deltas and lacustrine fringes. The broad, low gradient valleys allow the streams to develop into anastomosed-multi-channel streams, also referred to as a Da channel type (Rosgen 1996). Beaver activity is high in these systems. Several community components (a combination of soil type and associated plant community) related to the depth of water table, micro-topography, and different fluvial surfaces are present. The dominant soils are saturated late into summer, and have developed deep organic surfaces from decomposed sedge roots. A gradation of soil and plant communities exists from the wettest areas, with sedges and organic soils to drier grass and forb communities on poorly developed sandy soils. Willows, thinleaf alder, quaking aspen (*Populus tremuloides*), and Sierra lodgepole pine (*Pinus contorta* var. *murrayana*) have niches within this spectrum.

Hydrologic factors:

This ecological site occurs on low gradient (typically less than 1 percent) deltas of lakes or ponds. A braided network of channels, defined as Da channel type, is the natural stream type for this system. Beaver prefer the deep channels and lake access that this site provides for mobility, safety, and den building (Beier and Barrett 1987). They are abundant and very active in this ecological site. Dam building raises the water table above the dam, causing pond development and flooding across the floodplain. Below the beaver dam, or after dams are removed, the channel may become unstable and scour or widen depending upon the site. Aerial imagery from 1940 shows dendritic-branching patterns, indicating the natural branching Da system prior to that time. The branching system may have developed with historic beaver activity or from the natural tendency for channels to meander and anastomose in these river delta systems.

A properly functioning marsh system is an important filtration system. The broad, low gradient floodplain allows for suspended sediments to settle out of suspension in the slow moving flood waters and ponds. Subsurface flow is filtered as it moves through the thick organic surface. Nutrients such as phosphorus and nitrogen are also deposited with the sediments reducing the load distributed to the lake.

Beavers were once thought to be non-native to the Sierra Nevada, but carbon dating of old beaver dams has shown that beavers were in the Sierra Nevada since AD 580 (James and Lanman 2012, Landman et al. 2012). Beaver trapping and extirpation efforts eliminated beavers from the higher Sierra Nevada by mid-1800s. Beavers were reintroduced into the Lake Tahoe Basin from 1939 to 1949 (Beier and Barrett 1987) after 100 years or more of absence. Nine beavers were introduced from the Snake River in Idaho to the Truckee River. Since then populations have expanded to many watersheds around Lake Tahoe. In 1987 there were 0.72 colonies (3.5 beavers) per km of stream along the Truckee River (Beier and Barrett 1987).

Anastomosed channels can have high variability. They are typically deep and narrow, but wide ponds develop above beaver dams, and some channels, particularly below beaver dams may become wide and shallow. The soils are saturated to the surface for most of the summer. The dominant soils in this ecological site have a 20 to 40 cm thick histic (deep organic) horizon, composed of poorly decomposed peat, primarily from sedge roots and leaf material. Saturated conditions slow the decomposition of the plant material, allowing for accumulation of plant material over time. The thick organic material has high pore capacity and allows for easy surface flow in the upper soil, as well as filtering fine sediments and nutrients. Beaver dams may increase sediment and phosphorous retention during high flows (Muskopf 2007). Soils in higher or drier locations lack the deep histic horizon, but typically have a rich organic surface mixed with mineral soils. In some areas depositional episodes are evident by layers of buried horizons, identified by the organic layers below sandier deposits.

Disturbance Factors:

In addition to beaver activity, these areas have been impacted by dams, historic logging, grazing, road and marina development.

During the mid-1800's approximately 80 percent of the forests in Tahoe were clear-cut during the Comstock era (Elliott-Fisk et al. 1996). The removal of the trees increased erosion on the mountain slopes and increased the sediment supply to these river systems. The removal of trees would also have reduced water loss to evapotranspiration and increased water flow into the streams. The side effects of the logging era on the stream morphology are unknown, but likely increased sediment supply and stream volume in these systems, and may have caused channel progradation (Elliott-Fisk et al. 1996). Davis (DAVIS 1996) report increased erosion between 1850 and 1900 based on analysis of cores from Lake Tahoe. The erosion was attributed to logging, grazing and road development during this period. During the same period, there was an increase in *Pediastrum* a genus of algae, and an increase in sedge (*Carex* spp.) pollen, possibly due to increased nutrient loading into the lake and tributaries.

The meadows in the Lake Tahoe Basin were grazed in the past by cattle and sheep. Over 13 dairy farms were active in the basin, and most meadows were fenced for cattle grazing. Sheep roamed the mountains, and denuded much of the forbs and grasses (Elliott-Fisk et al. 1996). Cattle concentrate in riparian habitats because of the access to water and forage, as well as shade and gently landscapes (Kie and Boroski 1996). The cattle may have affected the present composition of vegetation by selectively grazing species and trampling susceptible species. Grazing can affect channel morphology by removing bank stabilizing vegetation, such as willows, and by trampling the stream banks with their hooves when accessing the stream.

Fine sediment and nutrient loading (which increases algal growth) are currently the primary factors attributed to reduced lake clarity in Lake Tahoe Basin (Lahontan Water Board and Nevada Division of Environmental Protection 2008). Development within many of these river mouths for marinas and housing have channelized and confined these streams and floodplains, creating a rapid transport system for sediments and nutrients, rather than the diffuse and slow depositional environment.

This hydrology of this ecological site is typically influenced by an adjacent lake or pond. The construction of the Lake Tahoe Dam between 1909 and 1913 raised the water level of the lake by a maximum of 6 feet (AECOM and ENTRIX 2013) above the natural rim (6,223'). Many smaller lakes in the area also have small dams. High lake levels can flood into these low marshes. The development of the Tahoe Keys has altered the groundwater gradient in the Truckee Marsh (AECOM and ENTRIX 2013) by shortening distance and increasing the hydrologic gradient from the marsh to lake level. Groundwater flows west to the marina water level rather than to the north to Lake Tahoe (AECOM and ENTRIX 2013). The consequence of these and other hydrological changes in the Truckee Marsh has been a lowering of the ground water table during dry years (when lake levels are low), and the development of an incised highly unstable F channel. Vegetation has shifted to drier upland species in many areas, and the area influenced by saturated conditions and frequent ponding and flooding has decreased.

Ecological sites associated with lotic stream systems are developed using channel evolution models. Stream systems are dynamic and continually evolve to reach a stable equilibrium. Streams develop identifiable stages of development based on channel morphology (Rosgen 1997). These stages are identified using a state and transition model based on stream evolution models.


The reference state is typically the pre-settlement, most successional advanced and hydrologically stable community phase (numbered 1.1), and the community phases that result from natural and human disturbances. However, the reference state is speculative in this case, and has been altered by conditions mentioned above, and is therefore referred to the representative state. Community phase 1.1 is deemed the phase representative of the most successional advanced pre-European plant/animal community including beaver activity and hydrologic conditions that influence its composition and production. Because this phase is partly determined from reconstruction and/or historic literature, some speculation is necessarily involved in describing it.

All tabular data listed for a specific community component within this ecological site description represent a summary of one or more field data collection plots taken in modal communities within the community component. Although such data are valuable in understanding the community component (kinds and amounts of ground and surface materials, canopy characteristics, community phase species, production and composition, and growth), they do not represent the absolute range of characteristics or an exhaustive listing of all species that may occur in that phase over the geographic range of the ecological site.

State and transition model

R022AX101CA, Frigid Delta System

Populus tremuloides / *Salix lemmonii* - *Alnus incana* ssp. *tenuifolia* / *Carex utriculata* - *Scirpus microcarpus*
(quaking aspen / Lemmon's willow - thinleaf alder / Northwest Territory sedge - panicked bulrush)



I.D.	Plant Association	Fluvial Surface/Landform
1	Aquatic community, yellow pond lily	Pond- basins
2	Thinleaf alder, Panicked bulrush, Northwest Territory sedge	Fen, Floodplains
3	Northwest Territory sedge, water sedge	Floodplains
4	Lemmon's willow, Geyers willow, Northwest Territory sedge	Floodplain step
5	Arctic rush, tufted hairgrass, chamisso arnica	Floodplain step, dry
6	Quaking aspen forest	Floodplain step, wet
7	Sierra lodgepole pine forest	Outwash Terrace

Figure 6. R022AX101CA STM

State 1

Da Channel

This state is the representative state for this ecological site. It is a Rosgen Da channel type, highly influenced by beaver activity.

Community 1.1

Da Channel



Figure 7. CC2



Figure 8. CC3



Figure 9. CC5



Figure 10. CC6



Figure 11. CC7



Figure 12. CC1

This phase is composed of a stable Da channel type, with several community components. Community Component 1 (CC1), Lagoons and ponds This community develops in lagoons and ponds formed by high lake levels, beaver dams, or other obstructions that cause ponding. There is variation among these open water areas. Beaver ponds typically lack aquatic vegetation. Species documented in data transects include yellow pond lily (*Nuphar lutea*), pondweeds (*Potamogeton* sp.), Quillwort (*Isoetes* sp.), white water crowfoot (*Ranunculus aquatilis*), bur-reed (*Sparganium* sp.), and lesser bladderwort (*Utricularia minor*). Other native species observed in these areas include watershield (*Brasenia schreberi*), and common mare's-tale (*Hippuris vulgaris*). Non non-native aquatic species include Eurasian watermilfoil (*Myriophyllum spicatum*) and curly pondweed (*Potamogeton crispus*). These non-native aquatic species have established in ponds and lagoons within this ecological site. They are being managed to reduce growth and new infestations, since they release nutrients into the water, which can contribute to algal growth and reduce the clarity of Lake Tahoe (USACE, 2009). CC2, Active channels and adjacent floodplain This community develops along the main channel. It is enhanced by beaver activity, and is occurs at the margins of beaver ponds and side drainages. Thinleaf alder (*Alnus incana* spp. *tenuifolia*) and Lemmon's willow (*Salix lemmonii*) are dominant shrubs. The understory is dominated by a mix of sedges or other grass-like plants including water sedge (*Carex aquatilis* var. *aquatilis*), panicled bulrush (*Scirpus microcarpus*), Northwest Territory sedge (*Carex utriculata*), blister sedge (*Carex vesicaria*), lakeshore sedge (*Carex lenticularis*), slimstem reedgrass (*Calamagrostis stricta*). These sedges are sometimes intermixed, but more often have distinct patches due to wetness or proximity to flowing or open water. Water sedge, blister sedge, and lakeshore sedge tend to establish along the greenline of the stream channel or along the boundary of the lake and pond. Panicled bulrush is often under the thinleaf alder in high beaver activity areas, while Northwest Territory sedge tends to establish in ponded areas away from the active channel. Forbs have relatively low cover in this community, but may include Pacific onion (*Allium validum*), fringed willowherb (*Epilobium ciliatum*), field horsetail (*Equisetum arvense*), and tinker's penny (*Hypericum anagalloides*). The Watah soils, when associated with this community component, have a deep histic horizon, up to 40 cm thick. Sometimes the soils have layers of histic horizons inter-mixed with sandy horizons. This may be due to high beaver activity, and associated stream alterations. CC3, Floodplains and depressions This community can develop large expanses of sedge dominated meadow. It occurs among the braided channels or low topographic positions, and is dominated by Northwest territory sedge and/or blister sedge. Other sedges and mountain rush (*Juncus arcticus* ssp. *littoralis*) are also present. Forbs and shrubs are uncommon. This community is associated with the Watah soils. CC4, Floodplain step This community is composed of a drier willow-sedge community. It is found on slightly higher topography than CC3 on soils that do not have a histic epipedon, and has more bare ground and forbs than the CC1 willow-sedge community. Shrubs may compose 15 to 30 percent cover. Lemmon's willow is typically dominant, with Geyer willow (*Salix geyeriana*) common. Woods' rose (*Rosa woodsii*) and whitestem gooseberry (*Ribes inerme*) can be found under or near the willows. CC5, Floodplain step, dry This community occurs at the edge of the active floodplain. The water table lowers quickly through the season, reaching depths of 60 to 90 cm in normal water years. This community has lower vegetative cover, and higher cover of bare ground. Mexican rush (*Juncus mexicanus*), tufted hairgrass (*Deschampsia cespitosa*), Nebraska sedge, brownhead rush (*Juncus phaeocephalus*), Kentucky bluegrass (*Poa pratensis*), Timothy (Phleum pretense) and a diversity of other sedges are present. There is some debate whether Kentucky bluegrass is native to California, and Timothy is a non-native grass. Forbs include upland early pioneer species such as Chamisso arnica (*Arnica chamissonis*), lupine (*Lupinus* sp.), slender cinquefoil (*Potentilla gracilis*), and longstalk clover (*Trifolium longipes*). CC6, Floodplain step, wet This community is not present in all locations, but when present it occurs where the stream meets the meadow, and along the meadow boundary in wet positions. Quaking aspen is dominant with 35 to 70 percent canopy cover. Canopy height is around 40 to 50 feet tall. Black cottonwood (*Populus balsamifera* ssp. *trichocarpa*) is occasionally present and overtops the aspen canopy at approximately 90 feet in height. Common understory shrubs include thinleaf alder, redosier dogwood (*Cornus sericea*), and whitestem gooseberry. Grass and grasslike species include blue wildrye, bentgrass (*Agrostis* sp.), water sedge, blue wildrye (*Elymus glaucus*), and mannagrass (*Glyceria* sp.). Forbs are diverse but typically include Pacific onion, common cowparsnip (*Heracleum maximum*), common mare's-tail, feathery false lily of the valley (*Maianthemum racemosum* ssp. *racemosum*), and California false hellebore (*Veratrum californicum*). This community component has some similarities to the steeper A-B stream complex, R022AX105CA. CC7, Outwash terrace This community component occurs at the edge of the meadow complex or on raised topography within the meadow. Sierra lodgepole pine is characteristic of this community, and can develop a multi-tiered forest with up to 25 percent cover. The understory is dominated by upland species, with some meadow species intermixed. This community develops on drier areas, but can encroach into the wetter soils of the meadow during years of lower precipitation and lower groundwater tables. Fire is important in maintaining open forests at the edge of the meadow, and initiates forest regeneration. This community is similar to F022AX100CA, which may be referred to for more information about the successional dynamics of the Sierra lodgepole pine forest.

Table 5. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	50	1800	3500
Shrub/Vine	30	1500	3000
Tree	0	20	350
Forb	0	60	280
Total	80	3380	7130

Table 6. Canopy structure (% cover)

Height Above Ground (Ft)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.5	—	—	—	1-15%
>0.5 <= 1	0-1%	—	—	1-5%
>1 <= 2	0-1%	—	10-90%	0-1%
>2 <= 4.5	0-1%	0-5%	0-5%	—
>4.5 <= 13	0-10%	0-25%	—	—
>13 <= 40	0-15%	—	—	—
>40 <= 80	0-25%	—	—	—
>80 <= 120	—	—	—	—
>120	—	—	—	—

State 2

G-F Channel

When these streams are straightened or confined, velocity and erosive power increase. These streams then head cut or down cut to form an entrenched low gradient “G” type channel. The unstable banks of the “G” type channel erode rapidly into a wider and shallower entrenched “F” type channel.

Community 2.1

G Channel

This is a transitional channel with steep unstable banks. G channels are entrenched, with low wide to depth ratio, and moderate sinuosity. Plant communities transition from state 2, phase 1 to state 2 phase 2 during this transition. The first notable change is the loss of the channel stabilizing sedge, willow and thinleaf alder roots on the stream banks. Obligate wetland species in CC2 and CC3, further from the channel, decline in cover over time as the water table in the meadow drops to meet the channel bottom.

Community 2.2

F Channel



Figure 14. Straightened Channel by Tahoe Keys

The “F” type channel is entrenched, with moderate to wide depth ratio, and moderate sinuosity. These channels are wide, and shallow with nearly vertical, unstable, poorly vegetated banks. The associated community components are similar to state 1, phase 1, but the composition of the plant communities shifts to dominance by the drier upland communities CC4 and CC5, and species composition is altered within communities. These confined channels maintain high stream velocity, and very rarely overtop the banks to reach the former floodplain. These channels provide poor habitat for beaver. The loss of vegetative cover on the banks leaves beavers exposed and requires farther travel for forage and dam building supplies. The stream velocity can also be too strong and burst the beaver dams. CC1, Lagoons and ponds This community declines in this phase. The entrenched stream is less likely to develop natural or beaver ponds, and the area that potentially develops lagoons from lake water is smaller. However, some areas such as Pope Marsh may have become more ponded by development. Pope Marsh has high cover of yellow pond-lily and common mare’s tail. CC2, Active channels and adjacent floodplain This community severely declines in this phase since it is dependent upon a high water table and frequent flooding or ponding. Thinleaf alder and panicked bulrush are nearly absent. Lemmon’s willow, Northwest Territory sedge, blister sedge (*Carex vesicaria*), and lakeshore sedge (*Carex lenticularis*) establish at low levels along the channel. This community exists in isolated patches along the channel and in old oxbows that are shallower to the ground water table. CC3, Floodplains and depressions This community remains extensive in lower areas. Species dominance shifts to Nebraska sedge and Northwest territory sedge. CC4, Floodplain step This community increases in extent as it establishes close to the channel, replacing the wetter willow community (CC2). CC5, Floodplain step, dry This community increases as it moves into the drier abandoned floodplain. Overall cover and production in this phase is lower than in state 2, phase 2. CC6, Floodplain step, wet This community is typically absent when in this phase due to a combination of a lower water table, heavy beaver browse, or grazing. CC7, Outwash terrace This community increases in this phase as Sierra lodgepole pine establishes on the drier abandoned floodplains.

Pathway 2.1a

Community 2.1 to 2.2

With time and continued bank erosion, the "G" channel widens into the wide, shallow, entrenched "F" channel.

State 3

C New Floodplain

A “C” type channel is slightly entrenched, with moderate to wide width to depth ratios, and moderate to high sinuosity. Since, these channels are less entrenched than the F channel, bank erosion is less severe, and vegetation can establish on the floodplains and banks, creating more stability. This state may coexist within reaches of state 2 as it develops.

Community 3.1

C Channel New Floodplain

Data is lacking for this phase, since most of the altered channel presently exists in State 2. However, based on observations in stream sections showing initial signs of C channel development, CC2 is re-establishing along the new channel banks and floodplains. CC3 and CC4 are persisting on the abandoned floodplains. CC5 may decline

slightly as the floodplain expands, and CC7 is unchanged. The development of this state and phase, creates a narrow band of wetter community types, but still has a large abandoned floodplain, which will continue to be dominated by drier community types than were historically in those positions.

Transition 1A State 1 to 2

Transition to State 2 occurs with the channelization, straightening or confinement of the stream channel, removal or absence of beavers, or blowout of a beaver dam. The alteration causes the channel to become unstable as the concentration or change in water course causes bank erosion and channel bottom incision. The lowering of the water table causes a decline in wetland obligate species, and loss of vegetative structure along the banks, causing further bank instability. Past influences such as logging and grazing may have caused this transition in some areas. Since this system has braided channels, it is possible that a given meadow may have channels that exhibit both State 1 and State 2 channel types.

Restoration pathway 2a State 2 to 1

The Tahoe Conservancy and other agencies have developed and environment impact report for the Upper Truckee River and Marsh Restoration Project. The plans include several alternatives, but all of them focus on increasing the stream meander, and raising the elevation of the channel bed to increase flooding frequency on the original floodplain (AECOM and ENTRIX 2013).

Transition 2a State 2 to 3

This transition occurs when the stream has reached a relative equilibrium by eroding the banks to a sufficient width to contain regular flows. The bank full flows now erode a new channel deeper within wide F channel, creating a new, slightly-entrenched, C channel with a new floodplain.

Restoration pathway 3a State 3 to 1

The Tahoe Conservancy and other agencies have developed and environment impact report for the Upper Truckee River and Marsh Restoration Project. The plans include several alternatives, but all of them focus on increasing the stream meander, and raising the elevation of the channel bed to increase flooding frequency on the original floodplain (AECOM and ENTRIX 2013).

Additional community tables

Table 7. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
Forb					
1	Ponds			15–150	
	yellow pond-lily	NULU	<i>Nuphar lutea</i>	0	0–30
	pondweed	POTAM	<i>Potamogeton</i>	0	0–5
	white water crowfoot	RAAQ	<i>Ranunculus aquatilis</i>	0	0–3
	bur-reed	SPARG	<i>Sparganium</i>	0	0–2
	lesser bladderwort	UTMI	<i>Utricularia minor</i>	0	0–1
	quillwort	ISOET	<i>Isoetes</i>	0	0–1
2	Fen			0–12	
	Pacific onion	ALVA	<i>Allium validum</i>	0–10	0–1
	fringed willowherb	EPCI	<i>Epilobium ciliatum</i>	0–1	0–3

	field horsetail	EQAR	<i>Equisetum arvense</i>	0–1	0–1
	tinker's penny	HYAN2	<i>Hypericum anagalloides</i>	0–1	0–1
4	Floodplain step			0–130	
	primrose monkeyflower	MIPR	<i>Mimulus primuloides</i>	0–110	0–5
	curvepod yellowcress	ROCU	<i>Rorippa curvisiliqua</i>	0–20	0–2
	longleaf starwort	STLO	<i>Stellaria longifolia</i>	0–1	0–1
	fringed willowherb	EPCI	<i>Epilobium ciliatum</i>	0–1	0–1
5	Floodplain step, dry			10–60	
	Chamisso arnica	ARCH3	<i>Arnica chamissonis</i>	1–50	1–5
	lupine	LUPIN	<i>Lupinus</i>	1	1
	slender cinquefoil	POGR9	<i>Potentilla gracilis</i>	1	1
	longstalk clover	TRLO	<i>Trifolium longipes</i>	1	1
6	Floodplain step, wet			50–280	
	California false hellebore	VECA2	<i>Veratrum californicum</i>	0–120	0–8
	Pacific onion	ALVA	<i>Allium validum</i>	0–100	0–5
	feathery false lily of the valley	MARAR	<i>Maianthemum racemosum</i> ssp. <i>racemosum</i>	0–40	0–3
	common cowparsnip	HEMA80	<i>Heracleum maximum</i>	0–20	0–1
	common mare's-tail	HIVU2	<i>Hippuris vulgaris</i>	0–10	0–1
7	Outwash terrace			25–120	
	slender cinquefoil	POGR9	<i>Potentilla gracilis</i>	5–100	1–3
	longleaf starwort	STLO	<i>Stellaria longifolia</i>	0–5	0–1
	western mountain aster	SYSPS	<i>Symphotrichum spathulatum</i> var. <i>spathulatum</i>	0–5	0–1
	Sierra pea	LANE3	<i>Lathyrus nevadensis</i>	0–5	0–1
	feathery false lily of the valley	MARAR	<i>Maianthemum racemosum</i> ssp. <i>racemosum</i>	0–5	0–1
	sweetcicely	OSBE	<i>Osmorhiza berteroi</i>	0–5	0–1
	fireweed	CHANC	<i>Chamerion angustifolium</i> ssp. <i>circumvagum</i>	0–5	0–1
	willowherb	EPILO	<i>Epilobium</i>	0	0–1
	fleabane	ERIGE2	<i>Erigeron</i>	0	0–1
	Virginia strawberry	FRVI	<i>Fragaria virginiana</i>	0	0–1
	groundsmoke	GAYOP	<i>Gayophytum</i>	0	0–1
	tinker's penny	HYAN2	<i>Hypericum anagalloides</i>	0	0–1
	Chamisso arnica	ARCH3	<i>Arnica chamissonis</i>	0	0–1
	Gray's licorice-root	LIGR	<i>Ligusticum grayi</i>	0	0–1
	lupine	LUPIN	<i>Lupinus</i>	0	0–1
Shrub/Vine					
2	Fens			500–1500	
	thinleaf alder	ALINT	<i>Alnus incana</i> ssp. <i>tenuifolia</i>	500–1500	10–25
	Lemmon's willow	SALE	<i>Salix lemmonii</i>	10–500	1–5
4	Floodplain step			1200–3000	
	Lemmon's willow	SALE	<i>Salix lemmonii</i>	700–4000	3–50
	Geyer willow	SAGE2	<i>Salix geyeriana</i>	600–2200	5–15

	whitestem gooseberry	RIIN2	<i>Ribes inerme</i>	0–20	0–1
	Woods' rose	ROWO	<i>Rosa woodsii</i>	0–10	0–1
6	Floodplain step, wet			100–340	
	thinleaf alder	ALINT	<i>Alnus incana ssp. tenuifolia</i>	20–250	1–10
	whitestem gooseberry	RIIN2	<i>Ribes inerme</i>	10–90	1–5
	redosier dogwood	COSE16	<i>Cornus sericea</i>	10–50	1–2
7	Outwash terrace			30–200	
	whitestem gooseberry	RIIN2	<i>Ribes inerme</i>	10–60	1–5
	Woods' rose	ROWOU	<i>Rosa woodsii var. ultramontana</i>	0–60	0–5
	Lemmon's willow	SALE	<i>Salix lemmonii</i>	0–50	0–3
	Saskatoon serviceberry	AMAL2	<i>Amelanchier alnifolia</i>	0–30	0–2
	hollyleaved barberry	MAAQ2	<i>Mahonia aquifolium</i>	0–20	0–1
	wax currant	RICE	<i>Ribes cereum</i>	0–10	0–1
Grass/Grasslike					
2	Fens			1000–3500	
	Northwest Territory sedge	CAUT	<i>Carex utriculata</i>	1300–2800	32–65
	panicked bulrush	SCMI2	<i>Scirpus microcarpus</i>	75–2700	1–40
	blister sedge	CAVE6	<i>Carex vesicaria</i>	300–1800	10–48
	lakeshore sedge	CALE8	<i>Carex lenticularis</i>	0–1000	0–40
	water sedge	CAAQ	<i>Carex aquatilis</i>	0–900	0–30
	slimstem reedgrass	CAST36	<i>Calamagrostis stricta</i>	1–200	1–10
	sedge	CAREX	<i>Carex</i>	1–10	1–2
3	Floodplain			1000–3000	
	blister sedge	CAVE6	<i>Carex vesicaria</i>	0–2800	0–60
	Northwest Territory sedge	CAUT	<i>Carex utriculata</i>	1000–1800	35–50
	mountain rush	JUARL	<i>Juncus arcticus ssp. littoralis</i>	10–500	1–35
	sedge	CAREX	<i>Carex</i>	1–50	1–2
4	Floodplain step			800–2500	
	Northwest Territory sedge	CAUT	<i>Carex utriculata</i>	10–1900	1–45
	Nebraska sedge	CANE2	<i>Carex nebrascensis</i>	150–1800	5–55
	bigleaf sedge	CAAM10	<i>Carex amplifolia</i>	0–1000	0–80
	rough bentgrass	AGSC5	<i>Agrostis scabra</i>	0–488	0–20
	fowl mannagrass	GLST	<i>Glyceria striata</i>	0–100	0–4
5	Floodplain step, dry			500–1500	
	Mexican rush	JUME4	<i>Juncus mexicanus</i>	250–1000	12–45
	brownhead rush	JUPH	<i>Juncus phaeocephalus</i>	0–500	0–30
	tufted hairgrass	DECE	<i>Deschampsia cespitosa</i>	20–350	1–15
	Nebraska sedge	CANE2	<i>Carex nebrascensis</i>	10–200	1–8
	Kentucky bluegrass	POPR	<i>Poa pratensis</i>	0–200	0–5
	sedge	CAREX	<i>Carex</i>	10–150	1–5
	timothy	PHPR3	<i>Phleum pratense</i>	0–30	0–2
6	Floodplain step, wet			50–500	

	water sedge	CAAQ	<i>Carex aquatilis</i>	0–300	0–20
	blue wildrye	ELGL	<i>Elymus glaucus</i>	10–200	1–10
	bentgrass	AGROS2	<i>Agrostis</i>	0–30	0–3
	mannagrass	GLYCE	<i>Glyceria</i>	0–10	0–1
7	Outwash terrace			50–400	
	Kentucky bluegrass	POPR	<i>Poa pratensis</i>	10–250	1–15
	bentgrass	AGROS2	<i>Agrostis</i>	0–150	0–30
	mountain rush	JUARL	<i>Juncus arcticus ssp. littoralis</i>	10–100	1–3
	Pringle's phacelia	PHPR	<i>Phacelia pringlei</i>	0–60	0–3
	timothy	PHPR3	<i>Phleum pratense</i>	0–60	0–3
	California brome	BRCA5	<i>Bromus carinatus</i>	0–30	0–3
	analogue sedge	CASI2	<i>Carex simulata</i>	0–20	0–2
	meadow barley	HOBR2	<i>Hordeum brachyantherum</i>	0–10	0–1
	sedge	CAREX	<i>Carex</i>	0–10	0–1
Tree					
6	Floodplain step, wet			50–250	
	quaking aspen	POTR5	<i>Populus tremuloides</i>	50–180	5–15
	black cottonwood	POBAT	<i>Populus balsamifera ssp. trichocarpa</i>	0–75	0–3
7	Outwash terrace			10–100	
	Sierra lodgepole pine	PICOM	<i>Pinus contorta var. murrayana</i>	10–100	2–10

Animal community

This ecological site is important habitat for beaver. The deep low velocity channels allow beaver to build dams and lodges. The higher velocity and shallow waters of altered channels provide lower quality with a greater chance of dam failures.

A variety of birds nest in the cover of the sedges. Osprey and bald eagles nest in lodgepole pine snags, and are frequently seen in these areas. Coyotes are abundant in the drier areas of this site, and use the willows for cover.

The rivers provide habitat for a variety of fish. Lagoons support native frogs, but the American bullfrog has become an invasive pest in some areas.

Hydrological functions

This ecological site provides for water catchment, sediment and nutrient storage.

Recreational uses

These areas are typically wet, but great for viewing wildlife from the edges.

Inventory data references

The following NRCS vegetation plots were used to describe this ecological site.

State 1

CT1
094
mh03093

CT2
097
GR02210
GR03104
LO02210
Mh02667

CT3
Mh02670
Mh03092
Mh04003

CT4
Mh02676
Mh03093
Mh04020
Mh04019

CT5
Mh02543
Mh02668
Mh02669
Mh04021

CT6
124
110 (notes)

CT7
Mh02673
Mh02675
187

State 2

CT1
Mh02537

CT2
Lo04018Plot17
mh02536- Watah Type location

CT3
Lo04018Plot10
Lo04018Plot15
Lo04018Plot16
Lo04018Plot18
Lo04018Plot19
Lo04018Plot20
Lo04018Plot21
Lo04018Plot22
Lo04018Plot9
Mh02672

CT4
Lo04018Plot14
Lo04018Plot23
Mh02666

CT5
Lo04018Plot1
Lo04018Plot2&3
Lo04018Plot4&5
Lo04018Plot6
Lo04018Plot7
Lo04018Plot8
mh02535
Mh02671
Mh02678

CT7
Mh02673
Mh02675

Type locality

Location 1: El Dorado County, CA	
UTM zone	N
UTM northing	4313607
UTM easting	758359
General legal description	The Watah type location is between Pope Marsh and the Tahoe Keys, in the Lake Tahoe Basin.

Other references

AECOM, and C. ENTRIX. 2013. Upper Truckee River and Marsh Restoration Project (EIR).in C. T. Conservancy, editor.

Beier, P., and R. H. Barrett. 1987. Beaver habitat use and impact in Truckee River Basin, California. Journal of Wildlife Management 51:794-799.

Davis, O. K. 1996. Pollen Analysis of a Mid-Lake Core from Lake Tahoe, California: Historic Vegetation Change. Sierra Nevada Ecosystem Project: Final report to Congress, Addendum.

Elliott-Fisk, D. L., T. A. Cahill, O. K. Davis, L. Duan, C. R. Goldman, G. E. Gruell, R. Harris, R. Kattelman, R. Lacey, D. Leisz, S. Lindstrom, D. Machida, R. A. Rowntree, P. Rucks, D. A. Sharkey, S. L. Stephens, and D. S. Ziegler. 1996. Tahoe Case Study. Sierra Nevada Ecosystem Project: Final Report to Congress.

James, C. D., and R. B. Lanman. 2012. Novel physical evidence that beaver historically were native to the Sierra Nevada. California Fish and Game 98:129-132.

Kie, J. G., and B. B. Boroski. 1996. Cattle distribution, habitats, and diets in the Sierra Nevada of California. Journal of Range Management 49:482-488.

Lahontan Water Board, and Nevada Division of Environmental Protection. 2008. Charting the Course to Clarity.

Landman, R., H. Perryman, D. Brock, and C. D. James. 2012. The historical range of beaver in the Sierra Nevada: a review of the evidence. California Fish and Game 98:65 to 80.

Muskopf, S. 2007. The effect of beaver (*Castor canadensis*) dam removal on total phosphorus concentration in Taylor Creek and Wetland, South Lake Tahoe, California. . Natural Resources M.S.:35.

Rosgen, D. L. 1996. Applied river morphology. Wildland Hydrology, Pagosa Springs, Colo.

Rosgen, D. L. 1997. A Geomorphological Approach to Restoration of Incised Rivers in Proceedings of the Conference on Management of Landscapes Disturbed by Channel Incision.

Contributors

Marchel Munnecke

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

-
7. **Amount of litter movement (describe size and distance expected to travel):**
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-
14. **Average percent litter cover (%) and depth (in):**
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
