

Ecological site R023XY119OR BASIN WILLOW

Accessed: 05/18/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

Associated sites

R023XY104OR	LOAMY BOTTOM Loamy Bottom
R023XY116OR	SEMI-WET MARSH Semi-Wet Marsh
R023XY117OR	BASIN WET MEADOW Basin Wet Meadow
R023XY118OR	BASIN DRY MEADOW Basin Dry Meadow

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) Salix exigua
Herbaceous	Not specified

Physiographic features

This site occurs in dry lake basins and valleys. It typically occurs on banks of perennial or intermittent drainage

systems. Slopes range from 0 to 3 percent. Elevation varies from 4000 to 4600 feet.

Landforms	(1) Valley (2) Lakebed
Flooding frequency	Occasional
Elevation	1,219–1,402 m
Slope	0–3%
Water table depth	25–76 cm
Aspect	Aspect is not a significant factor

Table 2. Representative physiographic features

Climatic features

Influencing water features

Soil features

The soils of this site are very deep over lacustrine and alluvial sediments. Influenced by periodic flooding they are quite variable, often occuring as small naturally built-up levees along stream edges. Typically they have a loamy surface over a loamy to clay loam subsoil. Gravels are common. Along banks and berms localized seasonal surface drainage and aeration is good. This is desirable for willow growth and root structure. However, they are hydric with an overall drainage classification of somewhat poor. A seasonal water table occurs at 10 to 30 inches. Permeability is moderate. The available water holding capacity (AWC) is 6 to 10 inches.

Table 3. Representative soil features

Surface texture	(1) Gravelly loam
Family particle size	(1) Clayey
Drainage class	Somewhat poorly drained
Permeability class	Moderate
Available water capacity (0-101.6cm)	15.24–25.4 cm

Ecological dynamics

Range in Characteristics:

Seasonal availability and depth to groundwater influences the productivity of willow and other overstory species. With an increase in fire frequency, basin big sagebrush will decrease and basin wildrye will increase.

Response to Disturbance:

If the condition of the site deteriorates as a result of overgrazing or drainage, basin wildrye will decrease while creeping wildrye, basin big sagebrush, gray rabbitbrush, and greasewood will increase. Heavy suumer and fall use results in heavy hedging of willow. With further deterioration, banks become unstable, streams become incised and groundwater tables are lowered. Shrubs will continue to increase, annual invade, and areas of bareground increase. The stand of willows will develop a "tunnelled" appearance.

State and transition model



GENERAL MODEL FOR COOL-SEASON BUNCHGRASS RANGELANDS

State 1 Reference State

Community 1.1 Reference Plant Community

The potential native plant community is strongly dominated by willows. Coyote willow, a rhizomatous species, is clearly dominant followed by yellow and Pacific willow. Other shrubs and a variety of forbs and grasses are present. The potential vegetative composition is approximately 80 percent shrubs, 10 percent grass and grass-like, and 5 percent forbs. Approximate ground cover is 80-120 percent (basal and crown).

Table 4. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Shrub/Vine	1681	2522	3363
Grass/Grasslike	448	673	897
Forb	112	168	224
Total	2241	3363	4484

Additional community tables

Table 5. Community 1.1 plant community composition

Grass/like 1 Perennial, rhizomatous grass 168–336 1 beardless wildrye LETR5 Leymus triticoides 168–336 2 Other grasslike 135–538 2 Other grasslike Carex 67–269 5 Other perennial grasses 67–269 - 5 Other perennial grasses 67–269 - 6 shortawn foxtail ALAE Alopecurus aequalis 0–67 - 6 shortawn foxtail ALAE Alopecurus aequalis 0–67 - 1 teal lovegrass ERHY Eragrostis hypnoides 0–67 - 1 basin wildrye LEC14 Leymus cinereus 0–67 - 9 Other perennial forbs Poa 0–67 - - 9 Other perennial forbs Argentina anserina 0–67 - - 1 cowparsnip HERAC Heracleum 0–67 - - 1 dock RUMEX Rumex 0–67 - - 1 agwort </th		
1Perennial, rhizomatous grass168–336beardless wildryeLETR5Leymus triticoides168–3362Other grasslike135–538sedgeCAREXCarex67–2695Other perennial grasses67–269shortawn foxtailALAEAlopecurus aequalis0–67teal lovegrassERHYEragrostis hypnoides0–67basin wildryeLECl4Leymus cinereus0–67bluegrassPOAPoa0–67bluegrassPOAPoa0–67Forbsilverweed cinquefoilARAN7Argentina anserina0–67silverweed cinquefoilPOBI7Potentilla biennis0–67-dockRUMEXRumex0–67-gwortSENECSenecio0–67-ragwortSENECSenecio0–67-12Rhizomatous, deciduous, wetland obligate shrub673–1345-13Deciduous, wetland obligate shrub673–1345-14Deciduous, facultative wetland shrubSalix lucida ssp. lasiandria504–841-Pacific willowSALULSalix lucida ssp. lasiandria504–841-		
beardless wildryeLETR5Leymus triticoides168–336-2Other grasslike135–538sedgeCAREXCarex67–269-5Other perennial grasses67–269-shortawn foxtailALAEAlopecurus aequalis0–67-teal lovegrassERHYEragrostis hypnoides0–67-basin wildryeLEC14Leymus cinereus0–67-buegrassPOAPoa0–67-Forb9Other perennial forbs34–269silverweed cinquefoilARAN7Argentina anserina0–67cowparsnipHERACHeracleum0–67-biennial cinquefoilPOBI7Potentilla biennis0–67-dockRUMEXRumex0–67-ragwortSENECSenecio0–67-12Rhizomatous, deciduous, wetland obligate shrub673–1345-13Deciduous, wetland obligate shrub673–1345-14Deciduous, facultative wetland shrubSol4–841-14Deciduous, SALULSalix lucida ssp. lasiandra504–841-Pacific willowSALULSalix lucida ssp. lasiandra504–841-		
2Other grasslike135–538135–538sedgeCAREXCarex67–269 $-$ 5Other perennial grasse67–269 $-$ 135–538ALAEAlopecurus aequalis0–67 $-$ 14teal lovegrassERHYEragrostis hypnoides0–67 $-$ 15basin wildryeLECI4Leymus cinereus0–67 $-$ 15buegrassPOAPoa0–67 $-$ 16buegrassPOAPoa0–67 $-$ 17buegrassPOAPoa0–67 $-$ 18Verweed cinquefoilARAN7Argentina anserina0–67 $-$ 19Other perennial forbsVeracleum0–67 $-$ 10silverweed cinquefoilARAN7Argentina anserina0–67 $-$ 19Other perennial cinquefoilPOBI7Potentilla biennis0–67 $-$ 10biennial cinquefoilPOBI7Potentilla biennis0–67 $-$ 10dockRUMEXRumex0–67 $-$ 10dockRUMEXRumex0–67 $-$ 12Rhizomatous, deciduous, wetland obligate shrub673–1345 $-$ 13Deciduous, wetland obligate shrub673–1345 $-$ 13Deciduous, stalud size salux lutea504–841 $-$ 14Deciduous, facultative wetlant trubSalix lucida ssp. lasiandra504–841 $-$ 14DeciduousSALULSalix lucida ssp. lasiandra504–841<		
sedgeCAREXCarex $67-269$ $-$ 5Other perennial grasses $67-269$ $67-269$ shortawn foxtailALAEAlopecurus aequalis $0-67$ $-$ teal lovegrassERHYEragrostis hypnoides $0-67$ $-$ basin wildryeLECI4Leymus cinereus $0-67$ $-$ bluegrassPOAPoa $0-67$ $-$ Forb 9 Other perennial forbs $34-269$ $-$ silverweed cinquefoilARAN7Argentina anserina $0-67$ $-$ cowparsnipHERACHeracleum $0-67$ $-$ biennial cinquefoilPOBI7Potentilla biennis $0-67$ $-$ dockRUMEXRumex $0-67$ $-$ ragwortSENECSenecio $0-67$ $-$ Shrub/Vine $504-841$ $673-1345$ $-$ 13Deciduous, wetland \rightarrow bigate shrub $673-1345$ $-$ 14Deciduous, facultative wetland \rightarrow trub $504-841$ $-$ Pacific willowSALU2Salix lucia ssp. lasiandra $504-841$ $-$		
5Other perennial grasses $67-269$ shortawn foxtailALAEAlopecurus aequalis $0-67$ $-$ teal lovegrassERHYEragrostis hypnoides $0-67$ $-$ basin wildryeLECI4Leymus cinereus $0-67$ $-$ bluegrassPOAPoa $0-67$ $-$ Forb9Other perennial forbssilverweed cinquefoilARAN7Argentina anserina $0-67$ $-$ cowparsnipHERACHeracleum $0-67$ $-$ biennial cinquefoilPOBI7Potentilla biennis $0-67$ $-$ dockRUMEXRumex $0-67$ $-$ fagwortSENECSenecio $0-67$ $-$ 12Rhizomatous, decidu-s, wetlant obligate shrub $673-1345$ $-$ 13Deciduous, wetland obligate shrub $504-841$ $-$ 14Deciduous, facultative wetlant shrub $504-841$ $-$ Pacific willowSALULSalix lucida ssp. lasiandra $504-841$ $-$		
shortawn foxtailALAEAlopecurus aequalis $0-67$ $-$ teal lovegrassERHYEragrostis hypnoides $0-67$ $-$ bluegrassPOAPoa $0-67$ $-$ Forb POAPoa $0-67$ $-$ 9Other perennial forbs $34-269$ $-$ silverweed cinquefoilARAN7Argentina anserina $0-67$ $-$ cowparsnipHERACHeracleum $0-67$ $-$ biennial cinquefoilPOBI7Potentilla biennis $0-67$ $-$ dockRUMEXRumex $0-67$ $-$ fragwortSENECSenecio $0-67$ $-$ shizomatous, deciduus, wettand $0-67$ $ -$ 12Rhizomatous, deciduus, wettand obligate shrub $673-1345$ $-$ 13Deciduous, wettand obligate shrub $504-841$ $-$ 14Deciduous, facultative wetland shrub $504-841$ $-$ Pacific willowSALULSalix lucida ssp. lasiandra $504-841$ $-$		
teal lovegrassERHYEragrostis hypnoides0-67-basin wildryeLECI4Leymus cinereus0-67-bluegrassPOAPoa0-67-Forb9Other perennial forbs34-269silverweed cinquefoilARAN7Argentina anserina0-67-cowparsnipHERACHeracleum0-67-biennial cinquefoilPOBI7Potentilla biennis0-67-dockRUMEXRumex0-67-ragwortSENECSenecio0-67-ShrubVine12Rhizomatous, deciducus, wetland obligate shrub673-1345-13Deciduous, wetland obligate shrub504-841-14Deciduous, facultative wetland shrub504-841-Pacific willowSALULSalix lucida ssp. lasiandra504-841-		
basin wildryeLECl4Leymus cinereus $0-67$ $-$ bluegrassPOAPoa $0-67$ $-$ Forb9Other perennial forbs $34-269$ $silverweed cinquefoilARAN7Argentina anserina0-67 cowparsnipHERACHeracleum0-67 biennial cinquefoilPOBI7Potentilla biennis0-67 dockRUMEXRumex0-67 ragwortSENECSenecio0-67-StrubVine12Rhizomatous, deciduus, wetland obligate shrub673-1345-13Deciduous, wetland bigate shrub504-841-14Deciduous, facultative wetland shrub504-841-Pacific willowSALULSalix lucida ssp. lasiandra504-841-$		
bluegrassPOAPoa0-67-Forb9Other perennial forbs34-269silverweed cinquefoilARAN7Argentina anserina0-67-cowparsnipHERACHeracleum0-67-biennial cinquefoilPOBI7Potentilla biennis0-67-dockRUMEXRumex0-67-dockRUMEXRumex0-67-ragwortSENECSenecio0-67-Shrub/Vine12Rhizomatous, deciduus, wetland obligate shrub673-134513Deciduous, wetland obligate shrub504-841-14Deciduous, facultative wetland shrub504-841-14Pacific willowSALULSalix luciaa ssp. lasiandra504-841-		
Forb 9 Other perennial forbs 34–269 silverweed cinquefoil ARAN7 Argentina anserina 0–67 – cowparsnip HERAC Heracleum 0–67 – biennial cinquefoil POBI7 Potentilla biennis 0–67 – dock RUMEX Rumex 0–67 – ragwort SENEC Senecio 0–67 – Shrub/Vine SENEC Senecio 0–67 – 12 Rhizomatous, deciduus, wetland obligate shrub 673–1345 – 13 Deciduous, wetland obligate shrub 504–841 – 14 Deciduous, facultative wetland shrub 504–841 – Pacific willow SALU2 Salix lucida ssp. lasiandra 504–841 –		
9 Other perennial forbs 34-269 silverweed cinquefoil ARAN7 Argentina anserina 0-67 - cowparsnip HERAC Heracleum 0-67 - biennial cinquefoil POBI7 Potentilla biennis 0-67 - dock RUMEX Rumex 0-67 - ragwort SENEC Senecio 0-67 - Shrub/Vine 5ENEC Senecio 0-67 - 12 Rhizomatous, deciduus, wetland obligate shrub 673-1345 - 13 Deciduous, wetland obligate shrub 504-841 - 14 Deciduous, facultative wetland shrub 504-841 - Pacific willow SALUL Salix lucida ssp. lasiandra 504-841 -		
silverweed cinquefoilARAN7Argentina anserina0–67–cowparsnipHERACHeracleum0–67–biennial cinquefoilPOBI7Potentilla biennis0–67–dockRUMEXRumex0–67–ragwortSENECSenecio0–67–Shrub/Vine12Rhizomatous, deciduous, wetland obligate shrub673–134513Deciduous, wetland obligate shrub673–1345–13Deciduous, wetland obligate shrub504–841–14Deciduous, facultative wetland shrub504–841–Pacific willowSALULSalix lucida ssp. lasiandra504–841–		
cowparsnipHERACHeracleum0–67–biennial cinquefoilPOBI7Potentilla biennis0–67–dockRUMEXRumex0–67–ragwortSENECSenecio0–67–Shrub/Vine12Rhizomatous, deciduous, wetland obligate shrub673–1345narrowleaf willowSAEXSalix exigua673–1345–13Deciduous, wetland obligate shrub504–841–14Deciduous, facultative wetland shrub504–841–Pacific willowSALULSalix lucida ssp. lasiandra504–841–		
biennial cinquefoilPOBI7Potentilla biennis0–67–dockRUMEXRumex0–67–ragwortSENECSenecio0–67–Shrub/Vine12Rhizomatous, deciduous, wetland obligate shrub673–134513Deciduous, wetland obligate shrub504–841–14Deciduous, facultative wetland sp. lasiandra504–841–14Pacific willowSALULSalix lucida ssp. lasiandra504–841–		
dockRUMEXRumex0–67–ragwortSENECSenecio0–67–Shrub/Vine12Rhizomatous, deciduous, wetland obligate shrub673–134512narrowleaf willowSAEXSalix exigua673–134513Deciduous, wetland obligate shrub504–841–13yellow willowSALU2Salix lutea504–84114Deciduous, facultative wetland shrub504–841–14Pacific willowSALULSalix lucida ssp. lasiandra504–841		
ragwortSENECSenecio0–67–Shrub/Vine12Rhizomatous, deciduous, wetland obligate shrub673–134512narrowleaf willowSAEXSalix exigua13Deciduous, wetland obligate shrub504–84113yellow willowSALU2Salix lutea14Deciduous, facultative wetland shrub504–84114Pacific willowSALULSalix lucida ssp. lasiandra504–841–		
Shrub/Vine 12 Rhizomatous, deciduous, wetland obligate shrub 673–1345 narrowleaf willow SAEX Salix exigua 673–1345 – 13 Deciduous, wetland obligate shrub 504–841 – 14 Deciduous, facultative wetland shrub 504–841 – 14 Deciduous, facultative shrub 504–841 – 14 Deciduous, facultative shrub 504–841 –		
12 Rhizomatous, deciduous, wetland obligate shrub 673–1345 12 narrowleaf willow SAEX Salix exigua 673–1345 - 13 Deciduous, wetland obligate shrub 504–841 - - 14 Deciduous, facultative wetland shrub 504–841 - 14 Deciduous, facultative wetland shrub 504–841 - 14 Deciduous, facultative wetland shrub 504–841 -		
narrowleaf willow SAEX Salix exigua 673–1345 – 13 Deciduous, wetland obligate shrub 504–841 – 14 Deciduous, facultative wetland shrub 504–841 – 14 Deciduous, facultative wetland shrub 504–841 – 14 Deciduous, facultative wetland shrub 504–841 –		
13 Deciduous, wetland obligate shrub 504–841 yellow willow SALU2 Salix lutea 504–841 - 14 Deciduous, facultative wetland shrub 504–841 - Pacific willow SALUL Salix lucida ssp. lasiandra 504–841 -		
yellow willow SALU2 Salix lutea 504–841 – 14 Deciduous, facultative wetland shrub 504–841 – Pacific willow SALUL Salix lucida ssp. lasiandra 504–841 –		
14 Deciduous, facultative wetland shrub 504–841 Pacific willow SALUL Salix lucida ssp. lasiandra 504–841 –		
Pacific willow SALUL Salix lucida ssp. lasiandra 504–841 –		
15 Other shrubs 34–269		
gray alder ALIN2 Alnus incana 0–34 –		
golden currantRIAURibes aureum0-34-		
Woods' rose ROWO Rosa woodsii 0-34 -		
red elderberry SARA2 Sambucus racemosa 0-34 -		
Tree		
18 Deciduous 0-34		
balsam poplar POBA2 Populus balsamifera 0–34 –		

Animal community

Livestock Grazing:

This site is suited to summer, fall, and winter use by cattle and horses under a planned grazing system. The key species is willow. Palatable shrubs such as willow can be damaged if heavily grazed during periods when the herbaceous species are mature or dormant. Grazing prescriptions should be based on both vegetative and stream function goals.

Native Wildlife Associated with the Potential Climax Community:

Mammals and a variety of waterfowl and upland birds including savannah sparrows, bobolinks, and meadowlarks utilize this site for food and cover. It provides a high degree of diversity and high quality habitat for a large variety of

wildlife. Cover and habitat value is excellent when the ecological condition is high. The value of the site increases when it is near wet marshes, perennial streams, and open water areas.

Hydrological functions

The hydrologic cover condition is excellent when the ecological condition is high.

Other information

In fair condition with a remnant stand of willow, this site rapidly responds to good management.

Contributors

Alan V. Bahn Bob Gillaspy M. Parks (OSU)

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Jeff Repp
Contact for lead author	Oregon NRCS State Rangeland Management Specialist
Date	08/09/2012
Approved by	Bob Gillaspy
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills: None, moderate sheet & rill erosion hazard
- 2. **Presence of water flow patterns:** None to some. With reduced vegetative cover streambakns become unstable, streams become incised and groundwater tables are lowered.
- 3. Number and height of erosional pedestals or terracettes: None
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground): 0-5%
- 5. Number of gullies and erosion associated with gullies: None

- 6. Extent of wind scoured, blowouts and/or depositional areas: None, moderate wind erosion hazard
- 7. Amount of litter movement (describe size and distance expected to travel): Fine to coarse limited movement
- Soil surface (top few mm) resistance to erosion (stability values are averages most sites will show a range of values): Moderately resistant to erosion: aggregate stability = 3-5
- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Very deep somewhat poorly drained loams: Moderate to high OM (3-6%)
- Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: Significant ground cover (80-120%) and gentle slopes (0-3%) significantly limit rainfall impact and overland flow
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): None
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: Coyote willow > Yellow willow = Pacific willow > Creeping wildrye > Baltic rush = sedges = other grasses & grass-likes > other shrubs = forbs

Sub-dominant:

Other:

Additional:

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Normal decadence and mortality expected
- 14. Average percent litter cover (%) and depth (in):
- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annualproduction): Favorable: 4000, Normal: 3000, Unfavorable: 2000 lbs/acre/year at high RSI (HCPC)
- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if

their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Perennial brush species will increase with deterioration of plant community. Cheatgrass and Medusahead invade sites that have lost deep rooted perennial grass functional groups.

17. Perennial plant reproductive capability: All species should be capable of reproducing annually