

## Ecological site R025XY002ID STONY LOAM 16-22

Last updated: 4/24/2024  
Accessed: 05/06/2024

---

### Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	USDA/Natural Resources Conservation Service 9173 W. Barnes Drive, Suite C Boise, ID 83709 (208) 378-5700
Date	08/07/2007
Approved by	Kendra Moseley
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### Indicators

- 1. Number and extent of rills:** Rills are common on this site. If rills are present, they are likely to occur on slopes greater than 20 percent and immediately following wildfire. Rills are most likely to occur on soils with surface textures of silty clay loam.

---
- 2. Presence of water flow patterns:** Water-flow patterns are common on this site. When they occur, they are short and disrupted by cool season grasses, tall shrubs and surface cobbles. They are not extensive.

---
- 3. Number and height of erosional pedestals or terracettes:** Pedestals and/or terracettes occur on this site. In areas where slopes approach 20 percent and where flow patterns and/or rills are present, few pedestals may be expected. Terracettes are also common, especially on slopes greater than 20 percent.

---
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** On sites in mid-seral status, bare ground may range from 25-45 percent.

---
- 5. Number of gullies and erosion associated with gullies:** Gullies do not occur on this site.

- 
6. **Extent of wind scoured, blowouts and/or depositional areas:** Wind-scoured, blowouts and/or deposition areas usually not present. Immediately following wildfire, some soil movement may occur on lighter textured soils.
- 
7. **Amount of litter movement (describe size and distance expected to travel):** Fine litter in the interspaces may move up to 3 feet following a significant run-off event. Coarse litter generally does not move.
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Values should range from 4 to 6.
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** There is no data available on thickness of the A or A1 horizon. No data available on structure ranges. Soil organic matter (SOM) needs to be determined.
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Bunchgrasses, especially deep-rooted perennials, slow run-off and increase infiltration. Tall shrubs accumulate snow in the interspaces.
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** Compaction layer is not present.
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Cool season, deep-rooted perennial bunchgrasses>>tall shrubs.
- Sub-dominant: Perennial forbs>shallow rooted bunchgrasses.
- Other:
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Mountain big sagebrush and antelope bitterbrush will become decadent in the absence of fire and ungulate grazing. Grass and forb mortality will occur as tall shrubs increase.
- 
14. **Average percent litter cover (%) and depth ( in):** Additional litter cover data is needed but is expected to be 20-25 percent to a depth of 0.2 inches. Under mature shrubs, litter is >0.5 inches deep and is 90-100 percent ground cover.
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** Annual Production is 1500 pounds per acre (1670Kg/ha) in a year with normal precipitation and temperatures. Perennial grasses produce 65 percent of the total production, forbs 10 percent and shrubs 25 percent.
- 

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Invasive plants include cheatgrass, bulbous bluegrass, whitetop, rush skeletonweed, musk and scotch thistle and diffuse and spotted knapweed. In addition, Utah juniper and pinyon pine can invade on this site.
- 

17. **Perennial plant reproductive capability:** All functional groups have the potential to reproduce in most years.
-