

## **Ecological site R025XY032OR NORTH SLOPES 11-13 PZ**

Last updated: 4/25/2024  
Accessed: 05/05/2024

---

### **General information**

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### **MLRA notes**

Major Land Resource Area (MLRA): 025X–Owyhee High Plateau

The Owyhee High Plateau, MLRA 25, lies within the Intermontane Plateaus physiographic province. The southern half is found in the Great Basin while the northern half is located in the Columbia Plateaus. The southern section of the Owyhee High Plateau is characterized by isolated, uplifted fault-block mountain ranges separated by narrow, aggraded desert plains. This geologically older terrain has been dissected by numerous streams draining to the Humboldt River. The northern section forms the southern boundary of the extensive Columbia Plateau basalt flows. Deep, narrow canyons drain to the Snake River across the broad volcanic plain.

This MLRA is characteristically cooler and wetter than the neighboring MLRAs of the Great Basin. Elevation ranges from 3,000 to 7,550 feet on rolling plateaus and in gently sloping basins. It is more than 9,840 feet on some steep mountains. The average annual precipitation in most of this area is typically 11 to 22 inches. It increases to as much as 49 inches at the higher elevations. Precipitation occurs mainly as snow in winter. The supply of water from precipitation and streamflow is small and unreliable, except along major rivers. Streamflow depends largely on accumulated snow in the mountains.

The dominant soil orders in this MLRA are Aridisols and Mollisols. The soils in the area dominantly have a mesic or frigid temperature regime and an aridic, arid bordering on xeric, or xeric moisture regime. Most of the soils formed in mixed parent material. Volcanic ash and loess mantle the landscape. Surface soil textures are loam and silt loam, and have ashy texture modifiers in some cases. Argillic horizons occur on the more stable landforms.

### **Ecological site concept**

This ecological site is on north facing side slopes of plateaus and tablelands associated with volcanic plateau landscapes. Elevations range from 2,700 to 5,000 feet (823 to 1,524 meters) and slopes range from 12 to 70 percent.

The soils associated with this site are moderately deep to bedrock. The soil climate is mesic to mesic near frigid (soil temperature regime) and aridic bordering xeric (soil moisture regime). Since this site is on north aspects, it receives less solar insolation and is slightly wetter and colder than its non-aspect counterpart, resulting in increased resistance and resilience, and annual production.

The reference plant community is characterized by basin big sagebrush and Idaho fescue, bluebunch wheatgrass is also common in the understory.

### **Associated sites**

R025XY012OR	<b>LOAMY 11-13 PZ</b>
R025XY020OR	<b>SOUTH SLOPES 11-13 PZ</b>
R025XY038OR	<b>CLAYPAN NORTH SLOPES 11-13 PZ</b>

## Similar sites

R025XY034OR	<b>SHRUBBY NORTH SLOPES 13-16 PZ</b> Cooler and wetter; higher production; increased prevalence of bitterbrush and mountain big sagebrush; bedrock is more fractured, possibly different geologies - more likely to occur on paralithic than lithic soils.
-------------	---

**Table 1. Dominant plant species**

Tree	Not specified
Shrub	(1) <i>Artemisia tridentata subsp. tridentata</i>
Herbaceous	(1) <i>Festuca idahoensis</i> (2) <i>Pseudoroegneria spicata subsp. spicata</i>

## Physiographic features

This ecological site is associated with north facing canyon side slopes and tablelands in lava plateau landscapes. Slopes range from 12 to 70 percent with elevations of 2,700 to 5,000 feet (823 to 1,524 meters).

**Table 2. Representative physiographic features**

Landforms	(1) Lava plateau > Mountain slope
Runoff class	Medium to high
Flooding frequency	None
Ponding frequency	None
Elevation	2,700–5,000 ft
Slope	12–70%
Water table depth	100 in
Aspect	NW, N, NE


## Climatic features

The climate associated with this site is defined by hot dry summers and cold snowy winters. This site is characterized by less than 120 freeze-free days annually. Mean annual precipitation is 10 inches (25 cm), with effective precipitation between 11 to 13 inches (28 to 33 cm). Average snowfall is between 25 to 50 inches (63 to 127 cm) per year.


\*The above data was provided by the Danner, Rockville, and Mc Dermitt climate station based on elevation and precipitation. Additionally used to estimate climate normals is the National Soil Information System.

**Table 3. Representative climatic features**


Frost-free period (characteristic range)	70-110 days
Freeze-free period (characteristic range)	60-121 days
Precipitation total (characteristic range)	10-11 in
Frost-free period (actual range)	30-120 days
Freeze-free period (actual range)	40-129 days
Precipitation total (actual range)	9-11 in
Frost-free period (average)	80 days
Freeze-free period (average)	90 days
Precipitation total (average)	10 in




**Figure 1. Monthly precipitation range**



**Figure 2. Monthly minimum temperature range**



**Figure 3. Monthly maximum temperature range**



**Figure 4. Monthly average minimum and maximum temperature**




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

### Climate stations used

- (1) DANNER [USC00352135], Jordan Valley, OR
- (2) ROCKVILLE 5 N [USC00357277], Adrian, OR
- (3) MC DERMITT 26 N [USC00355335], Jordan Valley, OR

### Influencing water features

This site is not influenced by adjacent wetlands, streams or run-on.

### Wetland description

N/A

### Soil features

The soils associated with this site are well drained and formed in residuum, colluvium, slope alluvium derived from volcanic rock and ash. Surface textures are typically loamy. The soil profile is characterized by a dark surface horizon (mollic epipedon) and a layer of clay accumulation (argillic horizon) within 20 inches (51cm) of the soil surface. Soil are moderately deep with bedrock, ranging from weathered andesite to unweathered rhyolite, occurring within 40 inches (102cm) of the surface.

Representative soil components associated with this ecological site include Perla and Lerrow.

Table 4. Representative soil features

Parent material	(1) Colluvium–volcanic rock (2) Residuum–volcanic rock (3) Alluvium–volcanic rock
Surface texture	(1) Loam (2) Gravelly loam

Family particle size	(1) Fine
Drainage class	Well drained
Permeability class	Slow to moderately slow
Soil depth	18–40 in
Surface fragment cover ≤3"	5–15%
Surface fragment cover >3"	0–1%
Available water capacity (0–40in)	1.7–3.7 in
Soil reaction (1:1 water) (Depth not specified)	6.1–7.3
Subsurface fragment volume ≤3" (Depth not specified)	10–15%
Subsurface fragment volume >3" (Depth not specified)	5–15%

## Ecological dynamics

The reference plant community is dominated by basin big sagebrush with an understory of Idaho fescue. Bluebunch wheatgrass is also common in the understory. The site has moderately low resilience to disturbance and resistance to invasion. Resilience is a system's capacity to regain its structure, processes, and function following stressors or disturbance (e.g. drought or fire). Resistance is the capacity of the system to retain its structure, processes, and function despite stressors or disturbances (including pressure from invasive species) (Chambers 2014). Resilience increases with elevation, aspect, increased precipitation and increased nutrient availability (Stringham et al. 2015); where greater resource availability and more favorable environmental conditions exist for plant growth and reproduction (Chambers 2014).

This ecological site's lower effective precipitation (aridic bordering xeric soil moisture regime) limits site productivity; though since the site occurs on north aspects, it receives less solar insolation and thus is slightly wetter and colder than its non-aspect counterpart. The higher effective precipitation experienced on north slopes results in higher annual production and less open space for establishment of invasive annual grasses, increasing site resilience. The timing of precipitation on this site still favors invasive annual grasses that are particularly well adapted to cool wet winters and warm dry summers; beginning growth and utilizing resources prior to native species breaking dormancy. The site's warm soil temperature regime (mesic to mesic near frigid) gives the site moderately low resistance to disturbance, though being north aspect slightly increases resistance compared to a non-aspect site.

This ecological site is dominated by deep-rooted cool season, perennial bunchgrasses and long-lived shrubs (50+ years) with high root to shoot ratios. The dominant shrubs usually root to the full depth of the winter-spring soil moisture recharge, which ranges from 1.0 to over 3.0 m (Dobrowolski et al. 1990). Root length of mature sagebrush plants was measured to a depth of 2 meters in alluvial soils in Utah (Richards and Caldwell 1987). These shrubs have a flexible generalized root system with development of both deep taproots and laterals near the surface (Comstock and Ehleringer 1992).

Periodic drought regularly influences sagebrush ecosystems and drought duration and severity has increased throughout the 20th century in much of the Intermountain West. Major shifts away from historical precipitation patterns have the greatest potential to alter ecosystem function and productivity. Species composition and productivity can be altered by the timing of precipitation and water availability with the soil profile (Bates et al. 2006).

Big sagebrush and antelope bitterbrush are generally long-lived; therefore it is not necessary for new individuals to recruit every year for perpetuation of the stand. Infrequent large recruitment events and simultaneous low, continuous recruitment is the foundation of population maintenance (Noy-Meir 1973). Survival of the seedlings is dependent on adequate moisture conditions.

In the Great Basin, the majority of annual precipitation is received during winter and early spring. This continental semiarid climate regime favors growth and development of deep-rooted shrubs and herbaceous cool season plants

using the C3 photosynthetic pathway (Comstock and Ehleringer 1992). Winter precipitation and slow melting of snow results in deeper percolation of moisture into the soil profile. Herbaceous plants, more shallow-rooted than shrubs, grow earlier in the growing season and thrive on spring rains, while the deeper rooted shrubs lag in phenological development because they draw from deeply infiltrating moisture from snowmelt the previous winter.

Native insect outbreaks are also important drivers of ecosystem dynamics in sagebrush communities. Climate is generally believed to influence the timing of insect outbreaks especially with regard to Aroga moth (*Aroga websteri*), a sagebrush defoliator. Aroga moth infestations have occurred in the Great Basin in the 1960s, early 1970s, and have been ongoing in Nevada since 2004 (Bentz, et al 2008). Thousands of acres of big sagebrush have been impacted, with partial to complete die-off observed. Aroga moth can partially or entirely kill individual plants or entire stands of big sagebrush (Furniss and Barr 1975).

The Great Basin sagebrush communities have high spatial and temporal variability in precipitation both among years and within growing seasons. Nutrient availability is typically low but increases with elevation and closely follows moisture availability. The invasibility of plant communities is often linked to resource availability. Disturbance can decrease resource uptake due to damage or mortality of the native species and depressed competition or can increase resource pools by the decomposition of dead plant material following disturbance. The invasion of sagebrush communities by cheatgrass (*Bromus tectorum*) has been linked to disturbances (fire, abusive grazing) that have resulted in fluctuations in resources (Chambers et al. 2007).

The introduction of annual weedy species, like cheatgrass, may cause an increase in fire frequency and eventually lead to an annual state. Conversely, as fire frequency decreases, sagebrush will increase and with inappropriate grazing management the perennial bunchgrasses and forbs may be reduced.

#### Fire Ecology:

Big sagebrush is killed by fire and does not resprout (Blaisdell 1953). Post fire regeneration occurs from seed and will vary depending on site characteristics, seed source, and fire characteristics. Big sagebrush reinvades a site primarily by off-site seed or seed from plants that survive in unburned patches. Approximately 90% of big sagebrush seed is dispersed within 30 feet (9 m) of the parent shrub (Goodrich et al. 1985) with maximum seed dispersal at approximately 108 feet (33 m) from the parent shrub (Shumar and Anderson 1986). Mountain big sagebrush seedlings can grow rapidly and may reach reproductive maturity within 3 to 5 years (Bunting et al. 1987), while basin big sagebrush tends to be slower to regenerate, and it is often eliminated by frequent fires (Bunting et al. 1987). Regeneration of basin big sagebrush after stand replacing fires is difficult and dependent upon proximity of residual mature plants and favorable moisture conditions (Johnson and Payne 1968, Humphrey 1984).

The effect of fire on bunchgrasses relates to culm density, culm-leaf morphology, and the size of the plant. The initial condition of bunchgrasses within the site along with seasonality and intensity of the fire all factor into the individual species response. For most forbs and grasses the growing points are located at or below the soil surface providing relative protection from disturbances which decrease above ground biomass, such as grazing or fire. Thus, fire mortality is more correlated to duration and intensity of heat which is related to culm density, culm-leaf morphology, size of plant and abundance of old growth (Wright 1971, Young 1983).

Idaho fescue response to fire varies with condition and size of the plant, season and severity of fire, and ecological conditions. Mature Idaho fescue plants are commonly reported to be severely damaged by fire in all seasons (Wright et al. 1979). Initial mortality may be high (in excess of 75%) on severe burns, but usually varies from 20 to 50% (Barrington et al 1988). Rapid burns have been found to leave little damage to root crowns, and new tillers are produced with onset of fall moisture (Johnson et al. 1994). However, Wright and others (1979) found the dense, fine leaves of Idaho fescue provided enough fuel to burn for hours after a fire had passed, thereby killing or seriously injuring the plant regardless of the intensity of the fire (Wright et al. 1979).


Fire will remove aboveground biomass from bluebunch wheatgrass but plant mortality is generally low (Robberecht and Defossé 1995) because the buds are underground (Conrad and Poulton 1966) or protected by foliage. Uresk et al. (1976) reported burning increased vegetative and reproductive vigor of bluebunch wheatgrass. Thus, bluebunch wheatgrass is considered to experience slight damage to fire but is more susceptible in drought years (Young 1983). Plant response will vary depending on season, fire severity, fire intensity and post-fire soil moisture availability.

Sandberg bluegrass, a minor component of this ecological site, has been found to increase following fire likely due

to its low stature and productivity (Daubenmire 1975). Sandberg bluegrass may retard reestablishment of deeper rooted bunchgrass. Reduced bunchgrass vigor or density provides an opportunity for Sandberg bluegrass expansion and/or cheatgrass and other invasive species to occupy interspaces, leading to increased fire frequency and potentially an annual plant community.

Adapted from: Stringham, T.K., P. Novak-Echenique, P. Blackburn, D. Snyder, and A. Wartgow. 2015. Final Report for USDA Ecological Site Description State-and-Transition Models by Disturbance Response Groups, Major Land Resource Area 25 Nevada. University of Nevada Reno, Nevada Agricultural Experiment Station Research Report 2015-02. p. 569

## State and transition model



Reference State 1.0 Community Phase Pathways

- 1.1a: Low severity fire creates grass/sagebrush mosaic; high severity fire significantly reduces bitterbrush/sagebrush cover and leads to early/mid-seral community, dominated by grasses and forbs
- 1.1b: Time and lack of disturbance such as fire or drought. Excessive herbivory may also decrease perennial understory.
- 1.2a: Time and lack of disturbance allows for shrub regeneration.
- 1.3a: Low severity fire or Aroga moth infestation resulting in a mosaic pattern.
- 1.3b: High severity fire or Aroga moth significantly reduces sagebrush cover leading to early/mid-seral community.

Transition T1A: Introduction of non-native species.

Current Potential State 2.0 Community Phase Pathways

- 2.1a: Low severity fire creates grass/sagebrush mosaic; high severity fire significantly reduces sagebrush cover and leads to early/mid-seral community dominated by grasses and forbs; non-native annual species present.
- 2.1b: Time and lack of disturbance such as fire or drought. Inappropriate grazing management may also reduce perennial understory.
- 2.2a: Time and lack of disturbance allows for regeneration of sagebrush.
- 2.3a: Low severity fire or Aroga moth infestation creates shrub/grass mosaic. Brush treatment with minimal soil disturbance; late-fall/winter grazing causing mechanical damage to shrubs.
- 2.3b: High severity fire or Aroga moth significantly reduces shrub cover leading to a early/mid-seral community.

Transition T2A: Inappropriate grazing management favoring shrub dominance and reducing perennial bunchgrasses will lead to phase 3.1. Soil disturbing treatments will lead to phase 3.2.

Transition T2B: Catastrophic fire (to 4.1); inappropriate cattle/horse grazing management that removes bunchgrasses, favors shrubs and promotes the presence of non-native annual species (to 4.2)

Shrub State 3.0 Community Phase Pathways

- 3.1a: Low severity fire or Aroga moth infestation creates shrub/grass mosaic. Brush treatment with minimal soil disturbance; late-fall/winter grazing causing mechanical damage to shrubs.
- 3.2a: Time and lack of disturbance.

Restoration R3A: Brush management and seeding of native deep rooted bunchgrasses (probability of success is low).

Transition T3A: Fire and/or soil disturbing treatments.

Annual State 4.0 Community Phase Pathways

- 4.1a: Time and lack of disturbance. Big sagebrush is unlikely to reestablish and may take many years.
- 4.2a: High-severity fire.

(Adapted from Stringham, T.K. et al., 2015)

## State 1 Reference State

The Reference State 1.0 is a representative of the natural range of variability under pristine conditions. The reference state has three general community phases; a shrub-grass dominant phase, a perennial grass dominant phase and a shrub dominant phase. State dynamics are maintained by interactions between climatic patterns and disturbance regimes. Negative feedbacks enhance ecosystem resilience and contribute to the stability of the state. These include the presence of all structural and functional groups, low fine fuel loads, and retention of organic matter and nutrients. Plant community phase changes are primarily driven by fire, periodic drought and/or insect or disease attack.

**Characteristics and indicators.** Production will increase at the upper end of the precipitation zone and on deeper foot slope soils. Bluebunch wheatgrass increases on more westerly or easterly aspects. As a fire susceptible site, the amount of basin big sagebrush is influenced by fire frequency.

### Dominant plant species

- basin big sagebrush (*Artemisia tridentata* ssp. *tridentata*), shrub
- Idaho fescue (*Festuca idahoensis*), grass
- bluebunch wheatgrass (*Pseudoroegneria spicata* ssp. *spicata*), grass

## Community 1.1



Basin big sagebrush and Idaho fescue dominate the site. Bluebunch wheatgrass, other perennial grasses, and perennial forbs are also common on this site.

### Dominant plant species

- basin big sagebrush (*Artemisia tridentata* ssp. *tridentata*), shrub
- Idaho fescue (*Festuca idahoensis*), grass
- bluebunch wheatgrass (*Pseudoroegneria spicata* ssp. *spicata*), grass

Table 5. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	1050	1225	1400
Shrub/Vine	90	105	120
Forb	60	70	80
<b>Total</b>	<b>1200</b>	<b>1400</b>	<b>1600</b>

## Community 1.2

This community phase is characteristic of a post-disturbance, early to mid-seral community phase. Idaho fescue, bluebunch wheatgrass and other perennial grasses dominate. Rabbitbrush and other sprouting shrubs may be resprouting. Big sagebrush is killed by fire, therefore decreasing within the burned community. Depending on fire severity or intensity of Aroga moth infestations, patches of intact sagebrush may remain. Perennial forbs may increase post-fire but will likely return to pre-burn levels within a few years.

### Dominant plant species

- Idaho fescue (*Festuca idahoensis*), grass
- bluebunch wheatgrass (*Pseudoroegneria spicata* ssp. *spicata*), grass

## Community 1.3

Basin big sagebrush increases in the absence of disturbance. Decadent shrubs dominate the overstory and the deep-rooted perennial bunchgrasses in the understory are reduced either from competition with shrubs or from herbivory. Sandberg bluegrass will likely increase in the understory and may be the dominant grass on the site.

### Dominant plant species

- basin big sagebrush (*Artemisia tridentata* ssp. *tridentata*), shrub
- Idaho fescue (*Festuca idahoensis*), grass
- bluebunch wheatgrass (*Pseudoroegneria spicata* ssp. *spicata*), grass
- Sandberg bluegrass (*Poa secunda*), grass

## Pathway 1.1a

### Community 1.1 to 1.2

Fire decreases or eliminates the overstory of sagebrush and allows for the perennial bunchgrasses to dominate the site. Fires would typically be small and patchy due to low fuel loads. A fire following an unusually wet spring or a change in management may be more severe and reduce shrub cover to trace amounts. A severe infestation of Aroga moth could also cause a large decrease in sagebrush within the community, giving a competitive advantage to the perennial grasses and forbs.

## Pathway 1.1b

### Community 1.1 to 1.3

Long-term drought, time and/or herbivory favor an increase in shrubs over deep-rooted perennial bunchgrasses. Combinations of these would allow the shrub overstory to increase and dominate the site, causing a reduction in the

perennial bunchgrasses.

### **Pathway 1.2a**

#### **Community 1.2 to 1.1**

Time, absence of disturbance and natural regeneration over time allows sagebrush to recover. Recovery of sagebrush depends on the availability of a local seed source (patches of mature shrubs) as well as precipitation patterns favorable for germination and seedling recruitment. Sagebrush seedlings are susceptible to less than favorable conditions for several years. Completion of this community phase pathways may take decades.

### **Pathway 1.3a**

#### **Community 1.3 to 1.1**

Low intensity, patchy wildfire or insect infestation would reduce sagebrush overstory creating a mosaic on the landscape. Perennial bunchgrasses and forbs dominate disturbed patches due to an increase in light, moisture and nutrient resources.

### **Pathway 1.3b**

#### **Community 1.3 to 1.2**

Fire decreases or eliminates the overstory of sagebrush and allows for the perennial bunchgrasses to dominate the site. Fires would typically be small and patchy due to low fuel loads. A fire following an unusually wet spring may be more severe and reduce shrub cover to trace amounts. A severe infestation of Aroga moth could also cause a large decrease in sagebrush within the community, giving a competitive advantage to the perennial grasses and forbs.

## **State 2**

### **Current Potential**

This state is similar to the Reference State 1.0. Ecological function has not changed, however the resiliency of the state has been reduced by the presence of invasive weeds. This state has the same three general community phases. Negative feedbacks enhance ecosystem resilience and contribute to the stability of the state. These include the presence of all structural and functional groups, low fine fuel loads and retention of organic matter and nutrients. Positive feedbacks decrease ecosystem resilience and stability of the state. These include the non-natives high seed output, persistent seed bank, rapid growth rate, ability to cross pollinate and adaptations for seed dispersal. Additionally, the presence of highly flammable, non-native species reduces State resilience because these species can promote fire where historically fire has been infrequent leading to positive feedbacks that further the degradation of the system.

**Characteristics and indicators.** The introduction of annual weedy species, like cheatgrass, may cause an increase in fire frequency. Without targeted management actions state is at-risk of crossing a threshold to an annual dominated community.

**Resilience management.** Best management options would be to maintain high diversity of desired species to promote organic matter inputs and prevent the dispersal and seed production of the non-native invasive species. Inappropriate grazing management by livestock and feral horses will cause a decrease in deep-rooted perennial bunchgrasses, mainly Idaho fescue and bluebunch wheatgrass. Long-term inappropriate grazing management may result in an increase in Sandberg bluegrass (*Poa secunda*), balsamroot, lupine, sagebrush, and rabbitbrush (*Chrysothamnus viscidiflorus*).

### **Dominant plant species**

- basin big sagebrush (*Artemisia tridentata* ssp. *tridentata*), shrub
- bluebunch wheatgrass (*Pseudoroegneria spicata*), grass
- Idaho fescue (*Festuca idahoensis*), grass

## **Community 2.1**

Basin big sagebrush, and Idaho fescue dominate the site. Bluebunch wheatgrass, other perennial grasses, and

perennial forbs are also common on this site. Non-native annual species are present in minor amounts.

**Resilience management.** The presence of non-native annuals has reduced site resilience. Management actions should focus on maintaining the presence of all functional and structural groups and minimizing wildfire and soil disturbing practices.

## **Community 2.2**

This community phase is characteristic of a post-disturbance, early to mid-seral community phase. Idaho fescue, bluebunch wheatgrass, and other perennial grasses dominate. Rabbitbrush and other sprouting shrubs may be resprouting. Big sagebrush is killed by fire, therefore decreasing within the burned community. Depending on fire severity or intensity of Aroga moth infestations, patches of intact sagebrush may remain. Perennial forbs may increase post-fire but will likely return to pre-burn levels within a few years. Annual non-native species generally respond well after fire and may be stable or increasing within the community.

## **Community 2.3 (at risk)**

Basin big sagebrush increases in the absence of disturbance. Decadent shrubs dominate the overstory and the deep-rooted perennial bunchgrasses in the understory are reduced either from competition with shrubs or from herbivory. Sandberg bluegrass will likely increase in the understory and may be the dominant grass on the site. Annual non-native species present.

### **Pathway 2.1a Community 2.1 to 2.2**

Fire decreases or eliminates the overstory of sagebrush and allows for the perennial bunchgrasses to dominate the site. Fires would typically be small and patchy due to low fuel loads. A fire following an unusually wet spring may be more severe and reduce shrub cover to trace amounts. A severe infestation of Aroga moth could also cause a large decrease in sagebrush within the community, giving a competitive advantage to the perennial grasses and forbs.

### **Pathway 2.1b Community 2.1 to 2.3**

Time and lack of disturbance allows for sagebrush to increase and become decadent. Mature sagebrush is controlling the spatial and temporal distribution of moisture, nutrient and light resources. Native perennial bunchgrasses are reduced due to competition for these resources. Non-native annuals are stable to increasing.

### **Pathway 2.1a Community 2.2 to 2.1**

Time, lack of disturbance and natural regeneration of sagebrush. The establishment of sagebrush depends on presence of seed source and favorable weather patterns. It may take decades for sagebrush to recover to pre-disturbance levels.

### **Pathway 2.3a Community 2.3 to 2.1**

Low severity fire or Aroga moth infestation creates shrub/grass mosaic.

### **Pathway 2.3b Community 2.3 to 2.2**

Fire reduces the shrub overstory and allows for perennial bunchgrasses to dominate the site. Fires are typically low severity resulting in a mosaic pattern due to low fuel loads. A fire following an unusually wet spring favoring an increase in fine fuels may be more severe and reduce sagebrush cover to trace amounts. A severe infestation of Aroga moth could also cause a large decrease in sagebrush within the community, giving a competitive advantage

to the perennial grasses and forbs. Annual non-native species are likely to increase after fire.

### **State 3**

#### **Shrub State**

This state has two community phases; a sagebrush dominated phase and a sprouting shrub dominated phase. This state is a product of many years of heavy grazing during time periods harmful to perennial bunchgrasses. Sandberg bluegrass will increase with a reduction in deep rooted perennial bunchgrass competition and become the dominant grass. Sagebrush dominates the overstory and rabbitbrush may be a significant component. Sagebrush canopy cover is high and shrubs may be decadent, reflecting stand maturity and lack of seedling establishment due to competition with mature plants. The shrub overstory and Sandberg bluegrass understory dominate site resources such that soil water, nutrient capture, nutrient cycling and soil organic matter are temporally and spatially redistributed.

#### **Dominant plant species**

- cheatgrass (*Bromus tectorum*), grass

### **Community 3.1**

This site has crossed a biotic threshold and site processes (soil hydrology, nutrient cycling, and energy capture) are being controlled by the shrub component of the plant community along with Sandberg bluegrass in the understory. Decadent sagebrush dominates the overstory. Deep-rooted perennial bunchgrasses may be present in trace amounts or absent from the community. Sandberg bluegrass and annual non-native species increase, and the amount of bare ground increases.

### **Community 3.2**

Sandberg bluegrass dominates the understory; annual non-natives are present but are not dominant. Trace amounts of bitterbrush/sagebrush may be present. Sprouting shrubs dominate for a number of years following fire.

#### **Pathway 3.1a**

#### **Community 3.1 to 3.2**

Fire decreases or eliminates the overstory of sagebrush. A severe infestation of Aroga moth could also cause a large decrease in sagebrush within the community, giving a competitive advantage to the Sandberg bluegrass, forbs and sprouting shrubs. Heavy fall grazing causing mechanical damage to shrubs, and/or brush treatments with minimal soil disturbance, would greatly reduce the overstory shrubs and allow for Sandberg bluegrass to dominate the site.

#### **Pathway 3.2a**

#### **Community 3.2 to 3.1**

Absence of disturbance over time would allow for bitterbrush/sagebrush and other shrubs to recover.

### **State 4**

#### **Annual State**

This state has two community phases; one dominated by annual non-native species without shrubs, the other with a shrub overstory. Annual non-natives dominate site productivity and site resources. This state is characterized by the dominance of annual non-native species such as cheatgrass and tansy mustard in the understory. Sagebrush and/or rabbitbrush may dominate the overstory.

### **Community 4.1**

Annual non-native plants such as cheatgrass or tansy mustard dominate the site.

## **Community 4.2**

Sagebrush and/or sprouting shrubs remain in the overstory with annual non-native species, likely cheatgrass, dominating the understory. Trace amounts of desirable bunchgrasses may be present.

### **Pathway 4.1a**

#### **Community 4.1 to 4.2**

Time and lack of disturbance allows for shrubs to reestablish. Sprouting shrubs such as rabbitbrush will be the first to reappear after fire. Probability of sagebrush establishment is extremely low, especially if fire becomes frequent on the site.

### **Pathway 4.2a**

#### **Community 4.2 to 4.1**

Fire allows for annual non-native species to dominate site.

## **Transition T1A**

### **State 1 to 2**

Trigger: This transition is caused by the introduction of non-native species. Slow variables: Over time the annual non-native plants will increase within the community decreasing organic matter inputs from deep-rooted perennial bunchgrasses resulting in reductions in soil water availability for perennial bunchgrasses. Threshold: Any amount of introduced non-native species causes an immediate decrease in the resilience of the site. Annual non-native species cannot be easily removed from the system and have the potential to significantly alter disturbance regimes from their historic range of variation.

## **Transition T2A**

### **State 2 to 3**

Trigger: Inappropriate, long-term grazing of perennial bunchgrasses during growing season would favor shrubs and initiate transition to Community Phase 3.1. Severe fire would cause a transition to Community Phase 3.2. Slow variables: Long term decrease in deep-rooted perennial grass density resulting in a decrease in organic matter inputs and subsequent soil water decline. Threshold: Loss of deep-rooted perennial bunchgrasses changes spatial and temporal nutrient cycling and nutrient redistribution, and reduces soil organic matter.

## **Transition T2B**

### **State 2 to 4**

Trigger: Multiple fires lead to plant community phase 4.1. Inappropriate grazing management that favors shrubs in the presence of non-native annual species leads to community phase 4.2. Slow variables: Increased production and cover of non-native annual species. Threshold: Loss of deep-rooted perennial bunchgrasses and shrubs changes nutrient capture and cycling within the community. Increased, continuous fine fuels modify the fire regime by changing intensity, size and spatial variability of fires.

## **Restoration pathway R3A**

### **State 3 to 2**

Brush management, herbicide or sub-soiling of Sandberg bluegrass and seeding of desired perennial bunchgrass.

## **Transition T3A**

### **State 3 to 4**

Trigger: Severe fire will transition to 4.1. Inappropriate grazing management in the presence of annual non-native species will transition to 4.2. Slow variable: Increased seed production and cover of annual non-native species. Threshold: Increased, continuous fine fuels modify the fire regime by changing intensity, size and spatial variability of fires. Changes in plant community composition and spatial variability of vegetation due to the loss of perennial

bunchgrasses and sagebrush truncate energy capture and impact the nutrient cycling and distribution.

## Additional community tables

Table 6. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
<b>Grass/Grasslike</b>					
1	<b>Perennial Deep-Rooted Bunchgrasses</b>			994–1302	
	Idaho fescue	FEID	<i>Festuca idahoensis</i>	630–770	–
	bluebunch wheatgrass	PSSP6	<i>Pseudoroegneria spicata</i>	350–490	–
	Thurber's needlegrass	ACTH7	<i>Achnatherum thurberianum</i>	14–42	–
2	<b>Perennial Shallow-Rooted Bunchgrass</b>			28–70	
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	28–70	–
3	<b>Other Perennial Grasses</b>			0–84	
	squirreldail	ELEL5	<i>Elymus elymoides</i>	0–28	–
	basin wildrye	LECI4	<i>Leymus cinereus</i>	0–28	–
	Cusick's bluegrass	POCU3	<i>Poa cusickii</i>	0–28	–
<b>Forb</b>					
4	<b>Perennial Forbs</b>			28–56	
	milkvetch	ASTRA	<i>Astragalus</i>	14–28	–
	lupine	LUPIN	<i>Lupinus</i>	14–28	–
5	<b>Other Perennial Forbs</b>			0–56	
	desertparsley	LOMAT	<i>Lomatium</i>	0–14	–
	arrowleaf balsamroot	BASA3	<i>Balsamorhiza sagittata</i>	0–14	–
	fleabane	ERIGE2	<i>Erigeron</i>	0–14	–
	common yarrow	ACMI2	<i>Achillea millefolium</i>	0–14	–
	tapertip hawksbeard	CRAC2	<i>Crepis acuminata</i>	0–14	–
	phlox	PHLOX	<i>Phlox</i>	0–14	–
	buckwheat	ERIOG	<i>Eriogonum</i>	0–14	–
	stoneseed	LITHO3	<i>Lithospermum</i>	0–14	–
<b>Shrub/Vine</b>					
6	<b>Shrubs</b>			42–126	
	basin big sagebrush	ARTRT	<i>Artemisia tridentata ssp. tridentata</i>	28–98	–
	yellow rabbitbrush	CHVI8	<i>Chrysothamnus viscidiflorus</i>	14–28	–
7	<b>Other Shrubs</b>			0–42	
	antelope bitterbrush	PUTR2	<i>Purshia tridentata</i>	0–14	–
	rubber rabbitbrush	ERNA10	<i>Ericameria nauseosa</i>	0–14	–
	Wyoming big sagebrush	ARTRW8	<i>Artemisia tridentata ssp. wyomingensis</i>	0–14	–

## Animal community

This site is suitable for grazing. Grazing management considerations include timing, duration and intensity of grazing.

Overgrazing leads to an increase in sagebrush and a decline in understory plants like Idaho fescue and bluebunch

wheatgrass. Squirreldale or Sandberg bluegrass will increase temporarily with further degradation. Invasion of annual weedy forbs and cheatgrass could occur with further grazing degradation, leading to a decline in squirreldale and bluegrasses and an increase in bare ground. A combination of overgrazing and prolonged drought leads to soil erosion, increased bare ground and a loss in plant production.

Idaho fescue is valuable forage for livestock and wildlife. However, Idaho fescue decreases under heavy grazing by livestock (Eckert and Spencer 1986, Eckert and Spencer 1987) and wildlife (Gaffney 1941).

Bluebunch wheatgrass is moderately grazing tolerant and is very sensitive to defoliation during the active growth period (Blaisdell and Pechanec 1949, Laycock 1967, Anderson and Scherzinger 1975). Herbage and flower stalk production was reduced with clipping at all times during the growing season; however, clipping was most harmful during the boot stage (Blaisdell and Pechanec 1949, Britton et al. 1990) Tiller production and growth of bluebunch was greatly reduced when clipping was coupled with drought (Busso and Richards 1995). Mueggler (1975) estimated that low vigor bluebunch wheatgrass may need up to 8 years rest to recover. Although an important forage species, it is not always the preferred species by livestock and wildlife.

Reduced bunchgrass vigor or density provides an opportunity for Sandberg bluegrass expansion and/or cheatgrass and other invasive species such as halogeton (*Halogeton glomeratus*), bur buttercup (*Ceratocephala testiculata*) and annual mustards to occupy interspaces. Sandberg bluegrass increases under grazing pressure (Tisdale and Hironaka 1981) and is capable of co-existing with cheatgrass. Excessive sheep grazing favors Sandberg bluegrass; however, where cattle are the dominant grazers, cheatgrass often dominates (Daubenmire 1970). Thus, depending on the season of use, the grazer and site conditions, either Sandberg bluegrass or cheatgrass may become the dominant understory with inappropriate grazing management.

## Inventory data references

Old SS Manuscripts, Range Site Descriptions, etc.  
Vale District BLM Ecological Site Inventory  
NASIS component and pedon data  
Range Site Descriptions  
Field knowledge of range-trained personnel

## References

- Haubensak K. and D'Antonio C. 2009. Effects of fire and environmental variable on plant structure and composition in grazed salt desert shrublands of the Great Basin (USA). *Journal of Arid Environment*. Elsevier. 643–650.
- Michael J. Falkowski and Jeffrey S. Evans. January 2017. Mapping Tree Canopy Cover in Support of Proactive Prairie Grouse Conservation in Western North America. *Rangeland Ecology and Management* 70:15–24.
- Pilliod, D.S. and J.L. Welty. 2017. Refining the cheatgrass–fire cycle in the Great Basin: Precipitation timing and fine fuel composition predict wildfire trends. *Ecology and Evolution*. Wiley.
- Sapsis, D.B. and J.B. Kauffman. 1991. Fuel consumption and fire behavior associated with prescribed fires in sagebrush ecosystems. *Northwest Science* 65:173–179.
- Williams, C.J. and F.B. Pierson. 2018. Effectiveness of prescribed fire to re-establish sagebrush steppe vegetation and ecohydrologic function on woodland-encroached sagebrush rangelands, Great Basin, USA: Part I: Vegetation, hydrology, and erosion responses.
- Williamson, M.A. and E. Fleishman. 2019. Fire, livestock grazing, topography, and precipitation affect occurrence and prevalence of cheatgrass (*Bromus tectorum*) in the central Great Basin, USA.

## Other references

- Anderson, E. W. and R. J. Scherzinger. 1975. Improving quality of winter forage for elk by cattle grazing. *Journal of Range Management* 28:120-125.
- Barrington, M., S. Bunting, and G. Wright. 1988. A fire management plan for Craters of the Moon National Monument. Cooperative Agreement CA-9000-8-0005. Moscow, ID: University of Idaho, Range Resources Department. 52 p. Draft.
- Bates, J. D., T. Svejcar, R. F. Miller, and R. A. Angell. 2006. The effects of precipitation timing on sagebrush steppe vegetation. *Journal of Arid Environments* 64:670-697.
- Bentz, B., D. Alston, and T. Evans. 2008. Great Basin Insect Outbreaks. In: J. Chambers, N. Devoe, A. Evenden [eds]. Collaborative Management and Research in the Great Basin -- Examining the issues and developing a framework for action Gen. Tech. Rep. RMRS-GTR-204. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fort Collins, CO. p. 45-48
- Blaisdell, J. P. and J. F. Pechanec. 1949. Effects of Herbage Removal at Various Dates on Vigor of Bluebunch Wheatgrass and Arrowleaf Balsamroot. *Ecology* 30:298-305.
- Blaisdell, J. P. 1953. Ecological effects of planned burning of sagebrush-grass range on the upper Snake River Plains. US Dept. of Agriculture.
- Blaisdell, J. P. and W. F. Mueggler. 1956. Sprouting of bitterbrush (*Purshia tridentata*) following burning or top removal. *Ecology* 37:365-370.
- Blaisdell, J. P., R. B. Murray, and E. D. McArthur. 1982. Managing intermountain rangelands-sagebrush grass ranges. USDA Forest Serv. Intermountain Forest and Range Exp. Sta. Gen. Tech. Rep. INT-134.
- Britton, C. M., G. R. McPherson, and F. A. Sneva. 1990. Effects of burning and clipping on five bunchgrasses in eastern Oregon. *Great Basin Naturalist* 50:115-120.
- Bunting, S. C., B. M. Kilgore, and C. L. Bushey. 1987. Guidelines for prescribed burning sagebrush-grass rangelands in the northern Great Basin. US Department of Agriculture, Forest Service, Intermountain Research Station Ogden, UT, USA.
- Burkhardt, J. W. and E. Tisdale. 1969. Nature and successional status of western juniper vegetation in Idaho. *Journal of Range Management*:264-270.
- Busse, D., A. Simon, and M. Riegel. 2000. Tree-growth and understory responses to low-severity prescribed burning in thinned *Pinus ponderosa* forests of central Oregon. *Forest Science* 46:258- 268.
- Busso, C. A. and J. H. Richards. 1995. Drought and clipping effects on tiller demography and growth of two tussock grasses in Utah. *Journal of Arid Environments* 29:239-251.
- Chambers, J. C., B. A. Roundy, R. R. Blank, S. E. Meyer, and A. Whittaker. 2007. What makes great basin sagebrush ecosystems invasible by *Bromus tectorum*? *Ecological Monographs* 77:117-145.
- Chambers, Jeanne C.; Pyke, David A.; Maestas, Jeremy D.; Pellant, Mike; Boyd, Chad S.; Campbell, Steven B.; Espinosa, Shawn; Havlina, Douglas W.; Mayer, Kenneth E.; Wuenschel, Amarina. 2014. Using resistance and resilience concepts to reduce impacts of invasive annual grasses and altered fire regimes on the sagebrush ecosystem and greater sage-grouse: A strategic multi-scale approach. Gen. Tech. Rep. RMRS-GTR-326. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 73 p.
- Clark, R. G., M. B. Carlton, and F. A. Sneva. 1982. Mortality of bitterbrush after burning and clipping in eastern Oregon. *Journal of Range Management* 35:711-714.
- Clements, C. D. and J. A. Young. 2002. Restoring antelope bitterbrush. *Rangelands* 24:3-6.



- Comstock, J. P. and J. R. Ehleringer. 1992. Plant adaptation in the Great Basin and Colorado plateau. *Western North American Naturalist* 52:195-215.
- Conrad, C. E. and C. E. Poulton. 1966. Effect of a wildfire on Idaho fescue and bluebunch wheatgrass. *Journal of Range Management*:138-141.
- Cook, J. G., T. J. Hershey, and L. L. Irwin. 1994. Vegetative response to burning on Wyoming mountain shrub big game ranges.. *Journal of Range Management* 47:296-302.
- Daubenmire, R. 1970. *Steppe vegetation of Washington*. 131 pp.
- Daubenmire, R. 1975. Plant succession on abandoned fields, and fire influences in a steppe area in southeastern Washington. *Northwest Science* 49:36-48.
- Dobrowolski, J. P., M. M. Caldwell, and J. H. Richards. 1990. Basin hydrology and plant root systems. In: C. B. Osmand, L. F. Pitelka, G. M. Hildy [eds]. *Plant biology of the Basin and range*. *Ecological Studies*. 80: 243-292
- Driscoll, R. S. 1964. A Relict Area in the Central Oregon Juniper Zone. *Ecology* 45:345-353.
- Eckert Jr, R. E. and J. S. Spencer. 1986. Vegetation response on allotments grazed under rest-rotation management. *Journal of Range Management*:166-174.
- Eckert, R. E., Jr. and J. S. Spencer. 1987. Growth and reproduction of grasses heavily grazed under restoration management. *Journal of Range Management* 40:156-159.
- Furniss, M. M. and W. F. Barr. 1975. Insects affecting important native shrubs of the northwestern United States General Technical Report INT-19. Intermountain Forest and Range Experiment Station, U.S. Department of Agriculture, Forest Service. Ogden, UT. p. 68
- Gaffney, W. S. 1941. The effects of winter elk browsing, South Fork of the Flathead River, Montana. *The Journal of Wildlife Management* 5:427-453.
- Garrison, G. A. 1953. Effects of Clipping on Some Range Shrubs. *Journal of Range Management* 6:309-317.
- Houston, D. B. 1973. Wildfires in northern Yellowstone National Park. *Ecology* 54:1111-1117.
- Johnson, C. G., R. R. Clausnitzer, P. J. Mehringer, and C. Oilver. 1994. Biotic and abiotic processes of eastside ecosystems: The effects of management on plant and community ecology, and on stand and landscape vegetation dynamics. Forest Service general technical report. Forest Service, Portland, OR (United States). Pacific Northwest Research Station.
- Kerns, B. K., W. G. Thies, and C. G. Niwa. 2006. Season and severity of prescribed burn in ponderosa pine forests: implications for understory native and exotic plants. *Ecoscience* 13:44-55.
- Laycock, W. A. 1967. How heavy grazing and protection affect sagebrush-grass ranges. *Journal of Range Management*:206-213.
- McArthur, E. D., A. Blaner, A. P. Plummer, and R. Stevens. 1982. Characteristics and hybridization of important Intermountain shrubs: 3. Sunflower family. U.S. Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station. Research Paper INT-177 43.
- McConnell, B. R. and J. G. Smith. 1977. Influence of grazing on age-yield interactions in bitterbrush. *Journal of Range Management* 30:91-93.
- Miller, R. F. R. J. T. 2000. The role of fire in juniper and pinyon woodlands: a descriptive analysis. Pages p. 15-30 in *Proceedings of the invasive species workshop: the role of fire in the control and spread of invasive species.*, Tallahassee, Florida.

- Miller, R. F. and E. K. Heyerdahl. 2008. Fine-scale variation of historical fire regimes in sagebrush-steppe and juniper woodland: an example from California, USA. *International Journal of Wildland Fire* 17:245-254.
- Mueggler, W. F. 1975. Rate and Pattern of Vigor Recovery in Idaho Fescue and Bluebunch Wheatgrass. *Journal of Range Management* 28:198-204.
- Murray, R. 1983. Response of antelope bitterbrush to burning and spraying in southeastern Idaho. Tiedemann, Arthur R.; Johnson, Kendall L., compilers. Research and management of bitterbrush and cliffrose in western North America. General Technical Report INT-152. Ogden, UT: US Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station:142-152.
- Neuenschwander, L. 1980. Broadcast burning of sagebrush in the winter. *Journal of Range Management*:233-236.
- Noy-Meir, I. 1973. Desert Ecosystems: Environment and Producers. *Annual Review of Ecology and Systematics*. Vol. 4:25-51.
- Richards, J. H. and M. M. Caldwell. 1987. Hydraulic lift: Substantial nocturnal water transport between soil layers by *Artemisia tridentata* roots. *Oecologia* 73:486-489.
- Robberecht, R. and G. Defossé. 1995. The relative sensitivity of two bunchgrass species to fire. *International Journal of Wildland Fire* 5:127-134.
- Stringham, T.K., P. Novak-Echenique, P. Blackburn, D. Snyder, and A. Wartgow. 2015. Final Report for USDA Ecological Site Description State-and-Transition Models by Disturbance Response Groups, Major Land Resource Area 25 Nevada. University of Nevada Reno, Nevada Agricultural Experiment Station Research Report 2015-02. p. 569
- Tisdale, E. W. and M. Hironaka. 1981. The sagebrush-grass region: A review of the ecological literature. University of Idaho, Forest, Wildlife and Range Experiment Station.
- Uresk, D. W., J. F. Cline, and W. H. Rickard. 1976. Impact of wildfire on three perennial grasses in southcentral Washington. *Journal of Range Management* 29:309-310.
- Wood, M. K., Bruce A. Buchanan, & William Skeet. 1995. Shrub preference and utilization by big game on New Mexico reclaimed mine land. *Journal of Range Management* 48:431-437.
- Wright, H. A. 1971. Why Squirreltail Is More Tolerant to Burning than Needle-and-Thread. *Journal of Range Management* 24:277-284.
- Wright, H.A., L.F. Neuenschwander, and C.M. Britton. 1979. The role and use of fire in sagebrush-grass and pinyon-juniper plant communities: A state-of-the-art review. Gen. Tech. Rep. INT-58. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station. 48 p.
- Young, R. P. 1983. Fire as a vegetation management tool in rangelands of the intermountain region. Pages 18-31 in *Managing intermountain rangelands - improvement of range and wildlife habitats*. USDA, Forest Service.
- Ziegenhagen, L. L. 2003. Shrub reestablishment following fire in the mountain big sagebrush (*Artemisia tridentata* Nutt. ssp. *vaseyana* (Rydb.) Beetle) alliance. M.S. Oregon State University.
- Ziegenhagen, L. L. and R. F. Miller. 2009. Postfire recovery of two shrubs in the interiors of large burns in the Intermountain West, USA. *Western North American Naturalist* 69:195-205.

## Contributors

A.V. Bahn, R.H. Barrett  
Bob Gillaspy  
E. Ersch

## Approval

Kendra Moseley, 4/25/2024

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Old SS Manuscripts, Range Site Descriptions, etc.
Contact for lead author	
Date	05/05/2024
Approved by	Kendra Moseley
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:**

---

2. **Presence of water flow patterns:**

---

3. **Number and height of erosional pedestals or terracettes:**

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

---

5. **Number of gullies and erosion associated with gullies:**

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

---

7. **Amount of litter movement (describe size and distance expected to travel):**

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
- 
14. **Average percent litter cover (%) and depth ( in):**
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
- 
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
- 
17. **Perennial plant reproductive capability:**
-