

Ecological site R026XF606CA Sandy Upland Pinyon (BLM)

Last updated: 4/10/2024
Accessed: 04/25/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 026X–Carson Basin and Mountains

The area lies within western Nevada and eastern California, with about 69 percent being within Nevada, and 31 percent being within California. Almost all this area is in the Great Basin Section of the Basin and Range Province of the Intermontane Plateaus. Isolated north-south trending mountain ranges are separated by aggraded desert plains. The mountains are uplifted fault blocks with steep side slopes. Most of the valleys are drained by three major rivers flowing east across this MLRA. A narrow strip along the western border of the area is in the Sierra Nevada Section of the Cascade-Sierra Mountains Province of the Pacific Mountain System. The Sierra Nevada Mountains are primarily a large fault block that has been uplifted with a dominant tilt to the west. This structure leaves an impressive wall of mountains directly west of this area. This helps create a rain shadow affect to MLRA 26. Parts of this eastern face, but mostly just the foothills, mark the western boundary of this area. Elevations range from about 3,806 feet (1,160 meters) on the west shore of Pyramid Lake to 11,653 feet (3,552 meters) on the summit of Mount Patterson in the Sweetwater Mountains.

Valley areas are dominantly composed of Quaternary alluvial deposits with Quaternary playa or alluvial flat deposits often occupying the lowest valley bottoms in the internally drained valleys, and river deposited alluvium being dominant in externally drained valleys. Hills and mountains are dominantly Tertiary andesitic flows, breccias, ash flow tuffs, rhyolite tuffs or granodioritic rocks. Quaternary basalt flows are present in lesser amounts, and Jurassic and Triassic limestone and shale, and Precambrian limestone and dolomite are also present in very limited amounts. Also of limited extent are glacial till deposits along the east flank of the Sierra Nevada Mountains, the result of alpine glaciation.

The average annual precipitation in this area is 5 to 36 inches (125 to 915 millimeters), increasing with elevation. Most of the rainfall occurs as high-intensity, convective storms in spring and autumn. Precipitation is mostly snow in winter. Summers are dry. The average annual temperature is 37 to 54 degrees F (3 to 12 degrees C). The freeze-free period averages 115 days and ranges from 40 to 195 days, decreasing in length with elevation.

The dominant soil orders in this MLRA are Aridisols and Mollisols. The soils in the area dominantly have a mesic soil temperature regime, an aridic or xeric soil moisture regime, and mixed or smectitic mineralogy. They generally are well drained, are clayey or loamy and commonly skeletal, and are very shallow to moderately deep.

This area supports shrub-grass vegetation characterized by big sagebrush. Low sagebrush and Lahontan sagebrush occur on some soils. Antelope bitterbrush, squirreltail, desert needlegrass, Thurber needlegrass, and Indian ricegrass are important associated plants. Green ephedra, Sandberg bluegrass, Anderson peachbrush, and several forb species also are common. Juniper-pinyon woodland is typical on mountain slopes. Jeffrey pine, lodgepole pine, white fir, and manzanita grow on the highest mountain slopes. Shadscale is the typical plant in the drier parts of the area. Sedges, rushes, and moisture-loving grasses grow on the wettest parts of the wet flood plains and terraces. Basin wildrye, alkali sacaton, saltgrass, buffaloberry, black greasewood, and rubber rabbitbrush grow on the drier sites that have a high concentration of salts.

Some of the major wildlife species in this area are mule deer, coyote, beaver, muskrat, jackrabbit, cottontail, raptors, pheasant, chukar, blue grouse, mountain quail, and mourning dove. The species of fish in the area include trout and catfish. The Lahontan cutthroat trout in the Truckee River is a threatened and endangered species.

LRU notes

The Mono-Adobe-Long Valleys LRU is comprised of the basins surrounding Mono Lake, Adobe Valley, and Long Valley to the southeast. Pleistocene and Holocene age alluvium and lacustrine deposits predominate. Ash layers occur from eruptions of the numerous volcanic domes that are mostly in adjacent LRUs. Soil temperature regimes are mesic and soil moisture regimes are aridic. Elevations range from 1310 to 2680 meters and slopes are typically less than 10 percent, however there are some ecological sites within the Mono-Adobe-Long Valleys LRU that are greater than 10 percent. Frost free days (FFD) range from 97-125.

Ecological site concept

The Sandy Upland Pinyon (BLM) site is found on sideslope hills at slopes between 15 and 30 percent. The soil is deep to a bedrock layer. The soil surface texture is very stony loamy sand. The dominant plants are singleleaf pinyon (*Pinus monophylla*) and Utah juniper (*Juniperus osteosperma*).

Associated sites

R026XF605CA	Sandy Juniper Flat (BLM) Soil is moderately deep to a restrictive layer.
R026XF003CA	Sandy 8-12" P.Z. Site is located on slopes less than 15 percent.
R026XF004CA	Gravelly Coarse Loamy 8-12" P.Z. Site is located on slopes less than 15 percent.

Table 1. Dominant plant species

Tree	(1) <i>Pinus monophylla</i> (2) <i>Juniperus osteosperma</i>
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

The Sandy Upland Pinyon (BLM) site is found on sideslope hills at slopes between 15 and 30 percent. The elevation ranges from 6800 to 7300 feet.

Table 2. Representative physiographic features

Landforms	(1) Hill
Runoff class	Very high
Elevation	6,800–7,300 ft
Slope	15–30%

Climatic features

The climate on this site is characterized by cold winters (20 to 40 degrees F) and warm, mostly dry summers (45 to 85 degrees F). The average annual precipitation ranges from 8 to 12 inches, with most falling as snow from November to March.

Table 3. Representative climatic features

Frost-free period (characteristic range)	
Freeze-free period (characteristic range)	
Precipitation total (characteristic range)	8-12 in
Frost-free period (average)	108 days
Freeze-free period (average)	
Precipitation total (average)	12 in




Figure 1. Monthly precipitation range




Figure 2. Monthly minimum temperature range




Figure 3. Monthly maximum temperature range




Figure 4. Monthly average minimum and maximum temperature




Figure 5. Annual precipitation pattern




Figure 6. Annual average temperature pattern

Climate stations used

- (1) BODIE CA ST HISTORIC PARK [USC00040943], Bridgeport, CA

Influencing water features

The Sandy Upland Pinyon (BLM) is not influenced by water features.

Soil features

The soil is well drained and deep to bedrock. The soil was formed in volcanic ash over residuum weathered from basalt. The soil surface texture is very stony loamy sand. Rocks on the surface are over 35 percent cover. Rocks in the subsurface are over 35 percent volume. The associated soil series is Pizona (CA802 mapunits 140 and 143).

Table 4. Representative soil features

Parent material	(1) Volcanic ash (2) Residuum–basalt
-----------------	-----------------------------------------

Surface texture	(1) Very stony loamy sand
Drainage class	Well drained
Permeability class	Moderately slow
Depth to restrictive layer	40–60 in
Surface fragment cover <=3"	20%
Surface fragment cover >3"	26%
Available water capacity (Depth not specified)	3–4.4 in
Calcium carbonate equivalent (Depth not specified)	0%
Electrical conductivity (Depth not specified)	0 mmhos/cm
Sodium adsorption ratio (Depth not specified)	0
Soil reaction (1:1 water) (Depth not specified)	6.6–7.3
Subsurface fragment volume <=3" (Depth not specified)	18–19%
Subsurface fragment volume >3" (Depth not specified)	28–29%

Ecological dynamics

Disturbance Response Group (DRG) 18 consists of four ecological sites; F026XY062NV, F026XY064NV, F026XY092NV, and F026XY093NV (Stringham et al. 2021). Site R026XF606CA is similar to sites found in Disturbance Response Group 18. The group falls in the 8 to 14 inch precipitation zone. Elevations range from 4,500 to 8,000 feet and these sites are found on slopes ranging from 2 to 75 percent. The soils in this group are typically shallow to very shallow and available water holding capacity is low. These soils usually have high amounts of rock fragments at the soil surface which help to reduce evaporation and provide a stabilizing effect on erosion conditions. This group is dominated by singleleaf pinyon (*Pinus monophylla*) and/or Utah juniper (*Juniperus osteosperma*) with Wyoming big sagebrush (*Artemisia tridentata* ssp. *wyomingensis*) or low sagebrush (*Artemisia arbuscula*) as the primary understory shrub. Other shrubs in the group include antelope bitterbrush (*Purshia tridentata*), Douglas rabbitbrush (*Chrysothamnus viscidiflorus*), green ephedra (*Ephedra viridis*), and currant (*Ribes* spp.). The dominant understory grass of the group is Thurber's needlegrass (*Achnatherum thurberianum*). Other understory grasses include muttongrass (*Poa fendleriana*), Sandberg bluegrass (*Poa secunda*), and Indian ricegrass (*Achnatherum hymenoides*). Under medium canopy cover (11-30 percent, dependent on ecological site), understory production ranges from 75 to 400 lbs/ac.

General State and Transition Model Narrative for Group 18:

This is a text description of the states, phases, transitions, and community pathways possible in the State and Transition model for the MLRA 26 disturbance response group 18.

Reference State 1.0:

The Reference State 1.0 is representative of the natural range of variability under pristine conditions. This reference state has four general community phases: an old-growth tree phase, a shrub-herbaceous phase, an immature tree phase, and an infilled tree phase. State dynamics are maintained by interactions between climatic patterns and disturbance regimes. Negative feedbacks enhance ecosystem resilience and contribute to the stability of the state. These include the presence of all structural and functional groups, low fine fuel loads, and retention of organic matter and nutrients. Plant community phase changes are primarily driven by fire, periodic drought, and/or insect or disease attack. Fires within this community are infrequent and likely small and patchy due to low fuel loads. This fire type will create a plant community mosaic that will include all/most of the following community phases within this state.

Community Phase 1.1:

This phase is characterized by widely dispersed old-growth pinyon and juniper trees with a Wyoming big sagebrush, perennial bunchgrass understory. The visual aspect is dominated by singleleaf pinyon and Utah juniper with over 15 percent canopy cover (USDA 1997). Trees have reached maximal or near maximal heights for the site and many tree crowns may be flat- or round-topped. Thurber's needlegrass is most prevalent grass in the understory. Wyoming big sagebrush is the primary understory shrub. Forbs such as phlox, and eriogonum are minor components. Overall, the understory is sparse with production ranging between 200 to 400 pounds per acre.

Community Phase Pathway 1.1a, from Phase 1.1 to 1.2:

A high-severity crown fire will eliminate or reduce the singleleaf pinyon and Utah juniper overstory and the shrub component. This allows for the perennial bunchgrasses to dominate the site.

Community Phase Pathway 1.1b, from Phase 1.1 to 1.4:

Time without disturbances such as fire, drought, or disease will allow for the gradual infilling of singleleaf pinyon and Utah juniper.

Community Phase 1.2:

This community phase is characterized by a post-fire shrub and herbaceous community. Thurber's needlegrass and other perennial grasses dominate. Forbs may increase after a fire but will likely return to pre-burn levels within a few years. Pinyon and juniper seedlings up to 4 feet in height may be present. Wyoming big sagebrush may be present in unburned patches. Burned tree skeletons may be present; however, these have little or no effect on the understory vegetation.

Community Phase Pathway 1.2a, from Phase 1.2 to 1.3:

Time without disturbances such as fire, drought, or disease will allow for the gradual maturation of the singleleaf pinyon and Utah Juniper component. Wyoming big sagebrush reestablishes. Excessive herbivory may also reduce perennial grass understory.

Community Phase 1.3:

This community phase is characterized by an immature woodland, with pinyon and juniper trees averaging over 4.5 feet in height. Tree canopy cover is between 10 to 20 percent. Tree crowns are typically cone- or pyramidal-shaped. Understory vegetation is dominated by Wyoming big sagebrush and perennial bunchgrasses as well as smaller tree seedling and saplings.

Community Phase Pathway 1.3a, from 1.3 to 1.4:

Time without disturbances such as fire, drought, or disease will allow for the gradual maturation of singleleaf pinyon and Utah juniper. Infilling by younger trees continues. Excessive herbivory may also reduce the perennial grass understory.

Community Phase Pathway 1.3b, from Phase 1.3 to 1.2:

Fire reduces or eliminates tree canopy, allowing perennial grasses to dominate the site.

Community Phase 1.4 (at-risk):

This phase is dominated by singleleaf pinyon and Utah juniper. The stand exhibits mixed age classes and canopy cover may be 30percent or greater. The density and vigor of the Wyoming big sagebrush and perennial bunchgrass understory is decreased. Bare ground areas are likely to increase. Mat-forming forbs such as phlox may increase. This community is at risk of crossing a threshold; without proper management this phase will transition to the infilled tree state 3.0. This community phase is typically described as early Phase II woodland (Miller et al. 2008).

Community Phase Pathway 1.4a, from Phase 1.4 to 1.1:

Low intensity fire, insect infestation, or disease kills individual trees within the stand reducing canopy cover to less than 30percent. Over time young trees mature to replace and maintain the old-growth woodland. The Wyoming big sagebrush and perennial bunchgrass community increases in density and vigor.

Community Phase Pathway 1.4b, from Phase 1.4 to 1.2:

A high-severity crown fire will eliminate or reduce the singleleaf pinyon and Utah juniper overstory and the shrub component which will allow for the perennial bunchgrasses to dominate the site.

T1A: Transition from Reference State 1.0 to Current Potential State 2.0:

Trigger: Introduction of non-native annual species.

Slow variables: Over time the annual non-native plants will increase within the community.

Threshold: Any amount of introduced non-native species causes an immediate decrease in the resilience of the site.

Annual non-native species cannot be easily removed from the system and have the potential to significantly alter disturbance regimes from their historic range of variation.

T1B: Transition from Reference State 1.0 to Infilled Tree State 3.0

Trigger: Time and a lack of disturbance allow trees to dominate site resources; may be coupled with inappropriate herbivory that favors shrub and tree dominance.

Slow variables: Over time the abundance and size of trees will increase.

Threshold: Pinyon and juniper canopy cover is greater than 40percent. Little understory vegetation remains due to competition with trees for site resources.

Current Potential State 2.0:

This state is similar to the Reference State 1.0, with four general community phases: an old-growth tree phase, a shrub-herbaceous phase, an immature tree phase, and an infilled tree phase. Ecological function has not changed, however the resiliency of the state has been reduced by the presence of non-native species. These non-natives, particularly cheatgrass, can be highly flammable and promote fire where historically fire had been infrequent. Negative feedbacks enhance ecosystem resilience and contribute to the stability of the state. These include the presence of all structural and functional groups, low fine fuel loads and retention of organic matter and nutrients. Positive feedbacks decrease ecosystem resilience and stability of the state. These include the non-natives' high seed output, persistent seed bank, rapid growth rate, ability to cross pollinate, and adaptations for seed dispersal. Fires within this community with the small amount of non-native annual species present are likely still small and patchy due to low fuel loads. This fire type will create a plant community mosaic that will include all/most of the following community phases within this state.

Community Phase 2.1:

This phase is characterized by a widely dispersed old-growth pinyon and juniper trees with a Wyoming big sagebrush perennial bunchgrass understory. The visual aspect is dominated by singleleaf pinyon and Utah juniper with canopy cover of 15 percent or more (USDA 1997). Trees have reached maximal or near maximal heights for the site and many tree crowns may be flat- or round-topped. Thurber's needlegrass is the most prevalent grass in the understory. Wyoming big sagebrush is the primary understory shrub. Forbs such as phlox and eriogonum are minor components. Overall, the understory is sparse with production ranging between 200 to 400 lbs. per acre.

Community Phase Pathway 2.1a, from Phase 2.1 to 2.2:

A high-severity crown fire will eliminate or reduce the singleleaf pinyon and Utah juniper overstory and the shrub component. This allows for the perennial bunchgrasses to dominate the site.

Community Phase Pathway 2.1b, from Phase 2.1 to 2.4:

Time without disturbances such as fire, drought, or disease will allow for the gradual infilling of singleleaf pinyon and Utah juniper.

Community Phase 2.2:

This community phase is characterized by a post-fire shrub and herbaceous community. Thurber's needlegrass and other perennial grasses dominate. Forbs may increase post-fire but will likely return to pre-burn levels within a few years. Pinyon and juniper seedlings up to 4 feet in height may be present. Wyoming big sagebrush may be present in unburned patches. Burned tree skeletons may be present; however, these have little or no effect on the understory vegetation. Annual non-native species generally respond well after fire and may be stable or increasing within the community.

Community Phase Pathway 2.2a, from Phase 2.2 to 2.3:

Time without disturbances such as fire, drought, or disease will allow for the gradual maturation of the singleleaf pinyon and Utah Juniper component. Wyoming big sagebrush reestablishes. Excessive herbivory may also reduce perennial grass understory.

Community Phase 2.3:

This community phase is characterized by an immature woodland, with pinyon and juniper trees averaging over 4.5

feet in height. Tree canopy cover is between 10 to 20 percent. Tree crowns are typically cone- or pyramidal-shaped. Understory vegetation is dominated by Wyoming big sagebrush and perennial bunchgrasses as well as smaller tree seedling and saplings. Annual non-native species are present.

Community Phase Pathway 2.3a, from Phase 2.3 to 2.4:

Time without disturbances such as fire, drought, or disease will allow for the gradual maturation of singleleaf pinyon and Utah juniper. Infilling by younger trees continues.

Community Phase Pathway 2.3b, from Phase 2.3 to 2.2:

Fire reduces or eliminates tree canopy, allowing perennial grasses to dominate the site.

Community Phase 2.4 (at-risk):

This phase is dominated by singleleaf pinyon and Utah juniper. The stand exhibits mixed age classes and canopy cover exceeds 30 percent. The density and vigor of the Wyoming big sagebrush and perennial bunchgrass understory is decreased. Bare ground areas are likely to increase. Mat-forming forbs may increase. Annual non-native species are present primarily under tree canopies. This community is at risk of crossing a threshold, without proper management this phase will transition to the infilled tree state 3.0. This community phase is typically described as early Phase II woodland (Miller et al. 2008).

Community Phase Pathway 2.4a, from Phase 2.4 to 2.1:

Low intensity fire, insect infestation, or disease kills individual trees within the stand reducing canopy cover to less than 30 percent. Over time young trees mature to replace and maintain the old-growth woodland. The Wyoming big sagebrush and perennial bunchgrass community increases in density and vigor. Annual non-natives present in trace amounts.

Community Phase Pathway 2.4b, from Phase 2.4 to 2.2:

A high-severity crown fire will eliminate or reduce the singleleaf pinyon and Utah juniper overstory and the shrub component which will allow for the perennial bunchgrasses to dominate the site. Annual non-native grasses typically respond positively to fire and may increase in the post-fire community.

T2A: Transition from Current Potential State 2.0 to Infilled Tree State 3.0:

Trigger: Time and a lack of disturbance allow trees to dominate site resources; may be coupled with inappropriate grazing management that favors shrub and tree dominance.

Slow variables: Over time the abundance and size of trees will increase.

Threshold: Singleleaf pinyon and Utah juniper canopy cover is greater than 40 percent. Little understory vegetation remains due to competition with trees for site resources.

T2B: Transition from Current Potential State 2.0 to Annual State 4.0:

Trigger: Catastrophic crown fire facilitates the establishment of non-native, annual weeds.

Slow variables: Increase in tree crown cover, loss of perennial understory and an increase in annual non-native species.

Threshold: Cheatgrass or other non-native annuals dominate understory. Loss of deep-rooted perennial bunchgrasses changes spatial and temporal nutrient cycling and nutrient redistribution, and reduces soil organic matter. Increased canopy cover of trees allows severe stand-replacing fire. The increased seed bank of non-native, annual species responds positively to post-fire conditions facilitating the transition to an Annual State.

Infilled Tree State 3.0:

This state has two community phases that are characterized by the dominance of Utah juniper and singleleaf pinyon in the overstory. This state is identifiable by over 40 percent cover of Utah juniper and singleleaf pinyon, exhibiting a mixed age class. Older trees are at maximal height and upper crowns may be flat-topped or rounded. Younger trees are typically cone- or pyramidal-shaped. Understory vegetation is sparse due to increasing shade and competition from trees.

Community Phase 3.1:

Singleleaf pinyon and Utah juniper dominate the aspect. Understory vegetation is thinning. Perennial bunchgrasses are sparse and Wyoming big sagebrush skeletons are as common as live shrubs due to tree competition for soil water, overstory shading, and duff accumulation. Tree canopy cover is greater than 40 percent. Annual non-native

species are present or co-dominate in the understory. Bare ground areas are prevalent. This community phase is typically described as a Phase II woodland (Miller et al. 2008).

Community Phase Pathway 3.1a, from Phase 3.1 to 3.2:

Time without disturbances such as fire, drought, or disease will allow for the gradual maturation of singleleaf pinyon and Utah juniper. Infilling by younger trees continues.

Community Phase 3.2 (at risk):

Singleleaf pinyon and Utah juniper dominate the aspect. Tree canopy cover exceeds 40 percent. Understory vegetation is sparse to absent. Perennial bunchgrasses, if present exist in the drip line or under the canopy of trees. Wyoming big sagebrush skeletons are common or the sagebrush has been extinct long enough that only scattered limbs remain. Mat-forming forbs or Sandberg bluegrass (*Poa secunda*) may dominate interspaces. Annual non-native species are present and are typically found under the trees. Bare ground areas are large and interconnected. Soil redistribution may be extensive. This community phase is typically described as a Phase III woodland (Miller et al. 2008).

T3A: Transition from Infilled Tree State 3.0 to Annual State 4.0:

Trigger: Canopy fire reduces the pinyon and juniper overstory and facilitates the annual non-native species in the understory to dominate the site.

Slow variables: Over time, cover, production and seed bank of annual non-native species increases.

Threshold: Loss of deep-rooted perennial bunchgrasses and shrubs changes temporal and spatial nutrient capture and cycling within the community. Increase in canopy cover of trees increases rainfall interception and reduces soil moisture for understory species. Increased canopy cover of trees increases the risk for severe stand-replacing crown fire. The increased seed bank of non-native, annual species responds positively to post-fire conditions facilitating the transition to an Annual State.

R3A: Restoration from Infilled Tree state 3.0 to Current Potential State 2.0:

Manual or mechanical thinning of trees coupled with seeding. Probability of success is highest from community phase 3.1.


Annual State 4.0:

This state has one community phase that is characterized by the dominance of annual non-native species such as cheatgrass and tansy mustard in the understory. Time since fire may facilitate the maturation of sprouting shrubs such as rabbitbrush. Ecological dynamics are significantly altered in this state. Annual non-native species create a highly combustible fuel bed that shortens the fire return interval. Nutrient cycling is spatially and temporally truncated as annual plants contribute significantly less to deep soil carbon. This state was not seen in MLRA 26 during field work for this project, however it is possible given increased fire activity in these sites and their proximity to known annual states of sagebrush ecological sites. We refer the reader to the report for Disturbance Response Group 21 for MLRA 28A and 28B.

Community Phase 4.1:

Cheatgrass, mustards and other non-native annual species dominate the site. Trace amounts of perennial bunchgrasses may be present. Sprouting shrubs may increase. Burned tree skeletons present.

State and transition model



Reference State 1.0 Community Pathways

1.1a: High severity crown fire reduces or eliminates tree cover.

1.1b: Time and lack of disturbance such as fire, disease, or drought allows younger trees to infill.

1.2a: Time and lack of disturbance such as fire or drought. Excessive herbivory may also reduce perennial grass understory.

1.3a: Time and lack of disturbance such as fire or drought. Excessive herbivory may also reduce perennial grass understory.

1.3b: Fire.

1.4a: Low severity fire, insect infestation, or disease removes individual trees and reduces total tree cover.

1.4b: High severity crown fire reduces or eliminates tree cover.

Transition T1A: Introduction of non-native annual species.

Transition T1B: Time and a lack of disturbance allows for trees to dominate site resources; may be coupled with inappropriate grazing management that favors shrub and tree dominance.

Current Potential State 1.0 Community Pathways

2.1a: High severity crown fire reduces or eliminates tree cover.

2.1b: Time and lack of disturbance such as fire, disease, or drought allows younger trees to infill.

2.2a: Time and lack of disturbance such as fire or drought. Excessive herbivory may also reduce perennial grass understory.

2.3a: Time and lack of disturbance such as fire or drought. Excessive herbivory may also reduce perennial grass understory.

2.3b: Fire.

2.4a: Low severity fire, insect infestation, or disease removes individual trees and reduces total tree cover.

2.4b: High severity crown fire reduces or eliminates tree cover.

Transition T2A: Time and a lack of disturbance allows for trees to dominate site resources; may be coupled with inappropriate grazing management that favors shrub and tree dominance.

Transition T2B: Catastrophic fire.

Infilled Tree State 3.0 Community Pathways

3.1a: Time and lack of disturbance such as fire, disease, or drought allows younger trees to infill.

Transition T3A: Catastrophic fire.

Restoration Pathway R3A: Thinning of trees coupled with seeding. Success unlikely from phase 3.2.

Annual State 4.0 Community Pathways

None.

State 1

Reference State

Dominant plant species

- singleleaf pinyon (*Pinus monophylla*), tree
- Utah juniper (*Juniperus osteosperma*), tree

Inventory data references

NASIS data for soil survey CA802 mapunits 140 and 143.

References

Stringham, T.K., D. Snyder, P. Novak-Echenique, K. O'Neill, A. Lyons, and M. Johns. 2021. Great Basin Ecological Site Development Project: State-and-Transition Models for Major Land Resource Area 26, Nevada and Portions of California..

Contributors

Patti Novak-Echenique

Tamzen Stringham

Approval

Kendra Moseley, 4/10/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	04/25/2024
Approved by	Kendra Moseley
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial**

distribution on infiltration and runoff:

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
-

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
-