

Ecological site R026XY019NV CHURNING CLAY 10-12 P.Z.

Last updated: 4/10/2024
Accessed: 05/18/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 026X–Carson Basin and Mountains

The area lies within western Nevada and eastern California, with about 69 percent being within Nevada, and 31 percent being within California. Almost all this area is in the Great Basin Section of the Basin and Range Province of the Intermontane Plateaus. Isolated north-south trending mountain ranges are separated by aggraded desert plains. The mountains are uplifted fault blocks with steep side slopes. Most of the valleys are drained by three major rivers flowing east across this MLRA. A narrow strip along the western border of the area is in the Sierra Nevada Section of the Cascade-Sierra Mountains Province of the Pacific Mountain System. The Sierra Nevada Mountains are primarily a large fault block that has been uplifted with a dominant tilt to the west. This structure leaves an impressive wall of mountains directly west of this area. This helps create a rain shadow affect to MLRA 26. Parts of this eastern face, but mostly just the foothills, mark the western boundary of this area. Elevations range from about 3,806 feet (1,160 meters) on the west shore of Pyramid Lake to 11,653 feet (3,552 meters) on the summit of Mount Patterson in the Sweetwater Mountains.

Valley areas are dominantly composed of Quaternary alluvial deposits with Quaternary playa or alluvial flat deposits often occupying the lowest valley bottoms in the internally drained valleys, and river deposited alluvium being dominant in externally drained valleys. Hills and mountains are dominantly Tertiary andesitic flows, breccias, ash flow tuffs, rhyolite tuffs or granodioritic rocks. Quaternary basalt flows are present in lesser amounts, and Jurassic and Triassic limestone and shale, and Precambrian limestone and dolomite are also present in very limited amounts. Also of limited extent are glacial till deposits along the east flank of the Sierra Nevada Mountains, the result of alpine glaciation.

The average annual precipitation in this area is 5 to 36 inches (125 to 915 millimeters), increasing with elevation. Most of the rainfall occurs as high-intensity, convective storms in spring and autumn. Precipitation is mostly snow in winter. Summers are dry. The average annual temperature is 37 to 54 degrees F (3 to 12 degrees C). The freeze-free period averages 115 days and ranges from 40 to 195 days, decreasing in length with elevation.

The dominant soil orders in this MLRA are Aridisols and Mollisols. The soils in the area dominantly have a mesic soil temperature regime, an aridic or xeric soil moisture regime, and mixed or smectitic mineralogy. They generally are well drained, are clayey or loamy and commonly skeletal, and are very shallow to moderately deep.

This area supports shrub-grass vegetation characterized by big sagebrush. Low sagebrush and Lahontan sagebrush occur on some soils. Antelope bitterbrush, squirreltail, desert needlegrass, Thurber needlegrass, and Indian ricegrass are important associated plants. Green ephedra, Sandberg bluegrass, Anderson peachbrush, and several forb species also are common. Juniper-pinyon woodland is typical on mountain slopes. Jeffrey pine, lodgepole pine, white fir, and manzanita grow on the highest mountain slopes. Shadscale is the typical plant in the drier parts of the area. Sedges, rushes, and moisture-loving grasses grow on the wettest parts of the wet flood plains and terraces. Basin wildrye, alkali sacaton, saltgrass, buffaloberry, black greasewood, and rubber rabbitbrush grow on the drier sites that have a high concentration of salts.

Some of the major wildlife species in this area are mule deer, coyote, beaver, muskrat, jackrabbit, cottontail, raptors, pheasant, chukar, blue grouse, mountain quail, and mourning dove. The species of fish in the area include trout and catfish. The Lahontan cutthroat trout in the Truckee River is a threatened and endangered species.

LRU notes

The "XY" LRU contains all the sites that cross other LRU boundaries. The sites are typically found along waterways.

Ecological site concept

The CHURNING CLAY 10-12 P.Z. occurs on nearly level to steeply sloping piedmont slopes, hills, mountains, lake terraces, and fan remnants. As the soil dries, deep wide cracks develop in these heavy textured soils which may extend to a depth of 40 inches. These openings in the soil result in rapid loss of soil moisture by exposing the subsoil to the atmosphere. The vertical and horizontal movements of the soil from alternative wetting and drying, shears fine and very fine roots and the establishment of plants with extensive lateral root systems is restricted. The combination of rapid moisture loss and root shear adversely affects root development. Disturbance of the soil surface by cracking and a sloughing of the surface layer into the cracks, favors plants capable of rapid regeneration through production of highly viable seed or rhizomatous growth habit.

Associated sites

| | |
|-------------|---------------------------|
| R026XY010NV | LOAMY 10-12 P.Z. |
| R026XY023NV | CLAYPAN 10-12 P.Z. |

Similar sites

| | |
|-------------|--|
| R026XY027NV | CHURNING CLAY 8-10 P.Z. ARAR8-ARTRW8 codominant shrubs; less productive site |
|-------------|--|

Table 1. Dominant plant species

| | |
|------------|---|
| Tree | Not specified |
| Shrub | (1) <i>Artemisia tridentata ssp. wyomingensis</i> (2) <i>Tetradymia glabrata</i> |
| Herbaceous | (1) <i>Pascopyrum smithii</i> (2) <i>Elymus elymoides</i> |

Physiographic features

The Churning Clay 10-12 P.Z. site occurs on nearly level to steeply sloping piedmont slopes, hills, mountains, lake terraces, and fan remnants. Slopes range from 0 to 50 percent. Elevations range from 4700 to 6200 feet.

Table 2. Representative physiographic features

| | |
|-----------|---|
| Landforms | (1) Fan piedmont (2) Hill (3) Fan remnant |
| Elevation | 1,433–1,890 m |
| Slope | 0–50% |
| Aspect | Aspect is not a significant factor |

Climatic features

The climate associated with this site is semiarid, characterized by cool, moist winters and warm, dry summers. Average annual precipitation is 10 to 12 inches. Mean annual air temperature is 49 to 52 degrees F. The average growing season is about 90 to 100 days.

Nevada's climate is predominantly arid, with large daily ranges of temperature, infrequent severe storms, heavy snowfall in the higher mountains, and great location variations with elevation. Three basic geographical factors largely influence Nevada's climate: continentality, latitude, and elevation. Continentality is the most important factor. The strong continental effect is expressed in the form of both dryness and large temperature variations. Nevada lies on the eastern, lee side of the Sierra Nevada Range, a massive mountain barrier that markedly influences the climate of the State. The prevailing winds are from the west, and as the warm moist air from the Pacific Ocean ascend the western slopes of the Sierra Range, the air cools, condensation occurs and most of the moisture falls as precipitation. As the air descends the eastern slope, it is warmed by compression, and very little precipitation occurs. The effects of this mountain barrier are felt not only in the West but throughout the state, with the result that the lowlands of Nevada are largely desert or steppes. The temperature regime is also affected by the blocking of the inland-moving maritime air. Nevada sheltered from maritime winds, has a continental climate with well-developed seasons and the terrain responds quickly to changes in solar heating.

Nevada lies within the mid-latitude belt of prevailing westerly winds which occur most of the year. These winds bring frequent changes in weather during the late fall, winter and spring months, when most of the precipitation occurs. To the south of the mid-latitude westerlies, lies a zone of high pressure in subtropical latitudes, with a center over the Pacific Ocean. In the summer, this high-pressure belt shifts northward over the latitudes of Nevada, blocking storms from the ocean. The resulting weather is mostly clear and dry during the summer and early fall, with scattered thundershowers. The eastern portion of the state receives significant summer thunderstorms generated from monsoonal moisture pushed up from the Gulf of California, known as the North American monsoon. The monsoon system peaks in August and by October the monsoon high over the Western U.S. begins to weaken and the precipitation retreats southward towards the tropics (NOAA 2004).

Table 3. Representative climatic features

| | |
|--|------------|
| Frost-free period (characteristic range) | |
| Freeze-free period (characteristic range) | |
| Precipitation total (characteristic range) | 254-305 mm |
| Frost-free period (average) | 95 days |
| Freeze-free period (average) | |
| Precipitation total (average) | 279 mm |




Figure 1. Monthly average minimum and maximum temperature




Figure 2. Annual precipitation pattern

Influencing water features

There are no influencing water features associated with this site.

Soil features

The soils are moderately deep and well drained formed in mixed pedisediment and residuum from tuff. The available water capacity is moderate. On partial drying, deep wide cracks develop in these heavy textured soils which may extend to a depth of 40 inches. These openings in the soil result in rapid loss of soil moisture by exposing the subsoil to the atmosphere. The vertical and horizontal movements of the soil from alternative wetting and drying, shears fine and very fine roots and the establishment of plants with extensive lateral root systems is restricted. The combination of rapid moisture loss and root shear adversely affects root development. Disturbance of the soil surface by cracking and a sloughing of the surface layer into the cracks, favors plants capable of rapid regeneration through production of highly viable seed or rhizomatous growth habit. The soil series associated with this site include: Waspo.

Table 4. Representative soil features

| | |
|---|---|
| Parent material | (1) Pedisediment–tuff (2) Residuum–tuff |
| Surface texture | (1) Clay (2) Gravelly clay (3) Stony clay |
| Family particle size | (1) Clayey |
| Drainage class | Well drained |
| Permeability class | Very slow |
| Soil depth | 51–102 cm |
| Surface fragment cover ≤3" | 7–16% |
| Surface fragment cover >3" | 2–12% |
| Available water capacity (0-101.6cm) | 9.14–10.92 cm |
| Calcium carbonate equivalent (0-101.6cm) | 0% |
| Electrical conductivity (0-101.6cm) | 0 mmhos/cm |
| Sodium adsorption ratio (0-101.6cm) | 0 |
| Soil reaction (1:1 water) (0-101.6cm) | 6.1–7.8 |

| | |
|--|------|
| Subsurface fragment volume <=3" (Depth not specified) | 8–9% |
| Subsurface fragment volume >3" (Depth not specified) | 2–4% |

Ecological dynamics

Where management results in over use by livestock, Wyoming big sagebrush, rabbitbrush and littleleaf horsebrush become more dominant. Species most likely to invade this site are annual grasses and forbs.

Fire Ecology:

The fire return interval for Wyoming big sagebrush communities ranges from 10 to 70 years. Fire is the principal means of renewal for decadent stands of Wyoming big sagebrush. Wyoming big sagebrush is killed by fire and establishes after fire from a seedbank; from seed produced by remnant plants that escaped fire; and from plants adjacent to the burn that seed in. Fire top-kills littleleaf horsebrush. When top-killed by fire, littleleaf horsebrush establishes by sprouting from the root crown. Rubber rabbitbrush is often top-killed by fire. Rubber rabbitbrush is a fire-adapted species that is typically unharmed or enhanced by fire. Recovery time is often rapid to very rapid. Rubber rabbitbrush is often one of the first species to colonize burned areas by sprouting or from off-site seed. The major adaptation of western wheatgrass to fire is its rhizomatous growth form. During a fire the coarse culms usually burn fast with little or no heat transferred to the roots. Recovery takes about 2 to 5 years after a fire. Bottlebrush squirreltail's small size, coarse stems, and sparse leafy material aid in its tolerance of fire. Postfire regeneration occurs from surviving root crowns and from on- and off-site seed sources. Frequency of disturbance greatly influences postfire response of bottlebrush squirreltail. Undisturbed plants within a 6 to 9 year age class generally contain large amounts of dead material, increasing bottlebrush squirreltail's susceptibility to fire. Sandberg bluegrass is generally unharmed by fire. It produces little litter, and its small bunch size and sparse litter reduces the amount of heat transferred to perennating buds in the soil. Its rapid maturation in the spring also reduces fire damage, since it is dormant when most fires occur.

Description of MLRA 26 Disturbance Response Group 3

Disturbance Response Group (DRG) 3, consists of three ecological sites (Stringham et al. 2021). The precipitation for these sites ranges from 8 to 12 inches. The elevation range for this group is 4,500 to 6,500 feet. Slopes range from 2 to 30 percent, however, 2 to 8 percent is typical. The soils in this group are deep and well drained. The water holding capacity ranges from low to moderate. The soils are heavy textured throughout and are subject to extreme shrink and swell action as they fluctuate from wet to dry. The vertical and horizontal soil movement from alternate wetting and drying shears fine and very fine roots and the establishment of plants with extensive lateral root systems is restricted. Annual production in a normal year ranges from 300 to 600 lbs/ac for the group. The potential native plant community varies depending on precipitation, elevation and landform. The shrub component is dominated by low sagebrush (*Artemisia arbuscula*), Lahontan sagebrush (*Artemisia arbuscula* ssp. *longicaulis*), or Wyoming big sagebrush (*Artemisia tridentata* ssp. *wyomingensis*). Other shrubs include spiny hopsage (*Grayia spinosa*), shadscale (*Atriplex confertifolia*), and littleleaf horsebrush (*Tetradymia glabrata*). The understory is dominated by perennial bunchgrasses, primarily bottlebrush squirreltail (*Elymus elymoides*), Sandberg bluegrass (*Poa secunda*) and western wheatgrass (*Pascopyrum smithii*).

Ecological sites in this group have multiple sagebrush species listed as the dominant shrub on the sites. During our visits to these sites, we used the black light test to verify sagebrush species. We identified Lahontan sagebrush as the dominant shrub during one site visit, and Wyoming big sagebrush as the dominant on others. On one occasion they occurred together on the modal site. The concepts for the three sites in this group may need reevaluation. Lahontan sagebrush was only recently identified as a unique subspecies of low sagebrush, so it may not have been apparent at the time some of these ecological sites were established. Additionally, these sites are mapped in highly disturbed areas, so site concepts may have been created on alternate states. Due to the differences in palatability between low sagebrush and Lahontan sagebrush, as well as potential soil differences, we recommend a reevaluation of the low sagebrush ecological sites in MLRA 26.

Disturbance Response Group 3 Ecological Sites:

Churning Clay 8-10" — Modal Site R026XY027NV
 Churning Clay 10-12" R026XY019NV

Potential Resilience Differences with Other Ecological Sites:

Churning Clay 10-12" (R026XY019NV):

This site occurs on similar landforms and at similar elevations as the modal site, but has a much different plant community that is dominated by western wheatgrass and Wyoming big sagebrush. This site is also much more productive than the modal site with 600 lb/ac in a normal year. The soils are deep, well drained and have moderate available water capacity. Root development is restricted due to shearing and soil moisture loss from shrink and swell action of the soil. Therefore, plants that rapidly regenerate through seed or have rhizomatous growth habits are favored.

State and Transition Model Narrative Group 3

This is a text description of the states, phases, transitions, and community pathways possible in the State and Transition model for the MLRA 26 Disturbance Response Group 3.

Reference State 1.0:

The Reference State 1.0 is representative of the natural range of variability under pristine conditions. The reference state has three general community phases: a shrub-grass dominant phase, a perennial grass dominant phase and a shrub dominant phase. State dynamics are maintained by interactions between climatic patterns and disturbance regimes. Negative feedbacks enhance ecosystem resilience and contribute to the stability of the state. These include the presence of all structural and functional groups, low fine fuel loads, and retention of organic matter and nutrients. Plant community phase changes are primarily driven by fire, periodic drought and/or insect or disease attack.

Community Phase 1.1:

This community is dominated by low sagebrush, bottlebrush squirreltail and Sandberg bluegrass. Forbs and other grasses make up smaller components.

Community Phase Pathway 1.1a, from Phase 1.1 to 1.2:

Fire will decrease or eliminate the overstory of sagebrush and allow for the perennial bunchgrasses to dominate the site. Fires will typically be low severity resulting in a mosaic pattern due to low fuel loads. A fire following an unusually wet spring may be more severe and reduce sagebrush cover to trace amounts.

Community Phase Pathway 1.1b, from Phase 1.1 to 1.3:

Time and lack of disturbance such as fire allows for sagebrush to increase and become decadent. Long-term drought, herbivory, or combinations of these will cause a decline in perennial bunchgrasses and fine fuels leading to a reduced fire frequency and allowing sagebrush to dominate the site.

Community Phase 1.2:

This community phase is characteristic of a post-disturbance, early/mid-seral community. Bottlebrush squirreltail, Sandberg bluegrass and other perennial bunchgrasses dominate. Depending on fire severity patches of intact sagebrush may remain. Rabbitbrush and other sprouting shrubs may be sprouting. Perennial forbs may be a significant component for a number of years following fire.

Community Phase Pathway 1.2a, from Phase 1.2 to 1.1:

Time and lack of disturbance will allow sagebrush to increase.

Community Phase 1.3:

Sagebrush increases in the absence of disturbance. Decadent sagebrush dominates the overstory and the deep-rooted perennial bunchgrasses in the understory are reduced either from competition with shrubs and/or from herbivory.

Community Phase Pathway 1.3a, from Phase 1.3 to 1.2:

Fire will decrease or eliminate the overstory of sagebrush and allow for the perennial bunchgrasses to dominate the site. Fires may be high severity in this community phase due to the dominance of sagebrush resulting in removal of overstory shrub community.

T1A: Transition from the Reference State 1.0 to Current Potential State 2.0:

Trigger: This transition is caused by the introduction of non-native annual plants, such as cheatgrass, mustards, and bur buttercup (*Ceratocephala testiculata*).

Slow variables: Over time the annual non-native species will increase within the community.

Threshold: Any amount of introduced non-native species causes an immediate decrease in the resilience of the site.

Annual non-native species cannot be easily removed from the system and have the potential to significantly alter disturbance regimes from their historic range of variation.

Current Potential State 2.0:

This state is similar to the Reference State 1.0. Ecological function has not changed, however, the resiliency of the state has been reduced by the presence of invasive weeds. This state has the same three general community phases. These non-native species can be highly flammable, and promote fire where historically fire had been infrequent. Negative feedbacks enhance ecosystem resilience and contribute to the stability of the state. These feedbacks include the presence of all structural and functional groups, low fine fuel loads, and retention of organic matter and nutrients. Positive feedbacks decrease ecosystem resilience and stability of the state. These include the non-natives' high seed output, persistent seed bank, rapid growth rate, ability to cross pollinate, and adaptations for seed dispersal.

Community Phase 2.1:

This community phase is similar to the Reference State Community Phase 1.1, with the presence of non-native species in trace amounts. Sagebrush, bottlebrush squirreltail and Sandberg bluegrass dominate the site. Forbs and other shrubs and grasses make up smaller components of this site.

Community Phase Pathway 2.1a, from Phase 2.1 to 2.2:

Fire reduces the shrub overstory and allows for perennial bunchgrasses to dominate the site. Fires are typically low severity resulting in a mosaic pattern due to low fuel loads. A fire following an unusually wet spring or a change in management favoring an increase in fine fuels may be more severe and reduce sagebrush cover to trace amounts. Annual non-native species are likely to increase after fire.

Community Phase Pathway 2.1b, from Phase 2.1 to 2.3:

Time and lack of disturbance allows for sagebrush to increase and become decadent. Long-term drought reduces fine fuels and leads to a reduced fire frequency, allowing sagebrush to dominate the site. Inappropriate grazing management reduces the perennial grass understory.

Community Phase 2.2:

This community phase is characteristic of a post-disturbance, early to mid-seral community where annual non-native species are present. Sagebrush is present in trace amounts; bottlebrush squirreltail, Sandberg bluegrass and other perennial bunchgrasses dominate the site. Depending on fire severity patches of intact sagebrush may remain. Rabbitbrush may be sprouting or dominant in the community. Perennial forbs may be a significant component for a number of years following fire. Annual non-native species are stable or increasing within the community.

Community Phase Pathway 2.2a, from Phase 2.2 to 2.1:

Time and lack of disturbance and/or grazing management that favors the establishment and growth of sagebrush allows the shrub component to recover. The establishment of low sagebrush can take many years.

Community Phase 2.3 (At Risk):

This community is at risk of crossing a threshold to another state. Sagebrush dominates the overstory and perennial bunchgrasses in the understory are reduced, either from competition with shrubs, from inappropriate grazing, or from both. Rabbitbrush may be a significant component. Bare ground increases. Annual non-native species may be stable or increasing due to lack of competition with perennial bunchgrasses. Pinyon and/or juniper may be present. This site is susceptible to further degradation from grazing, drought, and fire.

Community Phase Pathway 2.3a, from Phase 2.3 to 2.2:

Fire eliminates/reduces the overstory of sagebrush and allows for the understory perennial grasses to increase. Fires may be high severity in this community phase due to the dominance of sagebrush resulting in removal of overstory shrub community. Annual non-native species respond well to fire and may increase post burn.

T2A: Transition from Current Potential State 2.0 to Shrub State 3.0:

Trigger: To Community Phase 3.1: Inappropriate grazing will decrease or eliminate deep rooted perennial bunchgrasses, increase Sandberg bluegrass and favor shrub growth and establishment. To Community Phase 3.2: Severe fire in Community Phase 2.3 will remove sagebrush overstory. Annual non-native species will increase. Slow variables: Long term decrease in deep-rooted perennial grass density.

Threshold: Loss of deep-rooted perennial bunchgrasses changes nutrient cycling, nutrient redistribution, and reduces soil organic matter.

T2B: Transition from Current Potential State 2.0 to Annual State 4.0:

Trigger: Fire or soil disturbing treatment would transition to Community Phase 4.1.

Slow variables: Increased production and cover of non-native annual species.

Threshold: Loss of deep-rooted perennial bunchgrasses and shrubs changes temporal and spatial nutrient capture and cycling within the community. Increased, continuous fine fuels modify the fire regime by increasing frequency, size and spatial variability of fires.

Shrub State 3.0:

This state has two community phases, a shrub-dominated phase and a grass-dominated phase. This state is a product of many years of heavy grazing during time periods harmful to perennial bunchgrasses. Sagebrush dominates the overstory and rabbitbrush may be a significant component. Sagebrush cover exceeds site concept and may be decadent, reflecting stand maturity and lack of seedling establishment due to competition with mature plants. The shrub overstory and bottlebrush squirreltail or western wheatgrass understory dominate site resources such that soil water, nutrient capture, nutrient cycling and soil organic matter are temporally and spatially redistributed.

Community Phase 3.1 (At Risk):

Decadent sagebrush dominates the overstory. Rabbitbrush may be a significant component. Deep-rooted perennial bunchgrasses may be present in trace amounts or absent from the community. Western wheatgrass and annual non-native species increase. Bare ground is significant. Pinyon and/or juniper may be present.

Community Phase Pathway 3.1a, from Phase 3.1 to 3.2:

Fire, heavy fall grazing causing mechanical damage to shrubs, and/or brush treatments with minimal soil disturbance, will greatly reduce the overstory shrubs to trace amounts and allow grasses to dominate the site.

Community Phase 3.2:

Bottlebrush squirreltail and/or western wheatgrass dominate the site; annual non-native species may be present but are not dominant. Trace amounts of sagebrush or rabbitbrush may be present. Bare ground may be significant.

Community Phase Pathway 3.2a, from Phase 3.2 to 3.1:

Time and lack of disturbance and/or grazing management that favors the establishment and growth of sagebrush allows the shrub component to recover. The establishment of low sagebrush can take many years.

T3A: Transition from Shrub State 3.0 to Annual State 4.0:

Trigger: Fire and/or treatments that disturb the soil and existing plant community.

Slow variables: Increased seed production (following a wet spring) and cover of annual non-native species.

Threshold: Increased, continuous fine fuels modify the fire regime by changing frequency, intensity, size and spatial variability of fires. Changes in plant community composition and spatial variability of vegetation due to the loss of perennial bunchgrasses and sagebrush truncate energy capture and impact the temporal and spatial aspects of nutrient cycling and distribution.


Annual State 4.0:

This state has one community phase dominated by annual plants. An abiotic threshold has been crossed and state dynamics are driven by fire and time. The herbaceous understory is dominated by annual non-native species such as cheatgrass, medusahead, Russian thistle, and mustards. Resiliency has declined and further degradation from fire facilitates a cheatgrass and sprouting shrub plant community. Fire return interval has shortened due to the dominance of annual grasses in the understory and is a driver in site dynamics.

Community Phase 4.1:

Annual plants like cheatgrass, medusahead, and Russian thistle dominate. Bottlebrush squirreltail and perennial forbs may still be present in trace amounts. Surface erosion may increase with summer convection storms and would be evidenced by increased pedestalling of plants, rill formation, or extensive water flow paths.

State and transition model



Reference State 1.0 Community Phase Pathways

1.1a: Low severity fire creates sagebrush/grass mosaic; high severity fire significantly reduces sagebrush cover and leads to early/mid-seral community, dominated by grasses and forbs.

1.1b: Time and lack of disturbance such as fire. Excessive herbivory and/or long-term drought may also reduce perennial understory.

1.2a: Time and lack of disturbance allows for shrub regeneration.

1.3a: Low severity fire would create sagebrush/grass mosaic.

1.3b: High severity fire significantly reduces sagebrush cover leading to early/mid-seral community.

Transition T1A: Introduction of non-native annual species.

Current Potential State 2.0 Community Phase Pathways

2.1a: Low severity fire creates sagebrush/grass mosaic; high severity fire significantly reduces sagebrush cover and leads to early/mid-seral community dominated by grasses and forbs; non-native annual species present.

2.1b: Time and lack of disturbance. Inappropriate grazing management and/or long-term drought may also reduce perennial understory.

2.2a: Time and lack of disturbance allows for regeneration of sagebrush.

2.3a: Low severity fire creates sagebrush/grass mosaic, herbivory or combinations. Brush management with minimal soil disturbance reduces sagebrush.

Transition T2A: Inappropriate grazing management (3.1), or high severity fire (3.2).

Transition T2B: Fire or brush management causing severe soil disturbance.

Shrub State 3.0 Community Phase Pathways

3.1a: Low severity fire.

3.2a: Time and lack of disturbance.

Transition T3A: Catastrophic fire and/or treatments that disturb the existing plant community.

Annual State 4.0 Community Phase Pathways

None.

State 1**Reference Plant Community****Community 1.1****Reference Plant Community**

The reference plant community is dominated by Wyoming big sagebrush, western wheatgrass, bottlebrush squirreltail, Sandberg's bluegrass and littleleaf horsebrush. Potential vegetative composition is about 50% grasses, 5% forbs and 45% shrubs. Approximate ground cover (basal and crown) is 15 to 25 percent.

Table 5. Annual production by plant type

| Plant Type | Low (Kg/Hectare) | Representative Value (Kg/Hectare) | High (Kg/Hectare) |
|-----------------|---------------------|--------------------------------------|----------------------|
| Grass/Grasslike | 280 | 336 | 448 |
| Shrub/Vine | 252 | 303 | 404 |
| Forb | 28 | 34 | 45 |
| Total | 560 | 673 | 897 |

Additional community tables

Table 6. Community 1.1 plant community composition

| Group | Common Name | Symbol | Scientific Name | Annual Production (Kg/Hectare) | Foliar Cover (%) |
|------------------------|------------------------------------|--------|------------------------------------|--------------------------------|------------------|
| Grass/Grasslike | | | | | |
| 1 | Primary Perennial Grasses | | | 168–404 | |
| | western wheatgrass | PASM | <i>Pascopyrum smithii</i> | 101–235 | – |
| | squirreltail | ELEL5 | <i>Elymus elymoides</i> | 34–101 | – |
| | Sandberg bluegrass | POSE | <i>Poa secunda</i> | 34–67 | – |
| 2 | Secondary Perennial Grasses | | | 13–34 | |
| | Thurber's needlegrass | ACTH7 | <i>Achnatherum thurberianum</i> | 3–13 | – |
| Forb | | | | | |
| 3 | Perennial | | | 13–54 | |
| | onion | ALLIU | <i>Allium</i> | 3–13 | – |
| | sunflower | HELIA3 | <i>Helianthus</i> | 3–13 | – |
| 4 | Annual | | | 0–20 | |
| Shrub/Vine | | | | | |
| 5 | Primary Shrubs | | | 81–323 | |
| | littleleaf horsebrush | TEGL | <i>Tetradymia glabrata</i> | 34–101 | – |
| | rubber rabbitbrush | ERNA10 | <i>Ericameria nauseosa</i> | 13–54 | – |
| 6 | Secondary Shrubs | | | 13–67 | |
| | yellow rabbitbrush | CHV18 | <i>Chrysothamnus viscidiflorus</i> | 7–20 | – |

Animal community

Livestock Interpretations:

This site has limited suitability for grazing by livestock during the summer and fall. Grazing management should be keyed to western wheatgrass and Sandberg's bluegrass. Western wheatgrass provides important forage for domestic sheep. Fall regrowth cures well on the stem, so western wheatgrass is good winter forage for domestic livestock. Bottlebrush squirreltail is very palatable winter forage for domestic sheep of Intermountain ranges. Domestic sheep relish the green foliage. Overall, bottlebrush squirreltail is considered moderately palatable to livestock. Sandberg bluegrass is a widespread forage grass. It is one of the earliest grasses in the spring and is sought by domestic livestock and several wildlife species. Sandberg bluegrass is a palatable species, but its production is closely tied to weather conditions. It produces little forage in drought years, making it a less dependable food source than other perennial bunchgrasses. Livestock browse Wyoming big sagebrush, but may use it only lightly when palatable herbaceous species are available. In general, livestock forage only lightly on rubber rabbitbrush during the summer, but winter use can be heavy in some locations. Fall use is variable, but flowers are often used by livestock. A few leaves and the more tender stems may also be used.

Stocking rates vary over time depending upon season of use, climate variations, site, and previous and current management goals. A safe starting stocking rate is an estimated stocking rate that is fine tuned by the client by adaptive management through the year and from year to year.

Wildlife Interpretations:

Small non-gamebirds, rodents and deer may use this site. Wyoming big sagebrush is preferred browse for wild ungulates, and Wyoming big sagebrush communities are important winter ranges for big game. Pronghorn usually browse Wyoming big sagebrush heavily. Wyoming big sagebrush communities are critical habitat for the birds. Sage-grouse: Sagebrush-grassland communities provide critical sage-grouse breeding and nesting habitats. Meadows surrounded by sagebrush may be used as feeding and strutting grounds. Sagebrush is a crucial component of their diet year-round, and sage-grouse select sagebrush almost exclusively for cover. Sage-grouse prefer mountain big sagebrush and Wyoming big sagebrush communities to basin big sagebrush communities. Wildlife forage only lightly on rubber rabbitbrush during the summer, but winter use can be heavy in some locations. Fall use is variable, but flowers are often used by wildlife. A few leaves and the more tender stems may also be used. The forage value of rubber rabbitbrush varies greatly among subspecies and ecotypes. Elk consume western

wheatgrass during the fall, winter, spring, and summer. Western wheatgrass is used by various small mammals. Bottlebrush squirreltail is a dietary component of several wildlife species. Bottlebrush squirreltail may provide forage for mule deer and pronghorn. Sandberg bluegrass is desirable for pronghorn antelope and mule deer in the spring and preferable in the spring, summer, and fall for elk and desirable as part of their winter range.

Hydrological functions

Runoff is very high. Permeability is very slow.

Recreational uses

Aesthetic value is derived from the diverse floral and faunal composition and the colorful flowering of wild flowers and shrubs during the spring and early summer. This site offers rewarding opportunities to photographers and for nature study. This site is used for camping and hiking and has potential for upland and big game hunting.

Other products

Native Americans made tea from big sagebrush leaves. They used the tea as a tonic, an antiseptic, for treating colds, diarrhea, and sore eyes and as a rinse to ward off ticks. Big sagebrush seeds were eaten raw or made into meal.

Other information

Wyoming big sagebrush is used for stabilizing slopes and gullies and for restoring degraded wildlife habitat, rangelands, mine spoils and other disturbed sites. It is particularly recommended on dry upland sites where other shrubs are difficult to establish.

Horsebrush provide critically needed ground cover and protection from erosion on dry sites that are otherwise often sparsely vegetated.

Bottlebrush squirreltail is tolerant of disturbance. Western wheatgrass is a good soil binder and is well suited for reclamation of disturbed sites such as erosion control and soil stabilization.

Inventory data references

NASIS data for soil survey areas NV628, NV772, and NV773.

Type locality

| | |
|------------------------------------|--|
| Location 1: Carson City County, NV | |
| General legal description | This site also occurs in Douglas, Lyon, Mineral, Storey and Washoe counties, Nevada. |

References

Stringham, T.K., D. Snyder, P. Novak-Echenique, K. O'Neill, A. Lyons, and M. Johns. 2021. Great Basin Ecological Site Development Project: State-and-Transition Models for Major Land Resource Area 26, Nevada and Portions of California..

Other references

Fire Effects Information System (Online; <http://www.fs.fed.us/database/feis/plants/>).

USDA-NRCS Plants Database (Online; <http://www.plants.usda.gov>).

Contributors

DK/FR

Tamzen Stringham

Approval

Kendra Moseley, 4/10/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

| | |
|---|---------------------------------------|
| Author(s)/participant(s) | Patti Novak-Echenique |
| Contact for lead author | State Rangeland Management Specialist |
| Date | 07/18/2013 |
| Approved by | Kendra Moseley |
| Approval date | |
| Composition (Indicators 10 and 12) based on | Annual Production |

Indicators

- 1. Number and extent of rills:** Rills are none. If rills do develop, they are broken up by the churning action of the soil.

- 2. Presence of water flow patterns:** Water flow patterns are rare. Water flow patterns are broken up by the churning action of the soil.

- 3. Number and height of erosional pedestals or terracettes:** Pedestals are none to rare. Occurrence is usually limited to areas of water flow patterns.

- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare Ground 30-50% depending on amount of surface rock fragments

- 5. Number of gullies and erosion associated with gullies:** None

- 6. Extent of wind scoured, blowouts and/or depositional areas:** None

- 7. Amount of litter movement (describe size and distance expected to travel):** Fine litter (foliage from grasses and annual & perennial forbs) expected to move distance of slope length during intense summer convection storms or rapid snowmelt events. Persistent litter (large woody material) will remain in place except during large rainfall events.

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil stability values should be 3 to 6 on most soil textures found on this site. (To be field tested.)
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Surface structure is typically medium prismatic. Soil surface colors are dark grayish-brown and soils have an ochric epipedon. Organic matter of the surface 2 to 4 inches is typically 1 to 3 percent dropping off quickly below. Organic matter content can be more or less depending on micro-topography.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Perennial herbaceous plant slow runoff and increase infiltration. Shrub canopy and associated litter break raindrop impact and provide opportunity for snow catch and accumulation on site.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None. Heavy clay horizons or prismatic structure are not to be interpreted as compacted.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Rhizomatous, cool season, perennial bunchgrasses (western wheatgrass)
- Sub-dominant: tall shrubs (big sagebrush)>> shallow-rooted, cool season, perennial bunchgrasses = associated shrubs
> deep-rooted, cool season perennial forbs > shallow-rooted, cool season, perennial and annual forbs
- Other:
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Dead branches within individual shrubs common and standing dead shrub canopy material may be as much as 25% of total woody canopy; some of the mature bunchgrasses (<20%) have dead centers.
-
14. **Average percent litter cover (%) and depth (in):** Between plant interspaces 25-35% and litter depth is < ¼ inch. Fine litter falls or blows into the soil surface cracks
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** For normal or average growing season (through June) ± 600 lbs/ac; Favorable years ± 800 lbs/ac and unfavorable years ± 500 lbs/ac
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that**

become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is **NOT** expected in the reference state for the ecological site: Potential invaders include cheatgrass, halogeton, Russian thistle, annual mustards and knapweeds.

17. **Perennial plant reproductive capability:** All functional groups should reproduce in average (or normal) and above average growing season years. Reduced growth and reproduction occur during extended or extreme drought periods.
-