

## Ecological site R028AY011NV SODIC DUNE

Accessed: 05/07/2024

### Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	P Novak-Echenique
Contact for lead author	State Rangeland Management Specialist
Date	10/24/2013
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### Indicators

1. **Number and extent of rills:** None
- 

2. **Presence of water flow patterns:** None
- 

3. **Number and height of erosional pedestals or terracettes:** Pedestals are few to common with occurrence due to wind scouring. After a wildfire, the remaining vegetation may become severely pedestalled.
- 

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare Ground up to 70%.
- 

5. **Number of gullies and erosion associated with gullies:** None
- 

6. **Extent of wind scoured, blowouts and/or depositional areas:** Slight to moderate wind scouring. Severe blowouts and flattening of dunes may occur after a severe wildfire and the resulting loss of vegetation.
- 

7. **Amount of litter movement (describe size and distance expected to travel):** Fine litter (foliage from grasses and

annual & perennial forbs) expected to move unsheltered distance during heavy wind. Persistent litter (large woody material) expected to remain in place except during intense summer convection storms.

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil stability values should be 1 to 4 on the sandy soil textures found on this site.  

---
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Surface structure is typically single grained. Soil surface colors are very pale browns and soils are typified by an ochric epipedon. Surface textures are fine sands. Organic matter of the surface 2 to 3 inches is typically less than 1.5 percent. Organic matter content can be more or less depending on micro-topography.  

---
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Perennial herbaceous plants (especially deep-rooted bunchgrasses [i.e., Indian ricegrass] slow runoff and increase infiltration. Shrub canopy and associated litter break raindrop impact and provide some opportunity for snow catch and accumulation on site.  

---
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** Compacted layers are none.  

---
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**  
  
Dominant: Reference State: tall salt-desert shrubs >  
  
Sub-dominant: deep-rooted, cool season, perennial bunchgrasses (i.e. Indian ricegrass) > warm season bunchgrasses > rhizomatous grasses > deep-rooted, cool season, perennial forbs.  
  
Other: Annual forbs  
  
Additional:  

---
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Dead branches within individual shrubs common and standing dead shrub canopy material may be as much as 30% of total woody canopy; some of the mature bunchgrasses ( $\pm 20\%$ ) have dead centers.  

---
14. **Average percent litter cover (%) and depth ( in):** Between plant interspaces ( $\pm 10\text{-}15\%$ ) and depth of litter is  $< \frac{1}{4}$  inch.  

---
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** For normal or average growing season (thru June)  $\pm 500$  lbs/ac; Favorable years  $\pm 700$  lb/ac and unfavorable years  $\pm 350$  lbs/ac.  

---

16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Potential invaders include cheatgrass, halogeton, Russian thistle and annual mustards.
- 

17. **Perennial plant reproductive capability:** All functional groups should reproduce in average (or normal) and above average growing season years. Little growth or reproduction occurs in extreme or extended drought periods.
-