

Ecological site R028AY236UT

Semidesert Shallow Loam (Black Sagebrush)

Accessed: 05/16/2024

General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

Associated sites

R028AY220UT	Semidesert Loam (Wyoming Big Sagebrush)
R028AY230UT	Semidesert Shallow Hardpan (Black Sagebrush)
R028AY234UT	Semidesert Shallow Loam (Utah juniper-Salina wildrye)

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Artemisia nova</i>
Herbaceous	Not specified

Physiographic features

This site occurs on hillsides, mountains, and to a lesser extent on pediments.

Table 2. Representative physiographic features

Landforms	(1) Hill (2) Mountain slope (3) Pediment
Elevation	1,372–2,042 m
Slope	5–45%

## Climatic features

The climate is semi-arid and characterized by cold snowy winters and warm dry summers. The average annual precipitation is 8 to 12 inches. Approximately 70 percent comes as rain from March through October. On the average, June through September are the driest months and March through May are the wettest months.

Mean Annual Air Temperature: 45-50

Mean Annual Soil Temperature: 47-52

**Table 3. Representative climatic features**

Frost-free period (average)	0 days
Freeze-free period (average)	150 days
Precipitation total (average)	305 mm

## Influencing water features

### Soil features

The characteristic soils in this site are 10 to 20 inches deep over bedrock and well drained.

They formed in colluvium and residuum derived mainly from limestone, sandstone, siltstone, and basalt parent materials the surface horizon is loam texture and 4 inches thick. About 55 percent of the soil surface is covered by rock fragments. The volume of rock fragments in the soil profile is 35 to 60 percent.

The rock fragments in these soils are dominated by pebbles in most soils, but cobbles and stones predominate in some soils. These soils are calcareous throughout and permeability is generally moderate. The available water capacity is 1 to 3 inches.

The water supplying capacity is 1 to 4 inches. Natural geologic erosion in potential is approximately 3 tons/acre/year.

**Table 4. Representative soil features**

Surface texture	(1) Loam
Drainage class	Well drained
Permeability class	Moderate
Soil depth	25–51 cm
Surface fragment cover ≤3"	28%
Surface fragment cover >3"	28%
Available water capacity (0-101.6cm)	2.54–7.62 cm
Subsurface fragment volume ≤3" (Depth not specified)	18–30%
Subsurface fragment volume >3" (Depth not specified)	18–30%

Ecological dynamics


As ecological condition deteriorates due to overgrazing, bluebunch wheatgrass and Indian ricegrass decrease, while black sagebrush, low rabbitbrush and snakeweed increase.

When the potential natural plant community is burned, bluebunch wheatgrass and Indian ricegrass decrease while low rabbitbrush and Sandberg bluegrass increase.


Cheatgrass and annual forbs are most likely to invade this site.

State and transition model

Ecosystem states



State 1 submodel, plant communities



State 1  
Reference State

Community 1.1  
Reference State

The dominant aspect of this plant community is black sagebrush. The composition by air-dry weight is approximately 45 percent perennial grasses, 5 percent forbs, and 50 percent shrubs.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Forb	196	295	392
Grass/Grasslike	177	265	353
Shrub/Vine	196	29	39
Total	569	589	784

Table 6. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	15-30%
Grass/grasslike foliar cover	10-30%
Forb foliar cover	2-3%
Non-vascular plants	0%
Biological crusts	0%
Litter	0%

Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%

Table 7. Canopy structure (% cover)

Height Above Ground (M)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.15	—	—	—	—
>0.15 <= 0.3	—	—	—	—
>0.3 <= 0.6	—	25-35%	25-35%	0-5%
>0.6 <= 1.4	—	—	—	—
>1.4 <= 4	—	—	—	—
>4 <= 12	—	—	—	—
>12 <= 24	—	—	—	—
>24 <= 37	—	—	—	—
>37	—	—	—	—

Figure 4. Plant community growth curve (percent production by month).  
UT2361, PNC. Excellent Condition.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	5	15	40	30	5	5	0	0	0	0

### Additional community tables

Table 8. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
<b>Shrub/Vine</b>					
0	<b>Primary Shrubs</b>			155–216	
	black sagebrush	ARNO4	<i>Artemisia nova</i>	155–216	–
3	<b>Secondary Shrubs</b>			62–93	
	shadscale saltbush	ATCO	<i>Atriplex confertifolia</i>	7–19	–
	yellow rabbitbrush	CHVI8	<i>Chrysothamnus viscidiflorus</i>	7–19	–
	Nevada jointfir	EPNE	<i>Ephedra nevadensis</i>	7–19	–
	broom snakeweed	GUSA2	<i>Gutierrezia sarothrae</i>	7–19	–
	winterfat	KRLA2	<i>Krascheninnikovia lanata</i>	7–19	–
	granite prickly phlox	LIPU11	<i>Linanthus pungens</i>	7–19	–
	plains pricklypear	OPPO	<i>Opuntia polyacantha</i>	7–19	–
	bud sagebrush	PIDE4	<i>Picrothamnus desertorum</i>	7–19	–
	Mexican cliffrose	PUME	<i>Purshia mexicana</i>	7–19	–
	stemless mock goldenweed	STAC	<i>Stenotus acaulis</i>	7–19	–
	Nuttall's horsebrush	TENU2	<i>Tetradymia nuttallii</i>	7–19	–
<b>Grass/Grasslike</b>					
0	<b>Primary Grasses</b>			216–308	
	bluebunch wheatgrass	PSSP6	<i>Pseudoroegneria spicata</i>	123–155	–
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	62–93	–
	needle and thread	HECO26	<i>Hesperostipa comata</i>	31–62	–
1	<b>Secondary Grasses</b>			19–31	
	purple threeawn	ARPU9	<i>Aristida purpurea</i>	7–19	–
	squirreltail	ELEL5	<i>Elymus elymoides</i>	7–19	–
	James' galleta	PLJA	<i>Pleuraphis jamesii</i>	7–19	–
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	7–19	–
<b>Forb</b>					
2	<b>Forbs</b>			31–62	
	freckled milkvetch	ASLE8	<i>Astragalus lentiginosus</i>	7–19	–
	Hooker's balsamroot	BAHO	<i>Balsamorhiza hookeri</i>	7–19	–
	cushion buckwheat	EROV	<i>Eriogonum ovalifolium</i>	7–19	–
	shaggy fleabane	ERPU2	<i>Erigeron pumilus</i>	7–19	–
	ballhead ipomopsis	IPCOC3	<i>Ipomopsis congesta</i> ssp. <i>congesta</i>	7–19	–
	woolly groundsel	PACA15	<i>Packera cana</i>	7–19	–
	low beardtongue	PEHU	<i>Penstemon humilis</i>	7–19	–
	spiny phlox	PHHO	<i>Phlox hoodii</i>	7–19	–
	gooseberryleaf globemallow	SPGR2	<i>Sphaeralcea grossulariifolia</i>	7–19	–
	heartleaf twistflower	STCO6	<i>Streptanthus cordatus</i>	7–19	–

## Animal community

This site is suited for grazing by cattle and sheep during fall, winter, and spring.

Wildlife using this site include rabbit, coyote, fox, badger, pronghorn antelope, mule deer, and dove.

This is a short list of the more common species found. Many other species are present as well and migratory birds are present at times.

## Hydrological functions

Soils are in hydrologic group D with runoff curves ranging from 80 to 89 depending on hydrologic condition.

## Recreational uses

Resources that have special aesthetic and landscape values are wildflowers. Some recreation uses of this site are hiking and horseback riding.

## Wood products

None

## Other information

Threatened and endangered species include plants and animals.

## Type locality

Location 1: Box Elder County, UT	
Township/Range/Section	T10N R14W S12
General legal description	S ½ of Section 12, Township 10N, Range 14W Box Elder County Utah Red Dome Allotment, Horse Hills, Photo Number 167 F-6. T. 10 N. R. 14 W. South ½ Section12

## Contributors

David J. Somorville  
DJS

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Jack Alexander, Range Specialist, Synergy Resource Solutions, Inc. Julia Kluck, Soil Scientist, Synergy Resource Solutions, Inc. Shane Green, State Range Specialist, Utah NRCS
Contact for lead author	Shane Green, Shane.Green@ut.usda.gov
Date	02/09/2010
Approved by	Shane A. Green
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:** Some rills present. Rill development may increase following large storm events, but should begin to heal during the following growing season. Frost heaving will accelerate recovery. Rill development may increase when run inflow enters site from other sites that produce large amounts of runoff (i.e. steeper sites, slickrock, rock outcrop).
- 

2. **Presence of water flow patterns:** Water flow patterns will be short (2-5') and meandering; interrupted by plants and exposed rocks. Some evidence of erosion or deposition associated with flow patterns. Where slopes exceed 5%, water flow patterns may be longer (5-10').
- 

3. **Number and height of erosional pedestals or terracettes:** Plants may have small pedestals (1-3") where they are adjacent to water flow patterns, but without exposed roots. Terracettes should be few and stable. Terracettes should be small (1-3") and show little sign of active erosion. Some plants may appear to have a pedestal but rather than be formed by erosion, the only place litter accumulates and soil collects is at plant bases forming the appearance of a pedestal.

Well-developed biological crusts may appear pedestalled, but are actually a characteristic of the crust formation. Some plants may appear to have a pedestal but rather than be formed by erosion, the only place litter accumulates and soil collects is at plant bases forming the appearance of a pedestal.

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 20-35% bare ground (soil with no protection from raindrop impact). Herbaceous communities are most likely to have lower values. As species composition by shrubs increases, bare ground is likely to increase. Poorly developed biological soil crust that is susceptible to raindrop splash erosion should be recorded as bare ground. Very few if any bare spaces of greater than 1 square foot.
- 

5. **Number of gullies and erosion associated with gullies:** No gullies present.
- 

6. **Extent of wind scoured, blowouts and/or depositional areas:** Very minor evidence of active wind-generated soil movement. Wind scoured (blowouts) and depositional areas are rarely present. If present they have muted features and are mostly stabilized with vegetation and/or biological crust. Gravel or desert pavement protects the site from wind scour.
- 

7. **Amount of litter movement (describe size and distance expected to travel):** Most litter resides in place with some redistribution caused by water and wind movement. Very minor litter removal may occur in flow patterns and rills with deposition occurring at points of obstruction. The majority of litter accumulates at the base of plants. Some leaves, stems, and small twigs may accumulate in soil depressions adjacent to plants. Woody stems are not likely to move.
- 

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** This site should have an erosion rating of 5 or 6 under plant canopies and a rating of 4 to 5 in the interspaces with an average rating of 5 using the soil stability kit test.
- 

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** This

description is based on the modal soil (Checkett GR-L CB-L, soil survey area: 634, Iron-Washington). This site has 5 correlated soils, resulting in variation of each of these attributes. Unless working on a location with the modal soil, it is critical to supplement this description with the soil-specific information from the published soil survey.

Soil surface horizon is typically 2 to 3 inches deep. Structure is typically very thin platy. Color is typically pale brown (10YR 6/3), dark grayish brown (10YR 4/2) moist. An ochric horizon extends to a depth of 3 inches. An ochric horizon typically extends to a depth of 2 to 10 inches. The ochric horizon is a surface horizon lacking fine stratification and which is either light colored, or thin, or has a low organic carbon content, or is massive and (very) hard when dry. The A horizon would be expected to be more strongly developed under plant canopies. It is important if you are sampling to observe the A horizon under plant canopies as well as the interspaces.

- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Bunchgrasses and shrubs equally important for increasing infiltration and reducing runoff. Litter plays a role in increasing infiltration and decreasing runoff. Plants provide microhabitat for seedlings, catch litter and soil, and slow raindrops and runoff. Vascular plants and/or well-developed biological soil crusts (where present) will break raindrop impact and splash erosion. Spatial distribution of vascular plants and interspaces between well-developed biological soil crusts (where present) provide detention storage and surface roughness that slows runoff allowing time for infiltration. Interspaces between plants and any well-developed biological soil crusts (where present) may serve as water flow patterns during episodic runoff events, with natural erosion expected in severe storms. When perennial grasses decrease, reducing ground cover and increasing bare ground, runoff is expected to increase and any associated infiltration reduced. Shrubs catch snow, slow wind evaporation, and provide microhabitat for seedling establishment.

- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None. Lithic contact (the boundary with hard bedrock) may occur at about 19 inches. Naturally occurring soil horizons may be harder than the surface because of an accumulation of calcium carbonate and should not be considered as compaction layers.

- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: black sagebrush

Sub-dominant: bluebunch wheatgrass > Indian ricegrass > needle and thread

Other: other shrubs, other grasses, forbs

Additional: In the northern portion of the MLRA cool-season perennial grasses (Indian ricegrass, needle and thread) dominate. In the southernmost portion of the MLRA warm-season perennial grasses (galleta, sand dropseed) dominate. The two groups share dominance in the middle portion of the MLRA.

- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** During years with average to above average precipitation, there should be very little recent mortality or decadence apparent in either the shrubs or grasses. Some bunchgrass and shrub mortality may occur during severe droughts, particularly on the shallower and coarser soils associated with this site.
-

14. **Average percent litter cover (%) and depth ( in):** Litter cover includes litter under plants. Most litter will be fine litter. Depth should be 1-2 leaf thickness in the interspaces and up to 1/2" under canopies. Litter cover may increase to 15-25% following years with favorable growing conditions. Excess litter may accumulate in absence of disturbance. Vegetative production may be reduced if litter cover exceeds 40%.
- 

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 525#/acre.  
Even the most stable communities exhibit a range of production values. Production will vary between communities and across the MRLA. Refer to the community descriptions in the ESD. Production will differ across the MLRA due to the naturally occurring variability in weather, soils, and aspect. The biological processes on this site are complex; therefore, representative values are presented in a land management context.
- 

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Cheatgrass and annual forbs
- 

17. **Perennial plant reproductive capability:** All perennial plants should have the ability to reproduce sexually or asexually, except in drought years. Density of plants indicates that plants reproduce at level sufficient to fill available resource. Within capability of site there are no restrictions on seed or vegetative reproductive capacity.
-