

Ecological site R028BY050NV

WET SODIC BOTTOM

Last updated: 2/19/2025

Accessed: 12/22/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 028B—Central Nevada Basin and Range

MLRA 28B occurs entirely in Nevada and comprises about 23,555 square miles (61,035 square kilometers). More than nine-tenths of this MLRA is federally owned. This area is in the Great Basin Section of the Basin and Range Province of the Intermontane Plateaus. It is an area of nearly level, aggraded desert basins and valleys between a series of mountain ranges trending north to south. The basins are bordered by long, gently sloping to strongly sloping alluvial fans. The mountains are uplifted fault blocks with steep sideslopes. Many of the valleys are closed basins containing sinks or playas. Elevation ranges from 4,900 to 6,550 feet (1,495 to 1,995 meters) in the valleys and basins and from 6,550 to 11,900 feet (1,995 to 3,630 meters) in the mountains.

The mountains in the southern half are dominated by andesite and basalt rocks that were formed in the Miocene and Oligocene. Paleozoic and older carbonate rocks are prominent in the mountains to the north. Scattered outcrops of older Tertiary intrusives and very young tuffaceous sediments are throughout this area. The valleys consist mostly of alluvial fill, but lake deposits are at the lowest elevations in the closed basins. The alluvial valley fill consists of cobbles, gravel, and coarse sand near the mountains in the apex of the alluvial fans. Sands, silts, and clays are on the distal ends of the fans.

The average annual precipitation ranges from 4 to 12 inches (100 to 305 millimeters) in most areas on the valley floors. Average annual precipitation in the mountains ranges from 8 to 36 inches (205 to 915 millimeters) depending on elevation. The driest period is from midsummer to midautumn. The average annual temperature is 34 to 52 degrees F (1 to 11 degrees C). The freeze-free period averages 125 days and ranges from 80 to 170 days, decreasing in length with elevation.

The dominant soil orders in this MLRA are Aridisols, Entisols, and Mollisols. The soils in the area dominantly have a mesic soil temperature regime, an aridic or xeric soil moisture

regime, and mixed or carbonatic mineralogy. They generally are well drained, loamy or loamyskeletal, and shallow to very deep.

Nevada's climate is predominantly arid, with large daily ranges of temperature, infrequent severe storms and heavy snowfall in the higher mountains. Three basic geographical factors largely influence Nevada's climate: continentality, latitude, and elevation. The strong continental effect is expressed in the form of both dryness and large temperature variations. Nevada lies on the eastern, lee side of the Sierra Nevada Range, a massive mountain barrier that markedly influences the climate of the State. The prevailing winds are from the west, and as the warm moist air from the Pacific Ocean ascend the western slopes of the Sierra Range, the air cools, condensation occurs and most of the moisture falls as precipitation. As the air descends the eastern slope, it is warmed by compression, and very little precipitation occurs. The effects of this mountain barrier are felt not only in the West but throughout the state, as a result the lowlands of Nevada are largely desert or steppes.

The temperature regime is also affected by the blocking of the inland-moving maritime air. Nevada sheltered from maritime winds, has a continental climate with well-developed seasons and the terrain responds quickly to changes in solar heating. Nevada lies within the midlatitude belt of prevailing westerly winds which occur most of the year. These winds bring frequent changes in weather during the late fall, winter and spring months, when most of the precipitation occurs.

To the south of the mid-latitude westerlies, lies a zone of high pressure in subtropical latitudes, with a center over the Pacific Ocean. In the summer, this high-pressure belt shifts northward over the latitudes of Nevada, blocking storms from the ocean. The resulting weather is mostly clear and dry during the summer and early fall, with occasional thundershowers. The eastern portion of the state receives noteworthy summer thunderstorms generated from monsoonal moisture pushed up from the Gulf of California, known as the North American monsoon. The monsoon system peaks in August and by October the monsoon high over the Western U.S. begins to weaken and the precipitation retreats southward towards the tropics (NOAA 2004).

Ecological site concept

This site occurs on axial-stream floodplains. Slope gradients of 0 to 2 percent are most typical. Elevations are 4600 to 4900 feet.

Average annual precipitation is 5 to 8 inches. Mean annual air temperature is 45 to 50 degrees F. The average growing season is about 100 to 120 days.

The soils associated with this site are very deep and poorly drained. The soils are usually saturated with water during late winter and spring. During drier periods, the water table may lower to depths greater than 60 inches. In some cases, the soil surface is encrusted with water soluble salts, limiting seedling establishment.

The reference plant community is dominated by inland saltgrass. Potential vegetative composition is about 95% grasses, 2% forbs and 3% shrubs. Approximate ground cover

(basal and crown) is 60 to 80 percent.

Associated sites

R028BY020NV	SODIC FLAT 5-8 P.Z. SODIC FLAT 5-8 P.Z. Much less productive site; vegetation typically restricted to coppice mounds.
R028BY069NV	SODIC FLAT 8-10 P.Z. SODIC FLAT 8-10 P.Z. Less productive site; alkali sacaton minor grass; ground cover 10% or less.
R028BY004NV	SALINE BOTTOM This site occurs on lake plains and flood plains. Slopes gradients of 0 to 2 percent are typical and elevations range from 3400 to 7000 feet. The soils associated with this site are very deep, poorly drained, and are derived from silty alluvium, loess or lacustrine deposits from mixed rock sources. They are strongly salt and sodium affected in their upper profile with soil reaction and salinity decreasing with depth. Seasonally high water table can be found between 20 to 60 inches. The reference state is dominated by basin wildrye and alkali sacaton, with a component of black greasewood. The plant community is characterized by an open stand of perennial grasses, grass-like plants, forbs, with few shrubs. Production ranges from 800 to 2200 pounds per acre.

Similar sites

R028BY012NV	WET SALINE MEADOW PASM dominant grass; DISP increaser species; greater diversity of herbaceous vegetation
R028BY002NV	SALINE MEADOW SPAI-SPCR codominant grasses

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Distichlis spicata</i>

Physiographic features

This site occurs on axial-stream floodplains. Slope gradients of 0 to 2 percent are most typical. Elevations are 4600 to 4900 feet.

Table 2. Representative physiographic features

Landforms	(1) Flood plain
-----------	-----------------

Runoff class	High to very high
Flooding duration	Long (7 to 30 days)
Flooding frequency	Frequent
Ponding frequency	None
Elevation	1,402–1,494 m
Slope	0–2%
Water table depth	53–213 cm
Aspect	Aspect is not a significant factor

Climatic features

The climate associated with this site is semiarid, characterized by cold, moist winters and warm, dry summers.

Average annual precipitation ranges from 5 to 8 inches. Mean annual air temperature is about 45 to 50 degrees F. The average growing season is about 100 to 120 days.

Table 3. Representative climatic features

Frost-free period (average)	110 days
Freeze-free period (average)	
Precipitation total (average)	178 mm




Figure 1. Monthly average minimum and maximum temperature




Figure 2. Annual precipitation pattern

Influencing water features

This site is adjacent to streams and may be flooded for long periods during spring runoff.

Soil features

The soils associated with this site are very deep and poorly drained. The soils are usually saturated with water during late winter and spring. During drier periods, the water table may lower to depths greater than 60 inches. In some cases, the soil surface is encrusted with water soluble salts, limiting seedling establishment. The soil series associated with this site include: Ocala and Variant.

Table 4. Representative soil features

Parent material	(1) Alluvium
Surface texture	(1) Silty clay loam (2) Silty clay
Family particle size	(1) Loamy
Drainage class	Poorly drained
Permeability class	Very slow
Soil depth	183–213 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	18.03–18.29 cm
Calcium carbonate equivalent (0-101.6cm)	5–25%

Electrical conductivity (0-101.6cm)	0–32 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	13–45
Soil reaction (1:1 water) (0-101.6cm)	7.9–9
Subsurface fragment volume <=3" (Depth not specified)	0%
Subsurface fragment volume >3" (Depth not specified)	0%


Ecological dynamics

As ecological condition declines, rubber rabbitbrush, seepweed and black greasewood increase.

Fire Ecology:

Fire in dry meadow communities often involves stand replacing fires. The fire return interval occurs every 0 to 35 years. Saltgrass rhizomes occur deep in the soil where they are insulated from the heat of most fires. Saltgrass survives fire by sending up new growth from rhizomes.

State and transition model



Transition T1A: Introduction of non-native species such as Russian thistle.

Animal community

Livestock Interpretations:

This site is suitable for livestock grazing. Grazing management should be keyed to perennial grass production. Saltgrass's value as forage depends primarily on the relative availability of other grasses of higher nutritional value and palatability. It can be an especially important late summer grass in arid environments after other forage grasses have deceased. Saltgrass is rated as a fair to good forage species only because it stays green after most other grasses dry. Livestock generally avoid saltgrass due to its coarse foliage. Saltgrass is described as an “increaser” under grazing pressure.

Stocking rates vary over time depending upon season of use, climate variations, site, and previous and current management goals. A safe starting stocking rate is an estimated stocking rate that is fine tuned by the client by adaptive management through the year and from year to year.

Wildlife Interpretations:

Saltgrass provides cover for a variety of bird species, small mammals, and arthropods and is on occasion used as forage for several big game wildlife species.

Hydrological functions

Runoff is high to very high. Permeability is very slow.

Recreational uses

Aesthetic value is derived from the diverse floral and faunal composition and the colorful flowering of wild flowers and shrubs during the spring and early summer. This site offers rewarding opportunities to photographers and for nature study. This site is used for camping and hiking and has potential for upland and big game hunting.

Other information

Given its extensive system of rhizomes and roots which form a dense sod, saltgrass is considered a suitable species for controlling wind and water erosion.

Inventory data references

NASIS soil component data.

Type locality

Location 1: Elko County, NV	
Township/Range/Section	T40N R70E S19
Latitude	41° 20' 21"
Longitude	114° 5' 22"
General legal description	NW ¼ SW ¼, About 6 miles northeast of Montello, along Thousand Springs Creek leading into Dake Reservoir, Elko County, Nevada. This site also occurs in White Pine County, Nevada.

Other references

Fire Effects Information System (Online; <http://www.fs.fed.us/database/feis/plants/>).

Houghton, J.G., C.M. Sakamoto, and R.O. Gifford. 1975. Nevada's Weather and Climate, Special Publication 2. Nevada Bureau of Mines and Geology, Mackay School of Mines, University of Nevada, Reno, NV.

National Oceanic and Atmospheric Administration. 2004. The North American Monsoon. Reports to the Nation. National Weather Service, Climate Prediction Center. Available online: <http://www.weather.gov/>

USDA-NRCS Plants Database (Online; <http://www.plants.usda.gov>).

Contributors

HA/MD/RK

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	12/22/2025
Approved by	Kendra Moseley
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

14. **Average percent litter cover (%) and depth (in):**

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

17. **Perennial plant reproductive capability:**
