

Ecological site R030XA010CA Calcareous Hill 3-5

Last updated: 10/21/2024
Accessed: 11/13/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 030X–Mojave Basin and Range

The Mojave Desert Major Land Resource Area (MLRA 30) is found in southern California, southern Nevada, the extreme southwest corner of Utah and northwestern Arizona within the Basin and Range Province of the Intermontane Plateaus. The Mojave Desert is a transitional area between hot deserts and cold deserts where close proximity of these desert types exert enough influence on each other to distinguish these desert types from the hot and cold deserts beyond the Mojave. Kottek et. al 2006 defines hot deserts as areas where mean annual air temperatures are above 64 F (18 C) and cold deserts as areas where mean annual air temperatures are below 64 F (18 C). Steep elevation gradients within the Mojave create islands of low elevation hot desert areas surrounded by islands of high elevation cold desert areas.

The Mojave Desert receives less than 10 inches of mean annual precipitation. Mojave Desert low elevation areas are often hyper-arid while high elevation cold deserts are often semi-arid with the majority of the Mojave being an arid climate. Hyper-arid areas receive less than 4 inches of mean annual precipitation and semi-arid areas receive more than 8 inches of precipitation (Salem 1989). The western Mojave receives very little precipitation during the summer months while the eastern Mojave experiences some summer monsoonal activity.

In summary, the Mojave is a land of extremes. Elevation gradients contribute to extremely hot and dry summers and cold moist winters where temperature highs and lows can fluctuate greatly between day and night, from day to day and from winter to summer. Precipitation falls more consistently at higher elevations while lower elevations can experience long intervals without any precipitation. Lower elevations also experience a low frequency of precipitation events so that the majority of annual precipitation may come in only a couple precipitation events during the whole year. Hot desert areas influence cold desert areas by increasing the extreme highs and shortening the length of below freezing events. Cold desert areas influence hot desert areas by increasing the extreme lows and increasing the length of below freezing events. Average precipitation and temperature values contribute little understanding to the extremes which govern wildland plant communities across the Mojave.

LRU notes

The Mojave Desert is currently divided into 5 Land Resource Units (LRUs). This ecological site is within the arid portions of the Mojave where precipitation primarily occurs during the winter months. The Arid Western Mojave LRU is designated by the 'XA' symbol within the ecological site ID. The Arid Western Mojave LRU is roughly equivalent to Western Mojave Basins and Western Mojave Low Ranges and Arid Footslopes of EPA Level IV Ecoregions.

Elevations range from 1650 to 4000 feet and precipitation is between 4 to 8 inches per year. The Arid Western Mojave LRU is distinguished from the Arid Eastern Mojave (XB) by the lack of summer precipitation which excludes many warm season plant species from occurring in this LRU. The 'XA' LRU is generally west of the Mojave River and the 117 W meridian (Hereford et. al 2004). Vegetation includes creosote bush, shadscale saltbush, Nevada jointfir, Joshua tree, and burrobush. At the upper elevations of the LRU, plant production and diversity are greater and blackbrush is a common dominant shrub. The Arid Western Mojave LRU generally lacks the diversity of yucca,

cacti and warm season species found in the Arid Eastern Mojave.

Classification relationships

NDDDB/Holland, R.F. 1986. Preliminary descriptions of the terrestrial natural communities of California - Mojave Creosote Bush Scrub.; J.O. Sawyer and T. Keeler-Wolf. 1995. Manual of California Vegetation - Creosotebush Series.

Ecological site concept

Copy to R030XA043CA. The Calcareous Hill ecological site is found among the hill and mountains landscape below 3000 feet (915 m) on rock pediments between 5 and 15 percent slope . Soils are shallow, have an argillic horizon and formed in residuum from granodiorite.

Please refer to group concept R030XA043CA to view the provisional STM.

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Larrea tridentata</i> (2) <i>Atriplex confertifolia</i>
Herbaceous	(1) <i>Achnatherum speciosum</i>

Physiographic features

Table 2. Representative physiographic features

Landforms	(1) Hill (2) Pediment
Elevation	2,500–2,800 ft
Slope	5–15%
Aspect	Aspect is not a significant factor

Climatic features

Table 3. Representative climatic features

Frost-free period (average)	250 days
Freeze-free period (average)	223 days
Precipitation total (average)	5 in




Figure 1. Monthly average minimum and maximum temperature

Influencing water features

The plant community for this site is not influenced by water from a wetland or stream.

Soil features

Table 4. Representative soil features

Surface texture	(1) Gravelly sandy loam
Family particle size	(1) Loamy
Drainage class	Well drained
Permeability class	Moderately slow
Soil depth	17–20 in
Surface fragment cover ≤3"	15%
Surface fragment cover >3"	5%
Available water capacity (0-40in)	0.1–0.13 in
Calcium carbonate equivalent (0-40in)	0–5%
Electrical conductivity (0-40in)	0–4 mmhos/cm
Sodium adsorption ratio (0-40in)	0–5
Soil reaction (1:1 water) (0-40in)	7.7–8
Subsurface fragment volume ≤3" (Depth not specified)	15%
Subsurface fragment volume >3" (Depth not specified)	5%

Ecological dynamics

Please refer to group concept R030XA043CA to view the provisional STM.

State and transition model

Ecosystem states

1. Reference State - Plant Community 1.1

State 1 submodel, plant communities

1.1. Reference State - Plant Community 1.1

State 1

Reference State - Plant Community 1.1

Community 1.1

Reference State - Plant Community 1.1

This site is the historic climax plant community. This community is characterized by widely spaced shrubs, 0.5 to 2 meters tall. Approximate ground cover (basal and crown) ranges from 5 to 15 percent. Creosotebush, shadscale and white bursage form the most characteristic association. Perennial grasses include desert needlegrass, Indian ricegrass and Sandberg bluegrass. Perennial forbs include desert trumpet and desert globemallow. The majority of the annuals are winter annuals which are especially abundant after winters of above average precipitation. The following table lists the major plant species and percentages by weight, air dry, of the total plant community that each contributes in an average production year. Fluctuations in species composition and relative production may change from year to year dependent upon abnormal precipitation or other climatic factors.

Forest overstory. **Allow no more than 3% of each shrub species and no more than 10% in aggregate.

Forest understory. **Allow no more than 2% of each grass species and no more than 5% in aggregate.

Other perennial grasses comprise 2 to 5% composition (air-dry weight)

Other annual grasses comprise trace to 5% composition (air-dry weight)

**Allow no more than 2% of each forb species and no more than 8% in aggregate.

Other perennial forbs comprise 2 to 8% composition (air-dry weight)

Other annual forbs comprise trace to 5% composition (air-dry weight)

Table 5. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Shrub/Vine	40	120	200
Forb	5	15	25
Grass/Grasslike	5	15	25
Total	50	150	250

Table 6. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	4-12%
Grass/grasslike foliar cover	0-1%
Forb foliar cover	0-1%
Non-vascular plants	0%
Biological crusts	0%
Litter	0%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%




Figure 3. Plant community growth curve (percent production by month). CA3002, Creosote bush XY. Growth starts in early spring, flowering and seed set occur by July. Dormancy occurs during the hot summer months. With sufficient summer/fall precipitation, some vegetation may break dormancy and produce a flush of growth..




Figure 4. Plant community growth curve (percent production by month). CA3003, Shadscale. Growth starts in early spring. Flowering and seed set occur by July. Seeds stay on the shrub for several months. Dormancy occurs during the hot summer months..

Additional community tables

Animal community

Wildlife Interpretations: This site provides suitable habitat for small mammals such as kangaroo rats and ground squirrels and game and fur mammals such as coyotes and black-tailed jackrabbits. Raptors and ravens also frequent this site. Common reptiles include side-blotched lizards and western whiptails. The soils on this site are poorly suited for burrowing reptiles such as the desert tortoise where depths are less than 10 inches.

Management for this site would be to protect it from excessive disturbance and maintain existing plant cover. Water developments would increase the species diversity.

This site has limited value for livestock grazing due to low forage production, lack of stockwater and the hazard of wind erosion. Grazing is limited to a few weeks in the spring when annual forbs and grasses produce additional forage in favorable years.

Pounds/acre AC/AUM

air dry

Normal Years 150 40-60

Hydrological functions

Runoff is medium. Hydrologic soil group D - soils having very slow infiltration rates when thoroughly wetted and consisting chiefly of clay soils with a high swelling potential, soils with a claypan or clay layer at or near the surface, and shallow soils over nearly impervious material. Hydrologic conditions: good - >70% ground cover (includes litter, grass and brush overstory); fair - 30 to 70% ground cover; poor - <30% ground cover.

Soil Series: Sparkhule
Hydrologic group: D
Hydrologic Conditions and Runoff Curves:
Good 84; Fair 86; Poor 88

Recreational uses

This site has value for open space, especially for off-road vehicle enthusiasts. Flowering wildflowers may also attract visitors in the spring.

Other information

Military Operations - Land clearing or other disturbances that destroys the vegetation and soil structure can result in increased erosion, soil blowing and barren areas. Off-road vehicles should be limited to existing roads and trails. Native species indigenous to this site are recommended for any revegetation efforts.

Inventory data references

Sampling technique

3 NV-ECS-1
 SCS-Range 417
 Other

Type locality

Location 1: San Bernardino County, CA	
Township/Range/Section	T10N R7W S13
General legal description	SE 1/4 Section 13, T10N, R7W East of Leuhman Ridge, Edwards Air Force Base

Other references

Hickman, J.C. (ed) 1995. The Jepson Manual Higher Plants of California.

Contributors

P. Novak-Echenique

Approval

Kendra Moseley, 10/21/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	

Date	11/13/2024
Approved by	Kendra Moseley
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
-