

Ecological site R030XA115AZ Sandy Wash 3-6" p.z.

Accessed: 05/03/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 030X–Mojave Basin and Range

AZ CRA 30.1 – Lower Mohave Desert

Elevations range from 400 to 2500 feet and precipitation averages 3 to 6 inches per year. Vegetation includes creosotebush, white bursage, Mormon tea, and brittlebush. The soil temperature regime is hyperthermic and the soil moisture regime is typic aridic. This unit occurs within the Basin and Range Province and is characterized by broad basins, valleys, and old lakebeds. Widely spaced mountains trending north to south occur throughout the area. Isolated, short mountain ranges are separated by an aggraded desert plain. The mountains are fault blocks that have been tilted up. Long alluvial fans coalesce with dry lakebeds between some of the ranges.

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Hymenoclea salsola</i> (2) <i>Acacia greggii</i>
Herbaceous	(1) <i>Pleuraphis rigida</i> (2) <i>Muhlenbergia porteri</i>

Physiographic features

This ecological site occurs as ephemeral washes. It occasionally receives additional run-in moisture from the surrounding ecological sites. These flows are generally of a short duration, rarely more than 1 or 2 days in duration.

Table 2. Representative physiographic features

Landforms	(1) Wash
Flooding duration	Very brief (4 to 48 hours)
Flooding frequency	Occasional
Ponding frequency	None
Elevation	122–610 m
Slope	0–3%
Aspect	Aspect is not a significant factor

Climatic features

The 30-1AZ Lower Mohave Desert Shrub land resource unit is characterized by a hot, dry climate. The average annual rainfall is 3 to 6 inches, but it can be extremely variable (e.g. from 0 to 11 inches). There can be long periods when little or no precipitation is received. Most of the precipitation for the year could arrive in just a couple of storms. The soil moisture regime is typic aridic and the soil temperature regime is hyperthermic. Winter precipitation from November through April occurs as gentle rains from storms coming out of the Pacific Ocean. Snow is very rare and only falls in the highest mountains. A seasonal drought occurs in May and June. Summer/fall precipitation from July through October comes from spotty, unreliable, and sometimes violent thunderstorms. The moisture originates in the Gulf of Mexico (and the Pacific Ocean in the fall) and flows into the state on the north end of the Mexican monsoon. Strong winds are common, especially during the spring.

Table 3. Representative climatic features

Frost-free period (average)	325 days
Freeze-free period (average)	365 days
Precipitation total (average)	152 mm

Influencing water features

Soil features

The soil of this ecological site is deep to very deep to any plant root restricting layer. The surface texture is loamy sand to coarse sand and is gravelly to very cobbly. Subsurface textures are loamy sand to coarse sand and is gravelly to extremely gravelly.

A typical soil profile is:

0 to 3 inches; very gravelly loamy sand

3 to 60 inches; very gravelly coarse sand and very gravelly loamy sand

This ecological site has been correlated to map unit 623015, Carrizo Series, Shivwits Area, AZ, Part of Mohave County SSA, and map units 701108 and 701109, Carrizo Series, Grand Canyon Area, AZ, Parts of Coconino and Mohave Counties SSA

Table 4. Representative soil features

Surface texture	(1) Gravelly loamy sand (2) Very gravelly sand (3) Very cobbly sandy clay
-----------------	---

Family particle size	(1) Sandy
Drainage class	Excessively drained
Permeability class	Very rapid
Soil depth	152 cm
Surface fragment cover <=3"	50–60%
Surface fragment cover >3"	5–45%
Available water capacity (0-101.6cm)	2.54–7.62 cm
Calcium carbonate equivalent (0-101.6cm)	5–10%
Electrical conductivity (0-101.6cm)	0–2 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0
Soil reaction (1:1 water) (0-101.6cm)	7.9–8.4
Subsurface fragment volume <=3" (Depth not specified)	40–60%
Subsurface fragment volume >3" (Depth not specified)	5–20%

Ecological dynamics

The plant communities found on an ecological site are naturally variable. Composition and production will vary with yearly conditions, location, aspect, and the natural variability of the soils. The historical climax plant community represents the natural potential plant communities found on relict or relatively undisturbed sites. Other plant communities described here represent plant communities that are known to occur when the site is disturbed by factors such as grazing, fire, or drought.

Production data provided in this site description is standardized to air-dry weight at the end of the summer growing season. The plant communities described in this site description are based on near normal rainfall years.

NRCS uses a Similarity Index to compare existing plant communities to the plant communities described here. Similarity Index is determined by comparing the production and composition of a plant community to the production and composition of a plant community described in this site description. To determine Similarity Index, compare the production (air-dry weight) of each species to that shown in the plant community description. For each species, count no more than the maximum amount shown for the species, and for each group, count no more than the maximum shown for the group. Divide the resulting total by the total normal year production shown in the plant community description. If rainfall has been significantly above or below normal, use the total production shown for above or below normal years. If field data is not collected at the end of the summer growing season, then the field data must be corrected to the end of the year production before comparing it to the site description. The growth curve can be used as a guide for estimating production at the end of the summer growing season.

State and transition model



**State 1
Historic Climax Plant Community**

**Community 1.1
Historic Climax Plant Community**

The dominant aspect of the site is a desert-shrub community. Grasses and forbs are present among the shrubs. Catclaw acacia can be the tallest shrub, especially in the drainage channels. White burrobrush is the dominant shrub. Big galleta and bush muhly are the major perennial grasses.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Shrub/Vine	280	336	364
Grass/Grasslike	112	151	168
Forb	56	73	84
Total	448	560	616

Table 6. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	0-2%
Grass/grasslike foliar cover	0%

Forb foliar cover	0%
Non-vascular plants	0%
Biological crusts	0%
Litter	0%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%

Table 7. Canopy structure (% cover)

Height Above Ground (M)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.15	–	–	–	0-2%
>0.15 <= 0.3	–	–	1-3%	–
>0.3 <= 0.6	–	–	–	–
>0.6 <= 1.4	–	6-10%	–	–
>1.4 <= 4	–	–	–	–
>4 <= 12	–	–	–	–
>12 <= 24	–	–	–	–
>24 <= 37	–	–	–	–
>37	–	–	–	–

Figure 4. Plant community growth curve (percent production by month). AZ3011, 30.1 3-6" p.z. all sites. Growth begins in late winter, most growth occurs in the spring..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	7	30	32	13	7	3	5	2	1	0	0

Figure 5. Plant community growth curve (percent production by month). AZ3083, 30.27 3-6" p.z. white bursage. Growth begins in early spring. Dormancy occurs during the hot summer months. With sufficient summer/fall precipitation, some plants may break dormancy and produce a flush of growth. Flowers and sets seed by July..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	5	20	40	20	15	0	0	0	0	0	0

Figure 6. Plant community growth curve (percent production by month). AZ3086, 30.27 3-6" p.z. big galleta. Growth begins in the late winter and early spring, goes dormant in May through June, most growth occurs during the summer rainy season..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	10	20	10	0	0	15	40	5	0	0	0

Figure 7. Plant community growth curve (percent production by month). AZ3090, 30.27 3-6" p.z. bush muhly. Growth begins in the winter to early spring, goes semi-dormant in May through June, most growth occurs during the summer rainy season, green most of the year..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	10	10	10	5	5	15	20	20	5	0	0

Figure 8. Plant community growth curve (percent production by month). AZ3091, 30.23 6-10" p.z. catclaw acacia. Growth begins in spring and continues through summer. Flowers from May through June. Seeds from July through September..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	0	5	15	10	15	30	15	10	0	0

Figure 9. Plant community growth curve (percent production by month). AZ3092, 30.28 10-13" p.z. white burrobrush. Growth begins in the spring and continues through the summer. Flowers and sets seed during the summer rainy season..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	10	20	10	10	20	20	10	0	0	0

State 2

Historic Native Plant Community plus Introduced Annuals

Community 2.1

Historic Native Plant Community plus Introduced Annuals

This plant community resembles the historic climax plant community, but includes introduced annuals.

Table 8. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Shrub/Vine	280	342	364
Grass/Grasslike	112	140	168
Forb	56	78	84
Total	448	560	616

Table 9. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	0-2%
Grass/grasslike foliar cover	0%
Forb foliar cover	0%
Non-vascular plants	0%
Biological crusts	0%
Litter	0%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%

Table 10. Canopy structure (% cover)

Height Above Ground (M)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.15	–	–	–	0-2%
>0.15 <= 0.3	–	–	1-3%	–
>0.3 <= 0.6	–	–	–	–
>0.6 <= 1.4	–	6-10%	–	–
>1.4 <= 4	–	–	–	–
>4 <= 12	–	–	–	–
>12 <= 24	–	–	–	–
>24 <= 37	–	–	–	–
>37	–	–	–	–

Figure 11. Plant community growth curve (percent production by month). AZ3011, 30.1 3-6" p.z. all sites. Growth begins in late winter, most growth occurs in the spring..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	7	30	32	13	7	3	5	2	1	0	0

Figure 12. Plant community growth curve (percent production by month). AZ3080, 30.27 3-6" p.z. red brome (exotic). Germinates in October with fall precipitation, most growth occurs in February through April..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	15	60	20	0	0	0	0	0	5	0	0

Figure 13. Plant community growth curve (percent production by month). AZ3081, 30.27 3-6" p.z. redstem filaree (exotic). Germinates in October with fall precipitation, most growth occurs in February through April..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	15	60	20	0	0	0	0	0	5	0	0

Additional community tables

Table 11. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1				28–56	
	big galleta	PLRI3	<i>Pleuraphis rigida</i>	28–56	–
2				28–56	
	bush muhly	MUPO2	<i>Muhlenbergia porteri</i>	28–56	–
3				0–6	
	sixweeks grama	BOBA2	<i>Bouteloua barbata</i>	0–6	–
4				0–6	
	sixweeks fescue	VUOC	<i>Vulpia octoflora</i>	0–6	–
5				0–11	
	Parish's threeawn	ARPUP5	<i>Aristida purpurea var. parishii</i>	0–11	–
6				11–45	
	Grass, annual	2GA	<i>Grass, annual</i>	11–45	–
Forb					

7				28–56	
	desert Indianwheat	PLOV	<i>Plantago ovata</i>	28–56	–
8				6–45	
	Forb, annual	2FA	<i>Forb, annual</i>	6–45	–
9				6–11	
	Forb, perennial	2FP	<i>Forb, perennial</i>	6–11	–
Shrub/Vine					
10				28–56	
	catclaw acacia	ACGR	<i>Acacia greggii</i>	28–56	–
11				140–196	
	burrobrush	HYSA	<i>Hymenoclea salsola</i>	140–196	–
12				6–28	
	smoketree	PSSP3	<i>Psoralea spinosa</i>	6–28	–
13				6–28	
	creosote bush	LATR2	<i>Larrea tridentata</i>	6–28	–
14				6–28	
	burrobush	AMDU2	<i>Ambrosia dumosa</i>	6–28	–
15				0–17	
	yellow paloverde	PAMI5	<i>Parkinsonia microphylla</i>	0–17	–
16				6–28	
	water jacket	LYAN	<i>Lycium andersonii</i>	6–28	–
17				6–28	
	brittlebush	ENFA	<i>Encelia farinosa</i>	6–28	–
18				0–11	
	sweetbush	BEJU	<i>Bebbia juncea</i>	0–11	–
19				0–17	
	Mexican bladdersage	SAME	<i>Salazaria mexicana</i>	0–17	–
20				6–28	
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	6–28	–

Table 12. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1				28–56	
	big galleta	PLRI3	<i>Pleuraphis rigida</i>	28–56	–
2				28–56	
	bush muhly	MUPO2	<i>Muhlenbergia porteri</i>	28–56	–
3				6–28	
	red brome	BRRU2	<i>Bromus rubens</i>	6–28	–
4				0–6	
	sixweeks grama	BOBA2	<i>Bouteloua barbata</i>	0–6	–
5				0–6	
	sixweeks fescue	VUOC	<i>Vulpia octoflora</i>	0–6	–
6				0–11	

	Parish's threeawn	ARPUP5	<i>Aristida purpurea</i> var. <i>parishii</i>	0–11	–
7				6–17	
	Grass, annual	2GA	<i>Grass, annual</i>	6–17	–
Forb					
8				6–17	
	redstem stork's bill	ERCI6	<i>Erodium cicutarium</i>	6–17	–
9				28–56	
	desert Indianwheat	PLOV	<i>Plantago ovata</i>	28–56	–
10				0–28	
	Forb, annual	2FA	<i>Forb, annual</i>	0–28	–
11				6–11	
	Forb, perennial	2FP	<i>Forb, perennial</i>	6–11	–
Shrub/Vine					
12				28–56	
	catclaw acacia	ACGR	<i>Acacia greggii</i>	28–56	–
13				140–196	
	burrobrush	HYSA	<i>Hymenoclea salsola</i>	140–196	–
14				6–28	
	smoketree	PSSP3	<i>Psoralea argophylla</i>	6–28	–
15				6–28	
	creosote bush	LATR2	<i>Larrea tridentata</i>	6–28	–
16				6–28	
	burrobush	AMDU2	<i>Ambrosia dumosa</i>	6–28	–
17				0–17	
	yellow paloverde	PAMI5	<i>Parkinsonia microphylla</i>	0–17	–
18				6–28	
	water jacket	LYAN	<i>Lycium andersonii</i>	6–28	–
19				6–28	
	brittlebush	ENFA	<i>Encelia farinosa</i>	6–28	–
20				0–11	
	sweetbush	BEJU	<i>Bebbia juncea</i>	0–11	–
21				0–17	
	Mexican bladdersage	SAME	<i>Salazaria mexicana</i>	0–17	–
22				6–28	
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	6–28	–

Type locality

Location 1: Mohave County, AZ	
Township/Range/Section	T29 N. R21 W. S26
General legal description	Black Canyon Quad. - Detrital Wash Area East of Temple Bar Road; Sec 26, T. 29 N., R. 21 W.; Mohave County, Arizona

Contributors

Harmon Hodgkinson
Harmon S. Hodgkinson
Larry D. Ellicott
Stephen Cassady

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

14. **Average percent litter cover (%) and depth (in):**

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

17. **Perennial plant reproductive capability:**
