

Ecological site R030XB148CA Sandy Plain

Accessed: 05/20/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 030X–Mojave Basin and Range

MLRA Description:

Major Land Resource Area (MLRA) 30, Mojave Desert, is found in southern California, southern Nevada, the extreme southwest corner of Utah and northwestern Arizona within the Basin and Range Province of the Intermontane Plateaus. The climate of the area is hot (primarily hyperthermic and thermic; however at higher elevations, generally above 5000 feet, mesic, cryic and frigid) and dry (aridic). Elevations range from below sea level to over 12,000 feet in the higher mountain areas found within the MLRA. Due to the extreme elevational range found within this MLRA, Land Resource Units (LRUs) were designated to group the MLRA into similar land units.

LRU Description:

This LRU (designated by 'XB') is found across the eastern half of California, much of the mid-elevations of Nevada, the southernmost portions of western Utah, and the mid-elevations of northwestern Arizona. Elevations range from 1800 to 5000 feet and precipitation ranges from 4 to 9 inches per year, but is generally between 5-6 inches. This LRU is characterized primarily by the summer precipitation it receives, ranging from 18 – 35% but averages 25%. Summer precipitation falls between July and September in the form of rain, and winter precipitation falls starting in November and ends between February and March, also mostly in the form of rain; however it does receive between 0 and 3 inches of snow, with an average of 1 inch. The soil temperature regime is thermic and the soil moisture

regime is typic-aridic. Vegetation includes creosote bush, burrobush, Nevada jointfir, ratany, Mojave yucca, Joshua tree, chollas, cactus, big galleta grass and several other warm season grasses. At the upper portions of the LRU, plant production and diversity are greater and blackbrush is a common dominant shrub.

Ecological Site Concept -

This ecological site occurs on stabilized sand sheets, dunes, sand sheets on fan remnants, and fan aprons on fan remnants at elevations of 2240 to 2850 feet, and slopes ranging from 0 to 8 percent. Dominant soils are very deep fine sands that formed from eolian deposits and exhibit very little soil development. Soils may have an alluvial influence, but eolian processes dominate.

Production reference value (RV) is relatively high at 1000 pounds per acre, and depending on annual precipitation, ranges from 595 to 1400 pounds per acre. Perennial grasses dominate this site, with big galleta (*Pleuraphis rigida*) dominant, and Indian rice grass (*Achnatherum hymenoides*) an important secondary species. Creosote bush (*Larrea tridentata*) is the dominant shrub. Annual forbs are abundant during years of average to above average precipitation. Stable, deep fine sands with little soil development favors perennial grasses and the deep-rooted creosote bush.

The data in the following sections is from major (15% of mapunit or greater) components only.

Classification relationships

This ecological site is found within the *Pleuraphis rigida* Herbaceous Alliance (Sawyer et al. 2009), and includes the *Pleuraphis rigida*/*Larrea tridentata* Association.

Associated sites

R030XB005NV	Arid Active Alluvial Fans This ecological site occurs on adjacent fan aprons. Creosote bush (<i>Larrea tridentata</i>) and burrobush (<i>Ambrosia dumosa</i>) are co-dominant.
R030XB137CA	Granitic Loam This ecological site occurs on adjacent sand sheets over fan aprons. Creosote bush (<i>Larrea tridentata</i>), burrobush (<i>Ambrosia dumosa</i>), and big galleta (<i>Pleuraphis rigida</i>) are co-dominant.
R030XB150CA	Sandhill 3-5" P.Z. This ecological site occurs on adjacent sand hills. Big galleta (<i>Pleuraphis rigida</i>) and creosote bush (<i>Larrea tridentata</i>) are dominant.

Similar sites

R030XB137CA	Granitic Loam This ecological site occurs on more stable landforms. It is less productive, and burrobush is a dominant shrub.
R030XB039NV	LIMY FAN 5-7 P.Z. This ecological site occurs on inset fans, and landforms receiving additional moisture.
R030XD014CA	Hyperthermic Sandy Plains This ecological site occurs on soils with a hyperthermic soil temperature regime. Indian rice grass is not present.
R030XB150CA	Sandhill 3-5" P.Z. This ecological site occurs on steeper slopes, and is less productive.

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Larrea tridentata</i>
Herbaceous	(1) <i>Pleuraphis rigida</i>

Physiographic features

This site occurs on sand sheets, dunes, sand sheets on fan remnants and fan aprons on fan remnants at elevations of 2240 to 2850 feet, and slopes ranging from 0 to 8 percent. Runoff class is very low to low.

Table 2. Representative physiographic features

Landforms	(1) Sand sheet (2) Dune (3) Fan apron
Flooding frequency	None
Ponding frequency	None
Elevation	683–869 m
Slope	0–8%
Aspect	Aspect is not a significant factor

Climatic features

The climate on this site is arid, characterized by warm, moist winters (30 to 60 degrees F) and hot, somewhat dry summers (70 to 100 degrees F). The average annual precipitation ranges from 2 to 7 inches with most falling as rain from November to March. Approximately 45% of the annual precipitation occurs from July to September as a result of summer convection storms. Mean annual air temperature is 63 to 69 degrees F. The average frost-free period is 240 to 340 days. Freeze-free period was not entered and defaults to zero.

Table 3. Representative climatic features

Frost-free period (average)	340 days
Freeze-free period (average)	0 days
Precipitation total (average)	178 mm

Influencing water features

Soil features

The dominant soils associated with this ecological site are very deep, and formed in alluvium derived from granitic sources. Surface textures are loamy fine sand and sand, with sand, loamy sand and sandy loam subsurface textures. Surface gravels (< 3 mm in diameter) range from 0 to 10 percent, typically with no larger fragments. Subsurface gravels by volume (for a depth of 0 to 59 inches) range from 0 to 12 percent, typically with no larger fragments present. Soils are well to somewhat excessively drained with moderately rapid to rapid permeability.

This ecological site is associated with major components (15 percent or greater) of the following soil series: Bluepoint (Mixed, thermic Typic Torripsamments); Pipesflat (Loamy, mixed, superactive, thermic Arenic Haplargids); and Burntshack (Loamy, mixed, superactive, thermic Arenic Haplargids). It is also correlated with minor components of Rositas (Mixed, hyperthermic Typic Torripsamments), although this correlation needs to be corrected since Rositas is a hyperthermic soil.

The Bluepoint soils formed from eolian sand derived from mixed parent material, and consist of stratified layers of fine sand. The Pipesflat and Burntshack soils consist of soils that formed in eolian sand over alluvium derived from granite. These soils have an argillic horizon that occurs at depths of 20 to 40 inches. Pipesflat soils have eolian sands above the argillic horizon, while Burntshack soils have a calcic horizon above the argillic. The Rositas are hyperthermic and not typical for this ecological site.

This ecological site is correlated with the following soil survey areas, map units and soil components (Soil survey area; Mapunit symbol; Mapunit name; Component; phase; percent):

CA698 Mojave Desert Area, West Central Part

CA698;4710;Bluepoint loamy fine sand, 2 to 8 percent slopes;Bluepoint;loamy fine sand;80

CA698;100tp;Cajon-Pipeflat association, 2 to 8 percent slopes;Pipeflat;;25

CA698;4002;Daisy-Hypoint-Silvermine complex, 2 to 8 percent slopes;Bluepoint;;1

CA698;103tp;Cajon-Calcio-Edalphy complex, 2 to 4 percent slopes;Pipeflat;;5

CA698;4730;Calcic Haplosalids-Sodic Haplosalids-Typic Haplosalids complex, 0 to 2 percent slopes;Rositas;;3

CA794 Joshua Tree National Park

CA794;4041;Silvermine-Helendale-Burntshack association, 1 to 15 percent slopes;Burntshack;very rarely flooded;20

CA699 Twenynine Palms Marine Corps Air Ground Combat Center

CA699; 205;;Pipeflat; 30

CA699; 100;;Pipeflat, 25; Bluepoint; 10

CA699; 103;;Pipeflat; 5

Table 4. Representative soil features

Parent material	(1) Eolian sands–granite
Surface texture	(1) Loamy fine sand (2) Sand
Family particle size	(1) Sandy
Drainage class	Well drained to somewhat excessively drained
Permeability class	Moderately rapid to rapid
Soil depth	152 cm
Surface fragment cover <=3"	0–10%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	5.84–9.65 cm
Calcium carbonate equivalent (0-101.6cm)	0–5%
Electrical conductivity (0-101.6cm)	0–8 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0–5
Soil reaction (1:1 water) (0-101.6cm)	7.2–9
Subsurface fragment volume <=3" (Depth not specified)	0–12%
Subsurface fragment volume >3" (Depth not specified)	0%

Ecological dynamics

Abiotic Factors

This ecological site occurs on stabilized sandsheets on soils with a thermic soil temperature regime. Sandsheets are extensive, low relief accumulations of eolian sand deposits (Laity 2008). Stable, or dormant sandsheets are those where perennial vegetation cover is well-developed, and current rates of sand movement and deposition are low or absent, but may become active as a result of minor climate change or disturbance (Lancaster 1994). The stability of these landforms means that factors such as burial or abrasion by blowing sand does not restrict

vegetation to psammophiles (plants restricted to active eolian environments).

The plant community is strongly dominated by perennial grasses. Big galleta is the dominant species; big galleta is a highly drought-tolerant C4 grass that occurs on a range of soil types, but is dominant only on sandy soils where soil moisture is most readily available (McAuliffe 1994, Austin et al. 2004). Big galleta colonizes and stabilizes semi-stabilized eolian habitats with rhizomatous growth (Matthews 2000), and dominance by big galleta on these habitats is an indicator of eolian stability. Big galleta exhibits rapid growth and high productivity in response to temporal high moisture availability in these deep sands (Austin et al. 2004). Indian rice grass is a cool season perennial bunchgrass widely distributed throughout the western United States, but also reaches highest abundance on sandy soils (Tirmenstein 1999, Baldwin et al. 2002). It occurs in the thermic soil temperature regime of this ecological site, and disappears when the soil temperature regime transitions to hyperthermic. In arid regions, sand textured soils have greater water availability because water quickly infiltrates through sand to depths where it is not lost to evaporation, and because sandy surfaces form a physical crust that further reduces evaporation (Noy-Meir 1973, Hamerlynk et al. 2002). Thus, in desert regions, where the availability of soil water is the critical resource shaping plant communities in arid environments, productivity is highest on sandy soils (Noy-Meir 1973, McAuliffe 1994, Martre et al. 2002, Hamerlynk and McAuliffe 2002, Austin et al. 2004).

Creosote bush is a long-lived, deep-rooted evergreen shrub dominant across vast areas of the North American warm deserts. Creosote bush maintains its evergreen status by using water held in deep soil layers, and once established in this ecological site, individuals are large and productive. Creosote remains a secondary species in this site however, because of soil moisture restrictions and seedling sand abrasion during the establishment phase. Creosote bush establishes in response to warm season moisture; given limited warm season rain in this ecological site, the rapid infiltration of water, rapidly drying soil surfaces during the warm season, and increased erosion and abrasion during the summer, opportunities for successful establishment of creosote seedlings are rare.

Disturbance dynamics

Drought, invasion by nonnative species, and wind erosion are the primary disturbance affecting this ecological site.

Drought is an important shaping force in desert plant communities (Webb et al. 2003, Bowers 2005, Hereford et al. 2006, Miriti et al. 2007, Hamerlynk and McAuliffe 2008). The effects of drought may be particularly severe in deep sandy soils with little horizon development. High availability of soil moisture during normal to high precipitation conditions can lead to high growth rates and large individuals whose size cannot be sustained when water is no longer available (Hamerlynk and McAuliffe 2008). Short-lived shrubs and perennial grasses demonstrate the highest rates of drought-induced mortality (Webb et al. 2003, Bowers 2005, Hereford et al. 2006, Miriti et al. 2007), and annual species remain dormant in the soil seedbank (Beatley, 1974, 1976). Long-lived species are more likely to exhibit branch-pruning with limited recruitment during drought (Hereford et al. 2006, Miriti et al. 2007).

Non-native annual species such as red brome (*Bromus rubens*), Mediterranean grass (*Schismus barbatus*), redstem stork's bill (*Erodium cicutarium*) and Asian mustard (*Brassica tournefortii*) have become naturalized throughout the Mojave Desert over the past century (Rickard and Beatley 1965, D'Antonio and Vitousek 1992, Brooks 1999, Reid et al. 2006, Norton et al. 2007). Asian mustard and prickly Russian thistle (*Salsola tragus*) are threats in eolian habitats, with prickly Russian thistle abundant on disturbed areas or active sand, and Asian mustard most abundant on stabilized sand (Barrows et al. 2009). Like native annuals, nonnative annual cover and production is directly related to winter precipitation (Beatley 1969, Brooks and Berry 2006, Barrows et al. 2009). In this ecological site, Asian mustard and prickly Russian thistle invasion may be severe.

Wind erosion and deposition is the driver of eolian dynamics. Wind strength, precipitation, vegetation cover and disturbance influence the degree to which sand depositional surfaces are active or stable (Cooke et al. 1993, Lancaster 1994, 1997, Musick 1999). Drought may cause a stabilized sand surface to become active, due to losses in vegetation cover, and the increased erodibility of dry soils (Cooke et al. 1993, Lancaster 1994, Breed and Reheis 1999, Musick 1999). Similarly, other disturbances that cause a decline in vegetation cover, such as off-road vehicle use, grazing, and fire can reactivate a stable sand surface.

State and transition model




Figure 3. R030XB148CA

**State 1
Historic State**

State 1 represents the historic range of variability for this ecological site. This state no longer exists due to the ubiquitous naturalization of non-native species in the Mojave Desert. Periodic drought and rare fire were the natural disturbances influencing this ecological site. Fire would have been a very rare occurrence due to the lack of a continuous fine fuel layer between shrubs. Data for this State does not exist, but dynamics and composition would have been similar to State 2, except with only native species present. See State 2 narrative for more detailed

information.

State 2 Reference State

State 2 represents the current range of variability for this site. Non-native annuals, including red brome, Mediterranean grass, red-stem stork's bill, and Asian mustard are naturalized in this plant community. Their abundance varies with precipitation, but they are at least sparsely present (as current year's growth or present in the soil seedbank).

Community 2.1 Reference Plant Community



Figure 4. Community Phase 2.1

The reference plant community is characterized by an open two-tiered canopy less than 2 meters tall with creosote bush in the upper tier over a dense stand of big galleta and Indian ricegrass. Sand dropseed (*Sporobolus cryptandrus*) may also be present at low levels. Secondary shrubs may include burrobush (*Ambrosia dumosa*), white ratany (*Krameria grayi*), California ephedra (*Ephedra californica*), Wiggins' cholla (*Cylindropuntia echinocarpa*), Plummer's baccharis (*Baccharis plummerae*), and rayless goldenhead (*Acamptopappus sphaerocephalus*). The subshrubs California croton (*Croton californicus*) and desert globemallow (*Sphaeralcea ambigua*) are typically present. A spectacular display of annual forbs occurs during years of above-average precipitation. Common species typically include Esteve's pincushion (*Chaenactis steviodes*), smooth desertdandelion (*Malacothrix glabrata*), bristly fiddleneck (*Amsinckia tessellata*), and birdcage evening primrose (*Oenothera deltoides*). Non native species that may be present include Asian mustard, prickly Russian thistle, redstem stork's bill, and Mediterranean grass.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	549	785	1098
Shrub/Vine	118	168	235
Forb	—	168	235
Total	667	1121	1568

Table 6. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	2-3%
Grass/grasslike foliar cover	7-14%
Forb foliar cover	2-3%
Non-vascular plants	0%

Biological crusts	0%
Litter	0%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%

Figure 6. Plant community growth curve (percent production by month). CA3015, Creosote bush XB. Growth starts in early spring with flowering and seed set occurring by July. Dormancy occurs during the hot summer months. With sufficient summer/fall precipitation, some vegetation may break dormancy and produce a flush of growth..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	5	20	30	20	10	0	10	5	0	0	0

Figure 7. Plant community growth curve (percent production by month). CA3022, Indian ricegrass. Growth begins in late winter, flowering and fruiting finished by the hot summer months. Early fall rains can trigger a flush of new growth..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	5	15	30	35	5	0	0	5	5	0	0

Figure 8. Plant community growth curve (percent production by month). CA3024, Big galleta. Some green up in spring; dormant May and June; most growth occurs after summer rains..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	5	20	10	0	0	15	40	10	0	0	0

Community 2.2 Drought Response

This community phase is characterized by an overall decline in cover due to branch-pruning and lack of recruitment of longer-lived species, mortality of shorter-lived perennials, and lack of emergence of annual forbs and grasses. Big galleta and Indian ricegrass are likely to decline due to drought-induced mortality, while creosote bush remains stable. Big galleta may suffer very high rates of drought-induced mortality (Webb et al. 2003; Hereford et al. 2006); however, big galleta can respond very quickly to brief, intermittent rain during rare summer monsoonal events, which can buffer big galleta populations in the absence of more predictable winter rains. Creosote bush is an evergreen species capable of utilizing moisture at any time of the year. This ability buffers populations from the effects of drought that occur as the absence of the winter rains (the primary source of moisture for this ecological site). Further, creosote bush germinates in response to moisture during the warm season, so may still recruit if warm season rains occur during winter drought (Hereford et al. 2006). Creosote bush exhibits branch-pruning during severe drought, but mortality during drought in the Mojave Desert is very low (Webb et al. 2003, Griffiths et al. 2006). Nevertheless, during severe drought, creosote bush mortality may occur. This is an at-risk community. Reduced cover in this eolian landscape increases the risk of erosion, which can trigger a transition to State 3.

Community 2.3 Fire regeneration community

This community phase is characterized by increased dominance by big galleta and Indian rice grass, severe declines in creosote bush, and an increase in shrub diversity. Fire damage to big galleta varies depending on whether plants are dormant when burned; if plants are dry, damage may be severe because the live center may be burned out (Matthews 2000). However big galleta often increases after fire (Minnich 2003). Indian rice grass is

highly fire tolerant, and also increases after fire (Tirmenstein 1999). Creosote bush is generally killed by fire, and is slow to re-colonize burned areas due to specific recruitment requirements (Brown and Minnich 1986, Brooks et al. 2007, Steers and Allen 2011). The timing and severity of fire, as well as post-fire climate conditions determines trajectories of recovery (Brown and Minnich 1986, Steers and Allen 2011).

Pathway 2.1a **Community 2.1 to 2.2**

This pathway occurs with prolonged or severe drought.

Pathway 2.1b **Community 2.1 to 2.3**

This pathway occurs with moderate to severe fire.

Pathway 2.2a **Community 2.2 to 2.1**

This pathway occurs with a return to average or above average precipitation.

Pathway 2.2b **Community 2.2 to 2.3**

This pathway occurs with moderate to severe fire.

Pathway 2.3a **Community 2.3 to 2.1**

This pathway occurs with time without fire, adequate precipitation, and no other significant disturbances (e.g. grazing).

State 3 **Eroded State**

This State is characterized by the loss of sandsheet stability, with increased rates of wind erosion leading to deflation. This state has been significantly altered from the natural range of variability found in States 1 and 2. Increased wind erosion decreases the suitability of this ecological site for vegetation, killing established or recruiting individuals by abrasion and burial (Okin et al. 2001). Ongoing disturbance could result in complete loss of vegetation cover. Sand deflation could result in the accumulation of surface rock fragments, dramatically altering the soil and hydrological characteristics of this ecological site, and decreasing site suitability for annual species and big galleta. We do not have data for this State, and further research is necessary to describe the community phases and successional pathways that may exist within the state.

Transition 1 **State 1 to 2**

This transition occurred with the naturalization of non-native species in this ecological site. Non-native species were introduced with settlement of the Mojave Desert region in the 1860s. Post-settlement cattle and sheep grazing, as well as dryland farming, helped to spread and facilitate their establishment (Brooks and Pyke 2000, Brooks et al. 2007).

Transition 2 **State 2 to 3**

This transition occurs with a loss of vegetation cover, in combination with drought and/or extreme wind conditions and/or anthropogenic disturbance such as grazing or off-road vehicle use that increases wind erosion beyond the threshold that will sustain the reference plant community. It is difficult to pinpoint the precise combination of these

factors that will trigger this conversion (Cooke et al. 1993).

Additional community tables

Table 7. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1	Perennial Grasses			549–1098	
	big galleta	PLRI3	<i>Pleuraphis rigida</i>	560–729	–
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	112–168	–
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	22–56	–
4	Native Annual Grasses			0–56	
	sixweeks grama	BOBA2	<i>Bouteloua barbata</i>	0–56	–
5	Non-native annual grasses			0–123	
	common Mediterranean grass	SCBA	<i>Schismus barbatus</i>	0–123	–
Shrub/Vine					
2	Native shrubs			118–235	
	Plummer's baccharis	BAPL	<i>Baccharis plummerae</i>	0–224	–
	creosote bush	LATR2	<i>Larrea tridentata</i>	56–168	–
	rayless goldenhead	ACSP	<i>Acamptopappus sphaerocephalus</i>	22–112	–
	burrobush	AMDU2	<i>Ambrosia dumosa</i>	22–112	–
	Wiggins' cholla	CYEC3	<i>Cylindropuntia echinocarpa</i>	22–112	–
	California jointfir	EPCA2	<i>Ephedra californica</i>	22–112	–
	white ratany	KRGR	<i>Krameria grayi</i>	22–112	–
Forb					
3	Native Forbs			0–235	
	cryptantha	CRYPT	<i>Cryptantha</i>	0–224	–
	milkvetch	ASTRA	<i>Astragalus</i>	0–224	–
	browneyes	CACLC3	<i>Camissonia claviformis ssp. claviformis</i>	0–224	–
	Esteve's pincushion	CHST	<i>Chaenactis stevioides</i>	0–224	–
	birdcage evening primrose	OEDE2	<i>Oenothera deltoides</i>	0–224	–
	bristly fiddleneck	AMTE3	<i>Amsinckia tessellata</i>	0–191	–
	California croton	CRCA5	<i>Croton californicus</i>	22–90	–
	smooth desertydandelion	MAGL3	<i>Malacothrix glabrata</i>	0–67	–
	desert globemallow	SPAM2	<i>Sphaeralcea ambigua</i>	22–56	–
6	Non-native annual forbs			0–224	
	redstem stork's bill	ERCI6	<i>Erodium cicutarium</i>	0–112	–
	prickly Russian thistle	SATR12	<i>Salsola tragus</i>	0–112	–

Animal community

Small mammals occurring on this site include round-tailed ground squirrels, little pocket mice, and Merriam's and desert kangaroo rats. Black-tailed jackrabbits and coyotes are also common.

Reptiles occurring on this site include several species of lizards including Mojave fringe-toed lizards, long-tailed brush lizards, side-blotched lizards, and western whiptails. Common snakes include western shovel-nosed snakes, glossy snakes and sidewinders.

Birds common to this site include horned larks, common ravens, loggerhead shrikes, LeConte's thrashers and several species of sparrows. Raptors observed on this site include northern harriers, sharp-shinned hawks and American kestrels.

LIVESTOCK GRAZING:

Big galleta and Indian ricegrass are highly palatable to cattle and horses. Burrobush is fair browse for cattle and horses, and fair to good browse for goats. Sheep also use this shrub, feeding primarily on new growth and seeds. Creosote bush is unpalatable to livestock. Domestic sheep use this shrub for shade. During favorable years, annual forbs and grasses provide abundant forage.

Hydrological functions

Recreational uses

This site is highly valued for open space and those interested in desert ecology. Flowering wildflowers and shrubs may also attract visitors during the spring.

Other information

Military Operations - Management for this site would be to protect it from excessive disturbance and maintain existing plant cover. Land clearing or other disturbances that destroy the vegetation and soil structure can result in soil compaction reduced infiltration rates, accelerated erosion, severe soil blowing and barren areas.

Inventory data references

Sampling technique

1 NV-ECS-1

2 SCS-Range 417

3 Other

CA794:

MVAL-07

Type locality

Location 1: San Bernardino County, CA	
Township/Range/Section	T3N R6E S13
UTM zone	N
UTM northing	3800842
UTM easting	0564131
General legal description	SE1/4 Sec. 13 T3N R6E Approximately 15 miles north of Joshua Tree, CA Goat Mountain Quadrangle UTM 11S 0564131e 3800842n (Datum=NAS-C) San Bernardino Co., CA

Other references

Austin, A. T., L. Yahdjian, J. M. Stark, J. Belnap, A. Porporato, U. Norton, D. A. Ravetta, and S. M. Scheaeffer. 2004. Water pulses and biogeochemical cycles in arid and semiarid ecosystems. *Oecologia* 141:221-235.

- Baldwin, B. G., S. Boyd, B. J. Ertter, R. W. Patterson, T. J. Rosatti, and D. H. Wilken. 2002. *The Jepson Desert Manual*. University of California Press, Berkeley and Los Angeles, California.
- Barrows, C. W., E. B. Allen, M. L. Brooks, and M. F. Allen. 2009. Effects of an invasive plant on a desert sand dune landscape. *Biological Invasions* 11:673-686.
- Beatley, J. C. 1969. Dependence of desert rodents on winter annuals and precipitation. *Ecology* 50:721-724.
- Beatley, J. C. 1974. Effects of rainfall and temperature on the distribution and behavior of *Larrea tridentata* (Creosote-bush) in the Mojave Desert of Nevada. *Ecology* 55:245-261.
- Beatley, J. C. 1976. Rainfall and fluctuating plant populations in relation to distributions and numbers of desert rodents in southern Nevada. *Oecologia* 24:21-42.
- Bowers, J. E. 2005. Effects of drought on shrub survival and longevity in the northern Sonoran Desert. *Journal of the Torrey Botanical Society* 132:421-431.
- Breed, C. S. and M. C. Reheis. 1999. Desert winds: monitoring wind-related surface processes in Arizona, New Mexico, and California. Denver, CO.
- Brooks, M. L. 1999. Habitat invasibility and dominance by alien annual plants in the western Mojave Desert. *Biological Invasions* 1:325-337.
- Brooks, M. L. and K. H. Berry. 2006. Dominance and environmental correlates of alien annual plants in the Mojave Desert, USA. *Journal of Arid Environments* 67:100-124.
- Brooks, M. L., T. C. Esque, and T. Duck. 2007. Creosotebush, blackbrush, and interior chaparral shrublands. RMRS-GTR-202.
- Brooks, M. L. and D. A. Pyke. 2000. Invasive plants and fire in the deserts of North America. Pages 1-14 in *Fire conference 2000: the first national congress on fire ecology, prevention, and management*. Tall Timbers Research Station, Tallahassee, FL.
- Brown, D. E. and R. A. Minnich. 1986. Fire and Changes in Creosote Bush Scrub of the Western Sonoran Desert, California. *American Midland Naturalist* 116:411-422.
- Cooke, R. U., A. Warren, and A. S. Goudie. 1993. *Desert Geomorphology*. UCL Press, London.
- D'Antonio, C. M. and P. M. Vitousek. 1992. Biological invasions by exotic grasses, the grass/fire cycle, and global change. *Annual Review of Ecology and Systematics* 23:63-87.
- Griffeths, P. G., R. Hereford, and R. H. Webb. 2006. Sediment yield and runoff frequency of small drainage basins in the Mojave Desert, U.S.A. *Geomorphology* 74:232-244.
- Hamerlynk, E. P. and J. R. McAuliffe. 2008. Soil-dependent canopy die-back and plant mortality in two Mojave Desert shrubs. *Journal of Arid Environments* 72:1793-1802.
- Hamerlynk, E. P., J. R. McAuliffe, E. V. McDonald, and S. D. Smith. 2002. Ecological responses of two Mojave desert shrubs to soil horizon development and soil water dynamics. *Ecology* 83:768-779.
- Hereford, R., R. H. Webb, and C. I. Longpre. 2006. Precipitation history and ecosystem response to multidecadal precipitation variability in the Mojave Desert region, 1893-2001. *Journal of Arid Environments* 67:13-34.
- Laity, J. 2008. *Deserts and desert environments*. John Wiley & Sons.
- Lancaster, N. 1994. Controls on aeolian activity: some new perspectives from the Kelso Dunes, Mojave Desert, California. *Journal of Arid Environments* 27:113-125.
- Lancaster, N. 1997. Response of eolian geomorphic systems to minor climate change: examples from the southern

California deserts. *Geomorphology* 19:333-347.

Matthews, R. F. 2000. *Pleuraphis rigida*. In: Fire Effects Information System. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer).

McAuliffe, J. R. 1994. Landscape evolution, soil formation, and ecological patterns and processes in Sonoran Desert bajadas. *Ecological Monographs* 64:112-148.

Minnich, R. A. 2003. Fire and dynamics of temperature desert woodlands in Joshua Tree National Park. Contract, Joshua Tree National Park.

Miriti, M. N., S. Rodriguez-Buritica, S. J. Wright, and H. F. Howe. 2007. Episodic death across species of desert shrubs. *Ecology* 88:32-36.

Musick, H. B. 1999. Field monitoring of Vegetation Characteristics related to surface changes in the Yuma Desert, Arizona, and at the Jornada Experimental Range in the Chihuahuan Desert, New Mexico. Pages 78-91 in C. S. Breed and M. C. Reheis, editors. *Desert winds: Monitoring wind-related surface processes in Arizona, New Mexico, and California*. U.S. Geological Survey, Denver CO.

Norton, J. B., T. A. Monaco, and U. Norton. 2007. Mediterranean annual grasses in western North America: kids in a candy store. *Plant Soil* 298:1-5.

Noy-Meir, I. 1973. Desert ecosystems: environment and producers. *Annual Review of Ecology and Systematics* 4:25-51.

Okin, G. S., B. Murray, and W. H. Schlesinger. 2001. Degradation of sandy arid shrubland environments: observations, process modelling, and management implications. *Journal of Arid Environments* 47:123-144.

Reid, C. R., S. Goodrich, and J. E. Bowns. 2006. Cheatgrass and red brome: history and biology of two invaders. Pages 27-32 in *Shrublands under fire: disturbance and recovery in a changing world*. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Cedar City, Utah.

Rickard, W. H. and J. C. Beatley. 1965. Canopy-coverage of the desert shrub vegetation mosaic of the Nevada test site. *Ecology* 46:524-529.

Sawyer, J. O., T. Keeler-Woolf, and J. M. Evans. 2009. *A manual of California vegetation*. 2nd edition. California Native Plant Society, Sacramento, California.

Steers, R. J. and E. B. Allen. 2011. Fire effects on perennial vegetation in the western Colorado Desert, USA. *Fire Ecology* 7:59-74.

Tirmenstein, D. 1999. *Achnatherum hymenoides*. In: Fire Effects Information System. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer).

Webb, R. H., M. B. Muroy, T. C. Esque, D. E. Boyer, L. A. DeFalco, D. F. Haines, D. Oldershaw, S. J. Scoles, K. A. Thomas, J. B. Blainey, and P. A. Medica. 2003. Perennial vegetation data from permanent plots on the Nevada Test Site, Nye County, Nevada. U.S. Geological Society, Tucson, AZ.

Contributors

Alice Lee Miller

P. Novak-Echenique

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem

condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

14. **Average percent litter cover (%) and depth (in):**

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

17. **Perennial plant reproductive capability:**
