

Ecological site R030XB203AZ Basalt Slopes 6-9" p.z.

Accessed: 05/18/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 030X–Mojave Basin and Range

AZ CRA 30.2 - Middle Mohave Desert

Elevations range from 1500 to 3200 feet and precipitation averages 6 to 9 inches per year. Vegetation includes creosotebush, white bursage, yucca, prickly pear and cholla species, Mormon tea, flattop buckwheat, ratany, winterfat, bush muhly, threeawns, and big galleta. The soil temperature regime is thermic and the soil moisture regime is typic aridic. This unit occurs within the Basin and Range Province and is characterized by broad basins, valleys, and old lakebeds. Widely spaced mountains trending north to south occur throughout the area. Isolated, short mountain ranges are separated by an aggraded desert plain. The mountains are fault blocks that have been tilted up. Long alluvial fans coalesce with dry lakebeds between some of the ranges.

Associated sites

R030XB210AZ	Limestone Hills 6-9" p.z.
R030XB214AZ	Limy Upland 6-9" p.z.
R030XB218AZ	Sandy Wash 6-9" p.z.
R030XB222AZ	Gypsum Hills 6-9" p.z. Alkaline

Similar sites

R030XB202AZ Basalt Hills 6-9" p.z.

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) Ambrosia dumosa (2) Larrea tridentata
Herbaceous	(1) Pleuraphis rigida (2) Muhlenbergia porteri

Physiographic features

This ecological site occurs on escarpments below basalt flows. It is often in association with basalt talus slopes and exposed mudstone. Soil depth ranges from shallow to very deep. Slopes range from 10 to 40 percent.

Landforms	(1) Escarpment
Flooding frequency	None
Ponding frequency	None
Elevation	488–1,372 m
Slope	10–40%
Aspect	Aspect is not a significant factor

Table 2. Representative physiographic features

Climatic features

The climate is arid and warm. Annual precipitation ranges from 6 to 9 inches. About 65 percent of the rainfall comes from October through May as gentle rain from Pacific storms which may last for a couple of days. The rest of the rainfall comes during the summer monsoon season from July through September as spotty, brief, intense thunderstorms. Snow rarely falls, and only remains on the ground a few hours at most. Annual air temperature ranges from 59 to 70 degrees F. The average frost-free period ranges from 156 to 259 days.

Table 3. Representative climatic features

Frost-free period (average)	259 days
Freeze-free period (average)	290 days
Precipitation total (average)	229 mm

Influencing water features

Soil features

The soil of this ecological site is shallow to very deep. The surface texture ranges from very cobbly fine sandy loam to loam. Subsoil textures range from very cobbly loam to loamy sand. Basalt gravels, stone and cobbles cover much of the soil surface.

Soil map units correlated to this ecological site include 623056, Nikey family and Ruesh family, Shivwits Area, Arizona, Part of Mohave County SSA.

Surface texture	(1) Very cobbly fine sandy loam(2) Sandy loam(3) Loam
Family particle size	(1) Loamy
Drainage class	Well drained
Permeability class	Moderate to moderately rapid
Soil depth	51–152 cm
Surface fragment cover <=3"	10–25%
Surface fragment cover >3"	5–30%
Calcium carbonate equivalent (0-101.6cm)	10–40%
Electrical conductivity (0-101.6cm)	2–4 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0
Soil reaction (1:1 water) (0-101.6cm)	7.4–8.4
Subsurface fragment volume <=3" (Depth not specified)	5–25%
Subsurface fragment volume >3" (Depth not specified)	0–10%

Ecological dynamics

The historic climax plant community (HCPC) for a site in North America is the plant community that existed at the time of European immigration and settlement. It is the plant community that was best adapted to the unique combination of environmental factors associated with the site at that time. The HCPC was in dynamic equilibrium with its environment and was able to avoid displacement by the suite of disturbances and disturbance patterns (magnitude and frequency) that naturally occurred within the area occupied by the site. Natural disturbances, such as drought, fire, grazing of native fauna, and insects, were inherent in the development and maintenance of the plant community. The effects of these disturbances are part of the range of characteristics of the site that contribute to the dynamic equilibrium. Fluctuations in the plant community's structure and function caused by the effects of these natural disturbances establish the boundaries of dynamic equilibrium. They are accounted for as part of the range of characteristics for the ecological site. The HCPC is not a precise assemblage of species for which the proportions are the same from place to place or from year to year. Variability is apparent in productivity and occurrence of individual species.

The HCPC for this ecological site has been estimated by sampling relict or relatively undisturbed sites and/or reviewing historic records.

A plant community that is subjected to abnormal disturbances and physical site deterioration or that is protected from natural influences, such as fire and grazing, for long periods seldom typifies the HCPC. Any physical site deterioration caused by the abnormal disturbance may result in the crossing of a threshold or irreversible boundary to another state, or equilibrium, for the ecological site. There may be multiple thresholds and states possible for an ecological site, determined by the type and or severity of abnormal disturbance. The known states and transition pathways for this ecological site are described in the accompanying state and transition model.

The "Plant Community Plant Species Composition" table provides a list of species and each species or group of species' annual production in pounds per acre (air-dry weight) expected in a normal rainfall year. Low and high production yields represent the modal range of variability for that species or group of species across the extent of the ecological site.

The "Annual Production by Plant Type" table provides the median air-dry production and the fluctuations to be expected during favorable, normal, and unfavorable years.

The present plant community on an ecological site can be compared to the various common vegetation states that can exist on the site. The degree of similarity is expressed through a similarity index. To determine the similarity index, compare the production of each species to that shown in the plant community description. For each species, count no more than the maximum amount shown for the species, and for each group, count no more than the maximum shown for the group. Divide the resulting total by the total representative value shown in the "Annual Production by Plant Type" table for the reference plant community. Variations in production due to above or below normal rainfall, incomplete growing season or utilization must be corrected before comparing it to the site description. The "Worksheet for Determining Similarity Index" is useful in making these corrections. The accompanying growth curve can be used as a guide for estimating percent of growth completed.

State and transition model



State 1 Historic Climax Plant Community

Community 1.1 Historic Climax Plant Community

The dominant aspect of this site is a perennial grass - desert shrubland mix. The major grasses are bush muhly and big galleta. The major shrubs are white bursage and creosotebush. Grass and grasslike plants make up 20 to 35 percent of the plant community, forbs 5 to 10 percent, shrubs 55 to 75 percent and trees 0 to 1 percent.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Shrub/Vine	77	183	243
Grass/Grasslike	28	77	114
Forb	7	19	33
Tree	-	1	2
Total	112	280	392

Table 6. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	1-3%
Grass/grasslike foliar cover	0-2%
Forb foliar cover	0-1%
Non-vascular plants	0%
Biological crusts	0%
Litter	0%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%

Table 7. Canopy structure (% cover)

Height Above Ground (M)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.15	_	_	_	0-1%
>0.15 <= 0.3	-	-	1-3%	-
>0.3 <= 0.6	_	_	_	_
>0.6 <= 1.4	_	6-10%	_	_
>1.4 <= 4	_	_	_	_
>4 <= 12	_	_	_	_
>12 <= 24	_	_	_	_
>24 <= 37	_	_	_	_
>37	_	-	_	_

Figure 5. Plant community growth curve (percent production by month). AZ3022, 30.2 6-9" p.z. upland sites. Growth begins in the late winter, most growth occurs in the spring..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	2	19	33	18	7	7	11	3	0	0	0

Figure 6. Plant community growth curve (percent production by month). AZ3070, 30.23 6-9" p.z. big galleta. Growth begins in the spring, most growth occurs during the summer rainy season..

Jan	Feb	Mar	Apr	Мау	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	10	15	5	0	10	45	15	0	0	0

Figure 7. Plant community growth curve (percent production by month). AZ3074, 30.23, 6-9 p.z., Nevada Mormon Tea. Growth primarily in spring and early summer months..

Jan	Feb	Mar	Apr	Мау	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	15	25	20	15	15	10	0	0	0	0

Figure 8. Plant community growth curve (percent production by month). AZ3075, 30.23 6-9" p.z. white ratany. Most growth occurs in the spring, some growth occurs in the summer. Flowers in the spring..

Jan	Feb	Mar	Apr	Мау	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	5	20	35	5	10	15	10	0	0	0

Figure 9. Plant community growth curve (percent production by month). AZ3077, 30.23 6-9" p.z. bush muhly. Growth begins in the spring, most growth occurs during the summer rainy season, green most of the year.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	5	5	5	5	15	40	20	5	0	0

Additional community tables

 Table 8. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)		
Grass	Grass/Grasslike						
1				28–43			
	big galleta	PLRI3	Pleuraphis rigida	28–43	_		
2				13–20			
	bush muhly	MUPO2	Muhlenbergia porteri	13–20	_		
3				2–9			
	sand dropseed	SPCR	Sporobolus cryptandrus	2–9	_		
	mesa dropseed	SPFL2	Sporobolus flexuosus	0–9	_		
4				2–9			
	desert needlegrass	ACSP12	Achnatherum speciosum	2–9	_		
5				2–6			
	low woollygrass	DAPU7	Dasyochloa pulchella	2–6	_		
6		•	•	0–13			
	Grass, perennial	2GP	Grass, perennial	0–13	_		
	Indian ricegrass	ACHY	Achnatherum hymenoides	0–13	_		
	threeawn	ARIST	Aristida	0–13	_		
	slim tridens	TRMU	Tridens muticus	0–13	_		
7				0–13			
	Grass, annual	2GA	Grass, annual	0–13	-		
	sixweeks grama	BOBA2	Bouteloua barbata	0–13	-		
Forb							
8				2–9			
	desert trumpet	ERIN4	Eriogonum inflatum	2–9	-		
9				2–6			
	desert globemallow	SPAM2	Sphaeralcea ambigua	2–6	-		

I					
10				2–9	
	trailing windmills	ALIN	Allionia incarnata	2–9	_
11		-	1	2–9	
	Forb, perennial	2FP	Forb, perennial	2–9	-
12		-		2–9	
	Forb, annual	2FA	Forb, annual	0–9	_
	buckwheat	ERIOG	Eriogonum	0–9	_
	spurge	EUPHO	Euphorbia	0–9	_
	woolly tidestromia	TILA2	Tidestromia lanuginosa	0–9	_
Shrub	/Vine	-	-		
13				43–84	
	burrobush	AMDU2	Ambrosia dumosa	43–84	-
14				28–43	
	creosote bush	LATR2	Larrea tridentata	28–43	-
15		-	-	9–25	
	Nevada jointfir	EPNE	Ephedra nevadensis	9–25	_
16		-	•	2–13	
	white ratany	KRGR	Krameria grayi	2–13	_
17		-		0–13	
	water jacket	LYAN	Lycium andersonii	0–13	-
18				0–13	
	Fremont's dalea	PSFR	Psorothamnus fremontii	0–13	-
19				0–11	
	rayless goldenhead	ACSP	Acamptopappus sphaerocephalus	0–11	-
20		-	-	28–56	
	Shrub (>.5m)	2SHRUB	Shrub (>.5m)	0–28	_
	Wright's beebrush	ALWR	Aloysia wrightii	0–28	_
	fourwing saltbush	ATCA2	Atriplex canescens	0–28	_
	mormon tea	EPVI	Ephedra viridis	0–28	-
	Eastern Mojave buckwheat	ERFAP	Eriogonum fasciculatum var. polifolium	0–28	_
	burrobrush	HYSA	Hymenoclea salsola	0–28	_
	winterfat	KRLA2	Krascheninnikovia lanata	0–28	_
	pricklypear	OPUNT	Opuntia	0–28	-
	Mexican bladdersage	SAME	Salazaria mexicana	0–28	_
	American threefold	TRCA8	Trixis californica	0–28	
Tree					
21				0–2	
	Joshua tree	YUBR	Yucca brevifolia	0–2	_

Type locality

Location 1: Mohave County, AZ					
Township/Range/Section	T35 N. R16 W. S2				

Contributors

Larry D. Ellicott Stephen Cassady Steve Barker

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills:
- 2. Presence of water flow patterns:
- 3. Number and height of erosional pedestals or terracettes:
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):
- 5. Number of gullies and erosion associated with gullies:
- 6. Extent of wind scoured, blowouts and/or depositional areas:
- 7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of

- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):
- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):
- 14. Average percent litter cover (%) and depth (in):
- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annualproduction):
- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
- 17. Perennial plant reproductive capability: