

## Ecological site R030XC013NV LOAMY BOTTOM 11-13 P.Z.

Last updated: 2/25/2025  
Accessed: 12/18/2025

---

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

### Ecological site concept

This site occurs on inset fans associated with drainageways. Slopes range from 2 to 4 percent. Elevations range from 5500 to 7500. The soils are deep to very deep, and well drained. Surface soils are thick, fertile and moderately fine to medium textured.

Please refer to group concept R030XC033NV to view the provisional STM.

### Associated sites

F030XC240NV	<b>Pinus monophylla/Cercocarpus ledifolius-Quercus gambelii/Poa fendleriana</b>
R030XC024NV	<b>GRAVELLY FAN 9-11 P.Z.</b>

**Table 1. Dominant plant species**

Tree	Not specified
Shrub	(1) <i>Artemisia tridentata ssp. vaseyana</i>
Herbaceous	(1) <i>Leymus cinereus</i>

### Physiographic features

This site occurs on inset fans associated with drainageways. Slopes range from 2 to 4 percent. Elevations range from 5500 to 7500.

**Table 2. Representative physiographic features**

Landforms	(1) Inset fan
Flooding duration	Very brief (4 to 48 hours)
Flooding frequency	Rare
Elevation	1,676–2,286 m
Slope	2–4%
Aspect	Aspect is not a significant factor

## Climatic features

The primary air masses affecting the Spring Mountains are cold maritime polar air from the Gulf of Alaska and warmer, moist maritime subtropical air from lower latitudes. Occasionally there are invasions of cold continental polar air from northern Canada or the Rocky Mountains. Precipitation in the area results primarily from the passage of cyclones with associated fronts during fall, winter and spring; from closed cyclones in late winter and spring; and from the flow of moist tropical air from the southeast to the southwest quadrant in the summer.

The mean annual precipitation is 11 to 13 inches; mean annual air temperature is 45 to 50 degrees F., and the frost-free season is 90 to 130 days. There are no climate stations available for this site.

**Table 3. Representative climatic features**

Frost-free period (average)	130 days
Freeze-free period (average)	
Precipitation total (average)	330 mm

## Influencing water features

This site is influenced by runoff from snowmelt and flash flooding as a result of summer convective storms.

## Soil features

The soils are deep to very deep, and well drained. Surface soils are thick, fertile and moderately fine to medium textured. The available water capacity is low. Additional moisture is received on this site during flash flooding and as run-in from higher landscape positions. The soil temperature regime is mesic and soil moisture regime is xeric. The soils are classified as Xeric Haplocalcids.

**Table 4. Representative soil features**

Surface texture	(1) Gravelly sandy loam
Family particle size	(1) Loamy
Drainage class	Well drained
Permeability class	Moderate
Soil depth	183 cm
Surface fragment cover <=3"	15–25%
Surface fragment cover >3"	1–5%
Available water capacity (0-101.6cm)	5.33–5.59 cm
Calcium carbonate equivalent (0-101.6cm)	5–15%
Electrical conductivity (0-101.6cm)	0–2 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0–5
Soil reaction (1:1 water) (0-101.6cm)	7.4–9
Subsurface fragment volume <=3" (Depth not specified)	25–41%
Subsurface fragment volume >3" (Depth not specified)	3–8%

## Ecological dynamics

Please refer to group concept R030XC033NV to view the provisional STM.

Changes in plant composition can occur with improper grazing that will reduce perennial grass production and increase basin big sagebrush and rubber rabbitbrush. Species likely to invade this site include cheatgrass. Lack of natural occurring wildfires may result in an invasion of pinyon and juniper.


### Fire Ecology:

Grassland communities with a basin wildrye component historically experienced mostly infrequent to frequent stand replacing fires. Grassland vegetation types experienced both short fire intervals of less than 35 years as well as intervals ranging from 35 to 100 years, depending on climate and ignition sources. Presettlement fire return intervals in mountain big sagebrush communities varied from 15 to 25 years. Basin wildrye is top-killed by fire. Older basin wildrye plants with large proportions of dead material within the perennial crown can be expected to show higher mortality due to fire than younger plants having


little debris. Basin wildrye is generally tolerant of fire but may be damaged by early season fire combined with dry soil conditions. Creeping wildrye is top-killed by fire. Creeping wildrye is generally tolerant of fire but may be damaged by early season fire combined with dry soil conditions. Indian ricegrass can be killed by fire, depending on severity and season of burn. Indian ricegrass reestablishes on burned sites through seed dispersed from adjacent unburned areas. Mountain big sagebrush is highly susceptible to injury from fire. Plants are readily killed in all seasons, even light severity fires. Mountain big sagebrush plants top-killed by fire will not resprout. Rubber rabbitbrush is often top-killed by fire. Rubber rabbitbrush is a fire-adapted species that is typically unharmed or enhanced by fire. Recovery time is often rapid to very rapid. Rubber rabbitbrush is often one of the first species to colonize burned areas by sprouting or from off-site seed.

State and transition model

Ecosystem states



State 1 submodel, plant communities



State 1  
Reference Plant Community

Community 1.1  
Reference Plant Community

The reference plant community is dominated by basin wildrye. Creeping wildrye and basin big sagebrush are other important species. Potential vegetative composition is approximately 70 percent grasses and grass-like plants, 5 percent forbs and 25 percent shrubs. Approximate ground cover (basal and canopy) is 40 to 60 percent. Total annual air-dry production is 2500 pounds on favorable years, 1500 pounds on normal years, and 1000 pounds on unfavorable years.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	785	1177	1961
Shrub/Vine	280	420	701
Forb	56	84	140
<b>Total</b>	<b>1121</b>	<b>1681</b>	<b>2802</b>

**Table 6. Ground cover**

Tree foliar cover	0%
Shrub/vine/liana foliar cover	10-20%
Grass/grasslike foliar cover	30-45%
Forb foliar cover	1-5%
Non-vascular plants	0%
Biological crusts	0%
Litter	10-20%
Surface fragments >0.25" and <=3"	15-25%
Surface fragments >3"	1-5%
Bedrock	0%
Water	0%
Bare ground	15-30%

**Table 7. Canopy structure (% cover)**

Height Above Ground (M)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.15	—	1-5%	1-2%	1-2%
>0.15 <= 0.3	—	1-5%	1-5%	1-5%
>0.3 <= 0.6	—	5-10%	20-30%	—
>0.6 <= 1.4	—	10-20%	30-40%	—
>1.4 <= 4	—	—	—	—
>4 <= 12	—	—	—	—
>12 <= 24	—	—	—	—
>24 <= 37	—	—	—	—
>37	—	—	—	—

## Additional community tables

Table 8. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
<b>Grass/Grasslike</b>					
1	<b>Primary Perennial Grasses</b>			790–1177	
	basin wildrye	LECI4	<i>Leymus cinereus</i>	673–925	–
	beardless wildrye	LETR5	<i>Leymus triticoides</i>	84–168	–
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	34–84	–
2	<b>Secondary Perennial Grasses</b>			84–168	
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	9–50	–
	sedge	CAREX	<i>Carex</i>	9–50	–
	squirreltail	ELEL5	<i>Elymus elymoides</i>	9–50	–
<b>Forb</b>					
3	<b>Primary Perennial Forbs</b>			50–84	
	lupine	LUPIN	<i>Lupinus</i>	50–84	–
4	<b>Secondary Perennial Forbs</b>			1–84	
	milkvetch	ASTRA	<i>Astragalus</i>	9–34	–
	mariposa lily	CALOC	<i>Calochortus</i>	9–34	–
	cryptantha	CRYPT	<i>Cryptantha</i>	9–34	–
	phlox	PHLOX	<i>Phlox</i>	9–34	–
	desert globemallow	SPAM2	<i>Sphaeralcea ambigua</i>	9–34	–
<b>Shrub/Vine</b>					
5	<b>Primary Shrubs</b>			303–420	
	mountain big sagebrush	ARTRV	<i>Artemisia tridentata</i> ssp. <i>vaseyana</i>	252–336	–
	rubber rabbitbrush	ERNA10	<i>Ericameria nauseosa</i>	50–84	–
	lupine	LUPIN	<i>Lupinus</i>	56–84	–
	phlox	PHLOX	<i>Phlox</i>	0–6	–
	desert globemallow	SPAM2	<i>Sphaeralcea ambigua</i>	0–6	–
	milkvetch	ASTRA	<i>Astragalus</i>	0–6	–
	mariposa lily	CALOC	<i>Calochortus</i>	0–6	–

	cryptantha	CRYPT	<i>Cryptantha</i>	0–6	–
6	<b>Secondary Shrubs</b>			1–84	
	mormon tea	EPVI	<i>Ephedra viridis</i>	17–34	–
	Nevada broomsage	LELA4	<i>Lepidospartum latisquamum</i>	17–34	–
	rose	ROSA5	<i>Rosa</i>	17–34	–

## Animal community

### Livestock Interpretations:

This site is suited to livestock grazing. Wild horses may use this site year round if water is available. Grazing management should be keyed to basin wildrye production. The early growth and abundant production of basin wildrye make it a valuable source of forage for livestock. It is important forage for cattle and is readily grazed by cattle and horses in early spring and fall. Though coarse-textured during the winter, basin wildrye may be utilized more frequently by livestock and wildlife when snow has covered low shrubs and other grasses. Creeping wildrye can be used for forage and is very palatable to all livestock. Once established it is very rhizomatous and maintains stands for many years. Indian ricegrass is highly palatable to all classes of livestock in both green and cured condition. It supplies a source of green feed before most other native grasses have produced much new growth. Mountain big sagebrush is eaten by domestic livestock but has long been considered to be of low palatability, and a competitor to more desirable species. In general, livestock forage only lightly on rubber rabbitbrush during the summer, but winter use can be heavy in some locations. Fall use is variable, but flowers are often used by livestock. A few leaves and the more tender stems may also be used.

Stocking rates vary over time depending upon season of use, climate variations, site, and previous and current management goals. A safe starting stocking rate is an estimated stocking rate that is fine tuned by the client by adaptive management through the year and from year to year.

### Wildlife Interpretations:

Basin wildrye provides winter forage for mule deer, though use is often low compared to other native grasses. Basin wildrye provides summer forage for black-tailed jackrabbits. Because basin wildrye remains green throughout early summer, it remains available for small mammal forage for longer time than other grasses. Creeping wildrye is used for forage for many wildlife species and is often used for cover. Indian ricegrass is eaten by pronghorn in moderate amounts whenever available. A number of heteromyid rodents inhabiting desert rangelands show preference for seed of Indian ricegrass. Indian ricegrass is an important component of jackrabbit diets in spring and summer. Indian ricegrass seed provides food for many species of birds. Doves, for example, eat large amounts of shattered Indian ricegrass seed lying on the ground. Mountain big sagebrush is highly preferred and nutritious winter forage for mule deer and elk. Wildlife forage only lightly on rubber rabbitbrush during the summer, but winter use can be heavy in some

locations. Fall use is variable, but flowers are often used by wildlife. A few leaves and the more tender stems may also be used. The forage value of rubber rabbitbrush varies greatly among subspecies and ecotypes.

Sagebrush-grassland communities provide critical sage-grouse breeding and nesting habitats. Meadows surrounded by sagebrush may be used as feeding and strutting grounds. Sagebrush is a crucial component of their diet year-round, and sage-grouse select sagebrush almost exclusively for cover. Sage-grouse prefer mountain big sagebrush and Wyoming big sagebrush communities to basin big sagebrush communities.

## Hydrological functions

Rills – Few to none present due to high cover of vegetation.

Water flow Patterns – Short, not connected.

Pedestals and/or Terracettes – Few to none.

Gullies – None present.

Low runoff due to plant cover and composition. Plant community composition contributes to infiltration and reduces runoff.

## Recreational uses

This site is used for camping and hiking.

## Type locality

Location 1: Clark County, NV	
Township/Range/Section	T18 S R55 E S29 NW
UTM zone	N
UTM northing	4024973n
UTM easting	605292e
General legal description	Near the junction of Wheeler Pass and Wheeler Well roads, Spring Mountains, Clark County, Nevada.

## Other references

Clokey, I. 1951. Flora of the Charleston Mountains, Clark County, Nevada. University of California Press, Berkeley and Los Angeles.

Fire Effects Information System [Online] <http://www.fs.fed.us/feis>.

Glenne, G. and Johnson, D. 2002. Guide to Species of Concern in the Spring Mountains National Recreation Area, Clark and Nye Counties, Nevada. USFS, Las Vegas, NV.

Nachlinger, J. and G. Reese. 1996. Plant Community Classification of the Spring



Mountains National Recreation Area, Clark and Nye Counties, Nevada. The Nature Conservancy. Reno, Nevada.

## Contributors

TJ WOLFE

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	P NOVAK-ECHENIQUE
Contact for lead author	State Rangeland Management Specialist
Date	06/22/2006
Approved by	Sarah Quistberg
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:** None to rare. A few may occur after summer convection storms or rapid snowmelt.

- 
2. **Presence of water flow patterns:** None to few. A few may occur after summer convection storms or rapid snowmelt flow patterns are short (<1 m) and stable.

- 
3. **Number and height of erosional pedestals or terracettes:** None to rare. Occurrence is usually limited to areas of water flow patterns.
-

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare Ground  $\pm$  20-30%; surface rock fragments 30-40%.
- 
5. **Number of gullies and erosion associated with gullies:** Gullies are rare to common depending on severity of associated stream channel entrenchment. Gullies and head cuts are healing or stable.
- 
6. **Extent of wind scoured, blowouts and/or depositional areas:** None
- 
7. **Amount of litter movement (describe size and distance expected to travel):** Fine litter (foliage of grasses and annual & perennial forbs) only expected to move during periods of flooding by adjacent streams. Persistent litter (large woody material) will remain in place except during large flooding events.
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil stability values range from 4 to 6.
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Soil surface structure is typically very thick platy. Soil surface colors are dark browns and the soils have thick mollic epipedons. Organic matter can range from 3 to 5 percent for much of the upper 20 inches. (OM values derived from lab characterization data.)
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Deep-rooted perennial grasses slow runoff and increase infiltration. Tall stature and relatively coarse foliage of wildrye and associated litter break raindrop impact and provide opportunity for snow catch and snow accumulation on site.
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile**

**features which may be mistaken for compaction on this site):** None - Platy or subangular blocky structure are not to be interpreted as compaction.

---

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Tall-statured, deep-rooted, cool season, perennial bunchgrasses, basin wildrye

Sub-dominant: tall shrubs >warm season rhizomatous grasses > deep-rooted, cool season, perennial forbs >warm season rhiz fibrous, shallow-rooted, cool season, annual and perennial forbs

Other:

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Dead branches within individual shrubs common; standing dead shrub canopy material may be as much as 25% of total woody canopy.
- 

14. **Average percent litter cover (%) and depth ( in):** Between plant interspaces ( $\pm$  40-50%) and litter depth is  $> \frac{1}{2}$  inch.
- 

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** For normal or average growing season (through June)  $\pm$  1500 lbs/ac; Favorable years: 2500 lbs/ac; Unfavorable years: 1000 lbs/ac
- 

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing**

**what is NOT expected in the reference state for the ecological site:** Potential invaders include: annual mustards, cheatgrass, single-leaf pinyon and Utah juniper.

---

17. **Perennial plant reproductive capability:** All functional groups should reproduce in most years.
-