

Ecological site R030XC308AZ **Limy Fan 10-13" p.z.**

Accessed: 05/12/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 030X–Mojave Basin and Range

AZ CRA 30.3 – Upper Mohave Desert

Elevations range from 2800 to 4500 feet and precipitation averages 9 to 12 inches per year. Vegetation includes Joshua tree, blackbrush, creosotebush, ratany, bush muhly, big galleta, black grama, desert needlegrass, and Indian ricegrass. The soil temperature regime is thermic and the soil moisture regime is typic aridic. This unit occurs within the Basin and Range Province and is characterized by broad basins, valleys, and old lakebeds. Widely spaced mountains trending north to south occur throughout the area. Isolated, short mountain ranges are separated by an aggraded desert plain. The mountains are fault blocks that have been tilted up. Long alluvial fans coalesce with dry lakebeds between some of the ranges.

Associated sites

R030XC305AZ	Coarse Sandy Loam 10-13" p.z.
R030XC311AZ	Limy Upland 10-13" p.z.
R030XC317AZ	Sandy Bottom 10-13" p.z.

Table 1. Dominant plant species

Tree	Not specified
Shrub	(1) <i>Atriplex canescens</i>
Herbaceous	(1) <i>Pleuraphis rigida</i> (2) <i>Muhlenbergia porteri</i>

Physiographic features

This ecological site is found in an upland position on fan terraces and toeslopes of steep escarpments merging onto stream terraces.

Table 2. Representative physiographic features

Landforms	(1) Fan (2) Terrace
Flooding frequency	None
Ponding frequency	None
Elevation	914–1,280 m
Slope	1–6%
Aspect	Aspect is not a significant factor

Climatic features

The climate is arid and warm. Annual precipitation ranges from 10 to 13 inches. About 65 percent of the rainfall comes from October through May as gentle rain from Pacific storms which may last for a couple of days. The rest of the rainfall comes during the summer monsoon season from July through September as spotty, brief, intense thunderstorms. Snow rarely falls, and only remains on the ground a few hours at most. Annual air temperature ranges from 46 to 76 degrees F. The average frost-free period ranges from 121 to 231 days.

Table 3. Representative climatic features

Frost-free period (average)	231 days
Freeze-free period (average)	269 days
Precipitation total (average)	330 mm

Influencing water features

Soil features

The soil of this ecological site is moderately deep to deep. The soil surface textures include very fine sandy loam, sandy loam, very gravelly loam, and gravelly fine sandy loam. Subsoil textures are very fine sandy loam, fine sandy loam, silt loam cobbly very fine sandy loam, extremely very fine sandy loam, sandy clay loam, sandy loam, gravelly sandy loam, and gravelly sandy clay loam. The soil's available water capacity is low to high. Its erosion hazard by water is slight to severe and by wind is moderate to severe. The soil is non-saline, non-sodic with pH 7.4-8.6. The soil moisture regime is typical aridic and temperature regime is thermic.

A typical soil profile is:

0 to 2 inches; very fine sandy loam

2 to 27 inches; very fine sandy loam

27 to 37 inches; silt loam

37 to 53 inches; very fine sandy loam

53 to 60 inches; very gravelly fine sandy loam

Soil taxonomic classifications include Coarse-loamy, mixed thermic Typic Paleorthids and, Coarse-loamy, mixed, thermic Typic Camborthids

This ecological site has been correlated to map unit 627088, Poachie soil, Mohave County, AZ, Central Part SSA.

Table 4. Representative soil features

Surface texture	(1) Cobbly very fine sandy loam (2) Gravelly sandy loam (3) Loam
Family particle size	(1) Loamy
Drainage class	Well drained
Permeability class	Moderately slow to moderately rapid
Soil depth	152 cm
Surface fragment cover <=3"	45–70%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	13.97–20.32 cm
Calcium carbonate equivalent (0-101.6cm)	2–35%
Electrical conductivity (0-101.6cm)	0 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0
Soil reaction (1:1 water) (0-101.6cm)	7.4–8.4
Subsurface fragment volume <=3" (Depth not specified)	0–20%
Subsurface fragment volume >3" (Depth not specified)	0%

Ecological dynamics

The historic climax plant community (HCPC) for a site in North America is the plant community that existed at the time of European immigration and settlement. It is the plant community that was best adapted to the unique combination of environmental factors associated with the site. The historic climax plant community was in dynamic equilibrium with its environment. It is the plant community that was able to avoid displacement by the suite of disturbances and disturbance patterns (magnitude and frequency) that naturally occurred within the area occupied by the site. Natural disturbances, such as drought, fire, grazing by native fauna, and insects, were inherent in the development and maintenance of these plant communities. The effects of these disturbances are part of the range of characteristics of the site that contribute to that dynamic equilibrium. Fluctuations in plant community structure and function caused by the effects of these natural disturbances establish the boundaries of dynamic equilibrium. They are accounted for as part of the range of characteristics for an ecological site. Some sites may have a small range of variation, while others have a large range.

The historic climax plant community of an ecological site is not a precise assemblage of species for which the proportions are the same from place to place or from year to year. In all plant communities, variability is apparent in productivity and occurrence of individual species. Spatial boundaries of the communities; however, can be recognized by characteristic patterns of species composition, association, and community structure. The HCPC for this ecological site has been estimated by sampling relict or relatively undisturbed sites and/or reviewing historic records.

Plant communities that are subjected to abnormal disturbances and physical site deterioration or that are protected from natural influences, such as fire and grazing, for long periods seldom typify the historic climax plant community. The physical site deterioration caused by the abnormal disturbance results in the crossing of a threshold or

Historic Climax Plant Community

The dominant aspect of this plant community is grassland with shrubs. Big galleta and sand dropseed are the major grasses. Shrubs include fourwing saltbush, Anderson wolfberry, Nevada Mormon tea and winterfat.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	189	269	353
Shrub/Vine	127	191	256
Forb	20	45	64
Total	336	505	673

Table 6. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	1-3%
Grass/grasslike foliar cover	0-2%
Forb foliar cover	0-2%
Non-vascular plants	0%
Biological crusts	0%
Litter	0%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%

Table 7. Canopy structure (% cover)

Height Above Ground (M)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.15	—	—	—	1-3%
>0.15 <= 0.3	—	—	13-17%	—
>0.3 <= 0.6	—	—	—	—
>0.6 <= 1.4	—	9-11%	—	—
>1.4 <= 4	—	—	—	—
>4 <= 12	—	—	—	—
>12 <= 24	—	—	—	—
>24 <= 37	—	—	—	—
>37	—	—	—	—

Figure 5. Plant community growth curve (percent production by month). AZ3024, 30.3 10-13" p.z. upland sites. Growth begins in the spring and continues through the summer..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	1	8	18	18	11	14	20	8	2	0	0

Additional community tables

Table 8. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1				76–126	
	big galleta	PLRI3	<i>Pleuraphis rigida</i>	76–126	–
2				25–50	
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	25–50	–
3				25–50	
	bush muhly	MUPO2	<i>Muhlenbergia porteri</i>	25–50	–
4				4–25	
	black grama	BOER4	<i>Bouteloua eriopoda</i>	4–25	–
5				4–25	
	Grass, perennial	2GP	<i>Grass, perennial</i>	4–25	–
6				0–25	
	Grass, annual	2GA	<i>Grass, annual</i>	0–25	–
Forb					
7				4–25	
	desert globemallow	SPAM2	<i>Sphaeralcea ambigua</i>	4–25	–
8				4–16	
	Forb, perennial	2FP	<i>Forb, perennial</i>	4–16	–
9				4–10	
	Forb, annual	2FA	<i>Forb, annual</i>	4–10	–
Shrub/Vine					
10				25–50	
	fourwing saltbush	ATCA2	<i>Atriplex canescens</i>	25–50	–
11				4–25	
	water jacket	LYAN	<i>Lycium andersonii</i>	4–25	–
12				0–10	
	Whipple cholla	CYWH	<i>Cylindropuntia whipplei</i>	0–10	–
13				0–10	
	banana yucca	YUBA	<i>Yucca baccata</i>	0–10	–
14				4–25	
	Nevada jointfir	EPNE	<i>Ephedra nevadensis</i>	4–25	–
15				0–10	
	buckhorn cholla	CYACA2	<i>Cylindropuntia acanthocarpa</i> var. <i>acanthocarpa</i>	0–10	–
16				4–50	
	winterfat	KRLA2	<i>Krascheninnikovia lanata</i>	4–50	–
17				0–25	
	shadscale saltbush	ATCO	<i>Atriplex confertifolia</i>	0–25	–
18				0–4	
	catclaw acacia	ACGR	<i>Acacia greggii</i>	0–4	–

19				0–10	
	littleleaf ratany	KRER	<i>Krameria erecta</i>	0–10	–
20				0–25	
	creosote bush	LATR2	<i>Larrea tridentata</i>	0–25	–
21				4–25	
	Shrub, other	2S	<i>Shrub, other</i>	4–25	–

Animal community

Wildlife found on this ecological site includes desert cottontail rabbit, desert pocket mouse, lizards, blacktail jackrabbit, snakes and coyote.

Type locality

Location 1: Mohave County, AZ	
Township/Range/Section	T32N R8W S1
General legal description	Vulcas Throne Quad., north end of Prospect Valley on the Hualapai Indian Reservation.

Contributors

Harmon Hodgkinson
Harmon S. Hodgkinson
Larry D. Ellicott
Stephen Cassady

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

-
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**
-
5. **Number of gullies and erosion associated with gullies:**
-
6. **Extent of wind scoured, blowouts and/or depositional areas:**
-
7. **Amount of litter movement (describe size and distance expected to travel):**
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-
14. **Average percent litter cover (%) and depth (in):**
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-**

production):

16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
-

17. **Perennial plant reproductive capability:**
-