

Ecological site R030XC321AZ

Sandy Loam Upland 10-13" p.z. Fine

Accessed: 05/18/2024

General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 030X–Mojave Basin and Range

AZ CRA 30.3 – Upper Mohave Desert

Elevations range from 2800 to 4500 feet and precipitation averages 9 to 12 inches per year. Vegetation includes Joshua tree, blackbrush, creosotebush, ratany, bush muhly, big galleta, black grama, desert needlegrass, and Indian ricegrass. The soil temperature regime is thermic and the soil moisture regime is typic aridic. This unit occurs within the Basin and Range Province and is characterized by broad basins, valleys, and old lakebeds. Widely spaced mountains trending north to south occur throughout the area. Isolated, short mountain ranges are separated by an aggraded desert plain. The mountains are fault blocks that have been tilted up. Long alluvial fans coalesce with dry lakebeds between some of the ranges.

Associated sites

R030XC305AZ	Coarse Sandy Loam 10-13" p.z.
-------------	-------------------------------

Table 1. Dominant plant species

Tree	Not specified
------	---------------

Shrub	Not specified
Herbaceous	(1) <i>Pleuraphis rigida</i> (2) <i>Bouteloua eriopoda</i>

## Physiographic features

This ecological site is found in an upland position on shoulders, summits and sideslopes of fan terraces.

**Table 2. Representative physiographic features**

Landforms	(1) Terrace (2) Fan
Flooding frequency	None
Ponding frequency	None
Elevation	732–1,219 m
Slope	1–12%
Aspect	Aspect is not a significant factor

## Climatic features

The climate is arid and warm. Annual precipitation ranges from 10 to 13 inches. About 65 percent of the rainfall comes from October through May as gentle rain from Pacific storms which may last for a couple of days. The rest of the rainfall comes during the summer monsoon season from July through September as spotty, brief, intense thunderstorms. Snow rarely falls, and only remains on the ground a few hours at most. Annual air temperature ranges from 46 to 76 degrees F. The average frost-free period ranges from 121 to 231 days.

**Table 3. Representative climatic features**

Frost-free period (average)	231 days
Freeze-free period (average)	269 days
Precipitation total (average)	330 mm

## Influencing water features

### Soil features

The soil of this ecological site is deep to very deep with surface textures of sandy loam, gravelly sandy loam and gravelly loam. Subsoil textures are sandy clay loam, coarse sandy loam, gravelly sandy loam, gravelly sandy clay loam, very gravelly sandy loam, sandy loam, extremely gravelly sand, and gravelly loamy sand. The soil parent material is alluvium from granite and igneous rocks. Geologic formation varies. Soil available water capacity is moderate to high. The soil's erosion hazard from water is slight to moderate and from wind is slight. The soil is non-sodic, non-saline with pH of 7.4-8.0 (mildly to moderately alkaline). The soil moisture regime is typical aridic and temperature regime is thermic.

A typical soil profile is:

A-0 to 4 inches; gravelly loamy sand

BA-4 to 10 inches; gravelly sandy loam

2Bt1-10 to 26 inches; gravelly sandy clay

2Bt2-26 to 40 inches; gravelly sandy clay loam

3Bkn-40 to 60 inches; very gravelly sand

Soil taxonomic classifications include Fine, mixed, superactive, thermic Typic Haplargids; Fine-loamy, mixed, superactive, thermic Ustic Haplargids; Fine-loamy, mixed, superactive, thermic Typic Haplargids; Fine, mixed,

superactive, thermic Ustic Haplargids.

Soil mapping units correlated to this ecological site are 627125, Vekol family, Mohave County, AZ, Southern Part and 697057, 627090, 697165, Courtland family, Dutchflat, Whitehouse soils, Mohave County, AZ, Central Part.

**Table 4. Representative soil features**

Parent material	(1) Alluvium–granite
Surface texture	(1) Gravelly sandy loam (2) Gravelly loam (3) Sandy loam
Family particle size	(1) Loamy
Drainage class	Well drained
Permeability class	Moderately slow to moderate
Soil depth	152 cm
Surface fragment cover <=3"	20–45%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	12.7–17.78 cm
Calcium carbonate equivalent (0-101.6cm)	2–10%
Electrical conductivity (0-101.6cm)	0 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0
Soil reaction (1:1 water) (0-101.6cm)	7.4–8.4
Subsurface fragment volume <=3" (Depth not specified)	20–70%
Subsurface fragment volume >3" (Depth not specified)	0–5%

## Ecological dynamics

The historic climax plant community (HCPC) for a site in North America is the plant community that existed at the time of European immigration and settlement. It is the plant community that was best adapted to the unique combination of environmental factors associated with the site. The historic climax plant community was in dynamic equilibrium with its environment. It is the plant community that was able to avoid displacement by the suite of disturbances and disturbance patterns (magnitude and frequency) that naturally occurred within the area occupied by the site. Natural disturbances, such as drought, fire, grazing by native fauna, and insects, were inherent in the development and maintenance of these plant communities. The effects of these disturbances are part of the range of characteristics of the site that contribute to that dynamic equilibrium. Fluctuations in plant community structure and function caused by the effects of these natural disturbances establish the boundaries of dynamic equilibrium. They are accounted for as part of the range of characteristics for an ecological site. Some sites may have a small range of variation, while others have a large range.

The historic climax plant community of an ecological site is not a precise assemblage of species for which the proportions are the same from place to place or from year to year. In all plant communities, variability is apparent in productivity and occurrence of individual species. Spatial boundaries of the communities; however, can be recognized by characteristic patterns of species composition, association, and community structure. The HCPC for this ecological site has been estimated by sampling relict or relatively undisturbed sites and/or reviewing historic records.

Plant communities that are subjected to abnormal disturbances and physical site deterioration or that are protected

from natural influences, such as fire and grazing, for long periods seldom typify the historic climax plant community. The physical site deterioration caused by the abnormal disturbance results in the crossing of a threshold or irreversible boundary to another state, or equilibrium, for the ecological site. There may be multiple thresholds and states possible for an ecological site, determined by the type and or severity of abnormal disturbance. The known states and transition pathways for this ecological site are described in the accompanying state and transition model. The Plant Community Plant Species Composition table provides a list of species and each species or group of species' annual production in pounds per acre (air-dry weight) expected in a normal rainfall year. Low and high production yields represent the modal range of variability for that species or group of species across the extent of the ecological site.

## State 1

### Historic Climax Plant Community

Community 1.1  
Historic Climax Plant Community

The dominant aspect of this plant community is a desert grassland with shrubs. Major grasses are big galleta, bush muhly and black grama. Shrubs include fourwing saltbush, range ratany (*Krameria erecta*), Anderson wolfberry *Lycium Andesonii*), rayless goldenhead and buckhorn cholla.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	63	185	467
Shrub/Vine	43	129	351
Forb	7	22	78
Total	113	336	896

Table 6. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	0-2%
Grass/grasslike foliar cover	1-3%
Forb foliar cover	0%
Non-vascular plants	0%
Biological crusts	0%
Litter	0%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%

Table 7. Canopy structure (% cover)

Height Above Ground (M)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.15	—	—	—	0-2%
>0.15 <= 0.3	—	—	—	—
>0.3 <= 0.6	—	—	8-12%	—
>0.6 <= 1.4	—	8-12%	—	—
>1.4 <= 4	—	—	—	—
>4 <= 12	—	—	—	—
>12 <= 24	—	—	—	—
>24 <= 37	—	—	—	—
>37	—	—	—	—

Figure 5. Plant community growth curve (percent production by month).  
AZ3024, 30.3 10-13" p.z. upland sites. Growth begins in the spring and continues through the summer..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	1	8	18	18	11	14	20	8	2	0	0

## Additional community tables

Table 8. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
<b>Grass/Grasslike</b>					
1				50–84	
	big galleta	PLRI3	<i>Pleuraphis rigida</i>	50–84	–
2				17–34	
	black grama	BOER4	<i>Bouteloua eriopoda</i>	17–34	–
3				17–34	
	bush muhly	MUPO2	<i>Muhlenbergia porteri</i>	17–34	–
4				3–7	
	Parish's threeawn	ARPUP5	<i>Aristida purpurea</i> var. <i>parishii</i>	3–7	–
5				0–7	
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	0–7	–
6				3–17	
	spike dropseed	SPCO4	<i>Sporobolus contractus</i>	3–17	–
7				0–7	
	low woollygrass	DAPU7	<i>Dasyochloa pulchella</i>	0–7	–
8				3–17	
	Grass, perennial	2GP	<i>Grass, perennial</i>	3–17	–
9				3–17	
	Grass, annual	2GA	<i>Grass, annual</i>	3–17	–
<b>Forb</b>					
10				3–17	
	desert globemallow	SPAM2	<i>Sphaeralcea ambigua</i>	3–17	–
11				3–10	
	Forb, perennial	2FP	<i>Forb, perennial</i>	3–10	–
12				3–13	
	Forb, annual	2FA	<i>Forb, annual</i>	3–13	–
<b>Shrub/Vine</b>					
13				3–17	
	fourwing saltbush	ATCA2	<i>Atriplex canescens</i>	3–17	–
14				0–7	
	Eastern Mojave buckwheat	ERFAP	<i>Eriogonum fasciculatum</i> var. <i>polifolium</i>	0–7	–
15				3–10	
	Mexican bladdersage	SAME	<i>Salazaria mexicana</i>	3–10	–
16				3–17	
	rayless goldenhead	ACSP	<i>Acamptopappus sphaerocephalus</i>	3–17	–
17				3–7	
	Nevada jointfir	EPNE	<i>Ephedra nevadensis</i>	3–7	–
18				0–17	
	broom snakeweed	GUSA2	<i>Gutierrezia sarothrae</i>	0–17	–

19				3–10	
	burrobrush	HYSA	<i>Hymenoclea salsola</i>	3–10	–
20				3–17	
	littleleaf ratany	KRER	<i>Krameria erecta</i>	3–17	–
21				3–17	
	water jacket	LYAN	<i>Lycium andersonii</i>	3–17	–
22				3–10	
	banana yucca	YUBA	<i>Yucca baccata</i>	3–10	–
23				3–17	
	banana yucca	YUBA	<i>Yucca baccata</i>	3–17	–
24				0–3	
	beavertail pricklypear	OPBA2	<i>Opuntia basilaris</i>	0–3	–
25				3–10	
	branched pencil cholla	CYRA9	<i>Cylindropuntia ramosissima</i>	3–10	–
26				3–17	
	Shrub, other	2S	<i>Shrub, other</i>	3–17	–

## Animal community

Wildlife species found on this ecological site include coyote, lizards, desert cottontail, snakes, blacktail jackrabbit, and ants.

## Type locality

Location 1: Mohave County, AZ	
Township/Range/Section	T24N R17W S3
General legal description	Elements Canyon Quad. On west side of Stockton Hill Road and along road to Coarra Ranch Headquarters.

## Contributors

Harmon Hodgkinson  
Harmon S. Hodgkinson  
Karlynn Huling  
Larry D. Ellicott  
Stephen Cassady

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	

Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:**

---

2. **Presence of water flow patterns:**

---

3. **Number and height of erosional pedestals or terracettes:**

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

---

5. **Number of gullies and erosion associated with gullies:**

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

---

7. **Amount of litter movement (describe size and distance expected to travel):**

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

---

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

---

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

---

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

---

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
- 

14. **Average percent litter cover (%) and depth ( in):**
- 

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
- 

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
- 

17. **Perennial plant reproductive capability:**
-