

## Ecological site R030XD006CA Abandoned Fan

Accessed: 05/14/2024

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

### MLRA notes

Major Land Resource Area (MLRA): 030X–Mojave Basin and Range

MLRA Description:

Major Land Resource Area (MLRA) 30, Mojave Desert, is found in southern California, southern Nevada, the extreme southwest corner of Utah and northwestern Arizona within the Basin and Range Province of the Intermontane Plateaus. The climate of the area is hot (primarily hyperthermic and thermic; however at higher elevations, generally above 5000 feet, mesic, cryic and frigid) and dry (aridic). Elevations range from below sea level to over 12,000 feet in the higher mountain areas found within the MLRA. Due to the extreme elevational range found within this MLRA, Land Resource Units (LRUs) were designated to group the MLRA into similar land units.

LRU Description:

This Land Resource Unit (designated by 'XD') is found on the eastern side of California. Elevations range from 400 to 2200 feet on average, but may be found up to 3600 feet on southern exposures. Precipitation ranges from 1 to 6 inches per year, but averages between 2-4 inches. This LRU is characterized primarily by the extreme aridity, hot temperatures, hyperthermic soil temperatures and low stature of widely spaced vegetation. Temperatures can reach over 110 degrees Fahrenheit for several weeks in July and August. Summer precipitation falls between July and September, ranging from 20-33% in the form of rain, and winter precipitation falls starting in November and ends between February and March, ranging from 56-70%, also mostly in the form of rain. Vegetation is primarily small,

widely-spaced, low-producing creosote bush (*Larrea tridentata*), burrobush (*Ambrosia dumosa*), and brittlebush (*Encelia farinosa*).

#### Ecological Site Concept –

This ecological site occurs on gently sloping alluvial fans and fan aprons at elevations of 670 to 2620 feet. Soils are very deep, and are composed of layers of sand, coarse sand and loamy sand with varying amounts of gravel and cobbles. This ecological site tends to occupy distal fan positions, far from sources of run-on, and this site typically has no sheet-flow from flash-flooding events.

Production reference value (RV) is 67 pounds per acre, and ranges from 21 to 191 pounds per acre depending on annual precipitation and annual forb production. Vegetation is sparse, and dominated by low-statured creosote bush (*Larrea tridentata*). A hyperthermic climate with deep sandy soils with no additional run-on from sheet flooding drives the vegetation community of this ecological site. The deep-rooted creosote bush is the only shrub capable of tolerating the extremely arid conditions.

The data in the following sections is from major (15% of mapunit or greater) components only.

### Classification relationships

Mojave Creosote Bush (Holland, 1986)

*Larrea tridentata* Shrubland Alliance (Sawyer et al. 2009).

### Associated sites

|             |  |
|-------------|--|
| R030XD039CA | <b>Coarse Gravelly Fans</b><br>This ecological site is found on adjacent occasionally flooded alluvial fans. Creosote bush ( <i>Larrea tridentata</i> ), burrobush ( <i>Ambrosia dumosa</i> ) and brittlebush ( <i>Encelia farinosa</i> ) are dominant.  |
| R030XD004CA | <b>Low-Production Hyperthermic Hills</b><br>This ecological site is found on steep sideslopes of adjacent fan remnants. Sparse vegetation is dominated by creosote bush ( <i>Larrea tridentata</i> ).  |
| R030XD008CA | <b>Hyperthermic Sandhill</b><br>This ecological site is found on adjacent sandhills. Big galleta ( <i>Pleuraphis rigida</i> ) and creosote bush ( <i>Larrea tridentata</i> ) are dominant.   |
| R030XD014CA | <b>Hyperthermic Sandy Plains</b><br>This ecological site is found on adjacent semi-active sandsheets. Big galleta ( <i>Pleuraphis rigida</i> ) is dominant.  |
| R030XD015CA | <b>Hyper-Arid Fans</b><br>This ecological site is found on adjacent fan aprons with rare surface flooding. Creosote bush ( <i>Larrea tridentata</i> ) and burrobush ( <i>Ambrosia dumosa</i> ) dominate.   |
| R030XD025CA | <b>Hyperthermic Sandsheets</b><br>This ecological site is found on adjacent sandsheets. Creosote bush ( <i>Larrea tridentata</i> ) and big galleta ( <i>Pleuraphis rigida</i> ) dominate.  |
| R030XD042CA | <b>Hyperthermic Shallow To Moderately Deep Fan Remnants</b><br>This ecological site is found on adjacent very stable fan remnants. Creosote bush ( <i>Larrea tridentata</i> ) is dominant.   |
| R030XY001CA | <b>Occasionally Flooded, Hyperthermic, Diffuse Ephemeral Stream</b><br>This ecological site is found on adjacent occasionally flooded drainageways. Creosote bush ( <i>Larrea tridentata</i> ) and Schott's dalea ( <i>Psoralea schottii</i> ) dominate. |
| R030XY092NV | <b>DESERT PATINA</b><br>This ecological site is found on adjacent fan remnants with desert pavement surface. Very sparse vegetation is dominated by creosote bush ( <i>Larrea tridentata</i> ).  |

### Similar sites

|             |  |
|-------------|--|
| R030XD042CA | <b>Hyperthermic Shallow To Moderately Deep Fan Remnants</b><br>This ecological site occurs on very stable fan remnants with a high degree of soil horizon development. It has lower production and shrub density.  |
| R030XD015CA | <b>Hyper-Arid Fans</b><br>This ecological site occurs on landform positions receiving more surface flooding. This site is more productive, shrub density is higher, and burrobush ( <i>Ambrosia dumosa</i> ) is co-dominant with creosote bush ( <i>Larrea tridentata</i> ). |

**Table 1. Dominant plant species**

|            |                              |
|------------|------------------------------|
| Tree       | Not specified                |
| Shrub      | (1) <i>Larrea tridentata</i> |
| Herbaceous | (1) <i>Plantago ovata</i>    |

## Physiographic features

This ecological site occurs on alluvial fans and fan aprons at elevations of 670 to 2620 feet, and slopes ranging from 0 to 8 percent. This site typically does not receive flooding, and runoff class is negligible to very low.

**Table 2. Representative physiographic features**

|                   |                                    |
|-------------------|------------------------------------|
| Landforms         | (1) Alluvial fan<br>(2) Fan apron  |
| Flooding duration | Brief (2 to 7 days)                |
| Elevation         | 204–799 m                          |
| Slope             | 0–8%                               |
| Aspect            | Aspect is not a significant factor |

## Climatic features

The climate of this ecological site is characterized by hot temperatures, aridity, and a bimodal precipitation pattern. Precipitation falls as rain, with 30 percent falling in summer between July and October, and 65 percent falling in winter between November and March. The mean annual precipitation is 3 to 5 inches and mean annual air temperature is 68 to 73 degrees F. The frost free period is 300 to 340 days.

Maximum and minimum monthly climate data for this ESD were generated by the Climate Summarizer ([http://www.nm.nrcs.usda.gov/technical/handbooks/nrph/Climate\\_Summarizer.xls](http://www.nm.nrcs.usda.gov/technical/handbooks/nrph/Climate_Summarizer.xls)) using data from the following climate stations (results are unweighted averages):

42598, Eagle Mountain, CA (Period of record = 1933 to 2011) [1]

43855, Hayfield Reservoir, CA (Period of record = 1933 to 2011) [1]

049099, Twentynine Palms, California (Period of record = 1935 to 2011) [1]

The data from multiple weather were combined to most accurately reflect the climatic conditions of this ecological site.

**Table 3. Representative climatic features**

|                               |          |
|-------------------------------|----------|
| Frost-free period (average)   | 340 days |
| Freeze-free period (average)  | 0 days   |
| Precipitation total (average) | 127 mm   |

## Influencing water features

### Soil features

The soils associated with this ecological site are very deep, somewhat excessively to excessively drained, and formed in alluvium from predominantly granitoid rocks. These soils are sandy or sandy-skeletal in the particle size control section, and permeability is moderately rapid to rapid. Surface textures include fine sand, gravelly loamy fine sand, fine sandy loam. Subsurface horizons (1 to 59 inches) are composed of layers of fine sand and sand textures with gravelly to very gravelly modifiers. Surface gravels (< 3 mm in diameter) range from 30 to 75 percent, and larger fragments range from 0 to 11 percent. Subsurface gravels by volume (for a depth of 0 to 59 inches) range from 1 to 50 percent and larger fragments by volume range from 0 to 25 percent.

This ecological site is associated with Pintobasin soils (mixed, hyperthermic Typic Torripsamments); Carrizo soils (sandy-skeletal, mixed, hyperthermic Typic Torriorthents); and Joetree soils (mixed, hyperthermic Typic Torripsamments). Carrizo soils are sandy and have greater than 35 percent rock fragments in the particle control section. Joetree and Pintobasin soils are sandy throughout with few rock fragments. The difference between the Joetree and the Pintobasin soils is that Joetree soils have an argillic horizon from 40 to 59 inches with sandy loam or sandy clay loam textures.

This ecological site is correlated with the following map units and soil components in the Joshua Tree National Park Soil Survey:

Mapunit;Mapunit name;Component;Local phase;Component percent  
1513;Carrizo-Rubylee complex, 1 to 4 percent slopes;Carrizo; 60  
2065;Dalelake-Aquapeak-Coxpin association, 2 to 8 percent slopes;Carrizo; 1  
2076;Oldale-Carrizo complex, 2 to 8 percent slopes;Carrizo;30; Pintobasin;very rarely flooded;2  
2077;Oldale-Carrizo association, 2 to 8 percent slopes;Carrizo; 25  
2060;Joetree-Dalelake-Pintobasin complex, 0 to 2 percent slopes;Joetree;very rarely flooded;35  
1514;Carrizo-Pintobasin-Rubylee complex, 0 to 4 percent slopes;Pintobasin;fine sandy loam;30  
1515;Pintobasin-Carrizo complex, 2 to 8 percent slopes;Pintobasin; 80  
1516;Pintobasin fine sandy loam, 0 to 2 percent slopes;Pintobasin;fine sandy loam;90  
1517;Pintobasin-Dalelake complex, 2 to 8 percent slopes;Pintobasin; 65  
1522;Pintobasin sand, 1 to 3 percent slopes, rarely flooded;Pintobasin; 5  
1525;Pintobasin complex, 2 to 4 percent slopes, 10  
1530;Dalelake fine sand, 0 to 4 percent slopes;Pintobasin; 5  
2060;Joetree-Dalelake-Pintobasin complex, 0 to 2 percent slopes;Pintobasin;fine sandy loam;25  
2068;Aquapeak-Carpetflat-Pintobasin complex, 0 to 4 percent slopes;Pintobasin; 15  
2076;Oldale-Carrizo complex, 2 to 8 percent slopes;Pintobasin;very rarely flooded;2  
2715;Dalelake-Sheephole-Pintobasin complex, 2 to 8 percent slopes;Pintobasin; 25

**Table 4. Representative soil features**

|   |  |
|---|--|
| Parent material                         | (1) Alluvium–granite   |
| Surface texture                         | (1) Fine sandy loam<br>(2) Gravelly loamy fine sand<br>(3) Fine sand |
| Family particle size                    | (1) Sandy  |
| Drainage class                          | Somewhat excessively drained to excessively drained                  |
| Permeability class                      | Moderately rapid to rapid  |
| Soil depth                              | 150 cm   |
| Surface fragment cover <=3"             | 30–75%   |
| Surface fragment cover >3"              | 0–11%  |
| Available water capacity<br>(0-101.6cm) | 2.29–9.65 cm   |

|  |            |
|--|------------|
| Calcium carbonate equivalent<br>(0-101.6cm)              | 0–5%       |
| Electrical conductivity<br>(0-101.6cm)                   | 0 mmhos/cm |
| Sodium adsorption ratio<br>(0-101.6cm)                   | 0          |
| Soil reaction (1:1 water)<br>(0-101.6cm)                 | 6–8.8      |
| Subsurface fragment volume <=3"<br>(Depth not specified) | 1–50%      |
| Subsurface fragment volume >3"<br>(Depth not specified)  | 0–25%      |

## Ecological dynamics

### Abiotic Factors

The abiotic factors driving this site are This ecological site occurs on gently sloping alluvial fans and fan aprons at elevations of 670 to 2620 feet. Soils are very deep, and are composed of layers of sand, coarse sand and loamy sand with varying amounts of gravel and cobbles. This site typically has no sheet-flow from flash-flooding events. A hyperthermic climate with deep sandy soils with no additional run-on from sheet flooding drives the vegetation community of this ecological site. The deep-rooted creosote bush is the only shrub capable of tolerating the extremely arid conditions of this site.

Creosote bush shrublands dominate fan piedmont landscapes at elevations below 4000 feet in the Mojave Desert (Rundel and Gibson 1996). In arid regions, the availability of moisture is the key resource driving the productivity and composition of vegetation (Noy-Meir 1973, McAuliffe 1994, Hamerlynk et al. 2000, Martre et al. 2002, Austin et al. 2004). Where soil temperature regimes are thermic (above approximately 2800 feet) and soil moisture availability is higher, shrub production, cover, density and diversity are higher (Bedford et al. 2009). Where the soil temperature regime is hyperthermic and moisture becomes more limiting such as this ecological site, shrub production, cover, density and diversity decline. When soil moisture becomes even more limiting, due to factors such as very low elevations and hot temperatures, absence of sheet-flow, restrictive surface cover such as desert pavement, or the presence of subsurface horizons that limit infiltration, the shrub community typically becomes even sparser, and is restricted to widely spaced, small creosote bush.

This ecological site tends to occupy distal fan positions, far from sources of run-on. Infrequent precipitation rapidly infiltrates these sandy soils, with minimal loss due to run-off and evaporation (Noy-Meir 1973, Austin et al. 2004), leaving very little water available at near surface depths. Deep, free-draining soils promote dominance by the deep-rooted, long-lived evergreen creosote bush, which can access water held at deep levels (McAuliffe 1994, Hamerlynk et al. 2002, Hamerlynk and McAuliffe 2008). Shallow-rooted species like burrobush (*Ambrosia dumosa*), which is co-dominant with creosote bush in soils or environments with greater moisture availability, cannot survive in the arid conditions of this site.

### Disturbance dynamics

The primary disturbances influencing this ecological site are drought, invasion by non-native annual plants, and fire, all of which interact. Drought is an important shaping force in Mojave Desert plant communities (Webb et al. 2003, Bowers 2005, Hereford et al. 2006, Miriti et al. 2007). Short-lived perennial shrubs and perennial grasses demonstrate the highest rates of mortality (Webb et al. 2003, Bowers 2005, Hereford et al. 2006, Miriti et al. 2007), and annual species remain dormant in the soil seedbank (Beatley 1969, 1974, 1976). Long-lived shrubs and trees are more likely to exhibit branch-pruning, and or limited recruitment during drought (e.g. Hereford et al. 2006, Miriti et al. 2007), leading to reduced cover and biomass in drought-afflicted communities.

Non-native annual species such as red brome (*Bromus rubens*), Mediterranean grass (*Schismus barbatus*), redstem stork's bill (*Erodium cicutarium*) and Asian mustard (*Brassica tournefortii*) have become naturalized throughout the Mojave Desert over the past century (Rickard and Beatley 1965, D'Antonio and Vitousek 1992,

Brooks 1999, Reid et al. 2006, Norton et al. 2007). In lower elevations, where soil temperature regimes are hyperthermic and soil moisture is more limiting, Mediterranean grass is the dominant non-native grass (Brooks and Berry 2006). Like native annuals, nonnative annual cover and production is directly related to winter precipitation (Beatley 1969, Brooks and Berry 2006, Barrows et al. 2009). The fine sand textures that are typical for this ecological site provide preferred habitat for Asian mustard, and this site has a high potential for invasion by this species during wet years.

Invasion by non-native annual grasses has increased the flammability of Mojave Desert vegetation communities by providing a continuous fine fuel layer between widely spaced shrubs (Brown and Minnich 1986, Brooks 1999, Brooks et al. 2004, Rao and Allen 2010, Rao et al. 2010). After fire, these communities appear to be more susceptible to invasion by exotic grasses, leading to a grass-fire cycle (D'Antonio and Vitousek 1992). The low potential for high productivity of annual species in this ecological site means that it is relatively resilient to fire. However, after years of extremely high winter precipitation, this site may burn (Brown and Minnich 1986, Brooks et al. 2007).

## **State and transition model**

## R030XD006CA Dry Deep Sandy Fan Aprons




Figure 4. R030XD006CA

### State 1 Historic State

State 1 represents the historic range of variability for this ecological site. This state no longer exists due to the ubiquitous naturalization of non-native species in the Mojave Desert. Periodic drought and rare fire were the natural disturbances influencing this ecological site. Fire would have been a very rare occurrence due to the lack of a

continuous fine fuel layer between shrubs. Data for this State does not exist, but dynamics and composition would have been similar to State 2, except with only native species present. See State 2 narrative for more detailed information.

## State 2

### Reference State

State 2 represents the current range of variability for this site. Non-native annuals, including Mediterranean grass, red-stem stork’s bill, and Asian mustard are naturalized in this plant community. Their abundance varies with precipitation, but they are at least sparsely present (as current year’s growth or present in the soil seedbank).

## Community 2.1

### Reference plant community



Figure 5. Community Phase 2.1

The reference plant community is maintained by periods of average climatic conditions and the absence of fire. It is dominated by creosote bush. There are no important secondary shrubs, though burrobrush (*Ambrosia dumosa*) or white ratany (*Krameria grayi*) are occasionally present at trace levels. The perennial bunchgrass big galleta may be present at very low levels. Native annual forbs are seasonally present, and common species include *Cryptantha* (*Cryptantha* spp.), smooth desert dandelion (*Malacothrix glabrata*), desert Indianwheat (*Plantago ovata*), and devil’s spineflower (*Chorizanthe rigida*). Mediterranean grass and redstem stork’s bill are typically present at low levels, and Asian mustard may be present at up to 5 percent cover.

Table 5. Annual production by plant type

| Plant Type      | Low<br>(Kg/Hectare) | Representative Value<br>(Kg/Hectare) | High<br>(Kg/Hectare) |
|-----------------|---------------------|--------------------------------------|----------------------|
| Shrub/Vine      | 22                  | 53                                   | 128                  |
| Forb            | 1                   | 22                                   | 84                   |
| Grass/Grasslike | –                   | –                                    | 2                    |
| <b>Total</b>    | <b>23</b>           | <b>75</b>                            | <b>214</b>           |

## Community 2.2

### Drought response

This community phase is characterized by an overall decline in cover due to branch-pruning and lack of recruitment of creosote bush, and lack of emergence of annual forbs and grasses. Creosote bush is an evergreen species capable of utilizing moisture at any time of the year. This ability buffers populations from the effects of drought that occur as the absence of the winter rains (the primary source of moisture for this ecological site). Further, creosote bush germinates in response to moisture during the warm season, so may still recruit if warm season rains occur during winter drought (Hereford et al. 2006). Creosote bush exhibits branch-pruning during severe drought, but mortality during drought in the Mojave Desert is very low (Webb et al. 2003, Hereford et al. 2006). Nevertheless,



during severe drought, creosote bush mortality may occur. This community is at reduced risk of burning, and if it is ignited, will experience lower severity, smaller fires because of reductions in annual and perennial biomass (Minnich 2003). However, drought immediately after a period of heavy moisture, results in standing biomass of native fuels that may carry a fire one year post-production (Minnich, 2003), and standing dead biomass of non-native annuals that may provide fuel for 2 -3 years post-fire (Minnich 2003; Rao et al. 2010).

## **Community 2.3**

### **Fire regeneration community**

This community phase is characterized by severe declines in creosote bush. Creosote bush is generally killed by fire, and is slow to re-colonize burned areas due to specific recruitment requirements (Brown and Minnich 1986, Brooks et al. 2007, Steers and Allen 2011). Creosote bush communities in the Mojave Desert may resemble the natural range of variation found in pre-fire conditions in terms of species composition in as little as nineteen years (Engel and Abella 2011), but creosote communities in the Colorado Desert may show little recovery after 30 years (Steers and Allen 2011). The timing and severity of fire, as well as post-fire climate conditions determines trajectories of recovery (Brown and Minnich 1986, Steers and Allen 2011). The post-burn community is dominated by non-native and native annual grasses and forbs, and native subshrubs. Native annuals likely to be present include smooth desert dandelion, pincushion flower, and cryptantha, but many different species could be at a particular site. Subshrubs that often become dominant after fire include desert globemallow (*Sphaeralcea ambigua*), desert trumpet (*Eriogonum inflatum*), and desert marigold (*Baileya multiradiata*). With time, shrub cover increases with colonization by short-lived shrubs from off-site dispersal. Given the extremely arid conditions of this site, burrobrush (*Hymenoclea salsola*) may be the only short-lived shrub capable of colonizing this site. With a long period of time without fire, creosote bush begins to regain dominance (Vasek 1983, Abella 2009, Vamstad 2009). This community is an at-risk phase, as the increased cover and biomass of non-native annual grasses increases the likelihood of repeat burning. Further, the low cover in this ecological site, and the low colonization potential following fire means that the burned community is extremely susceptible to wind erosion, especially if drought follows fire. If the fire return interval is less than 20 years, or if extreme drought or additional disturbance affect the fire regeneration community, this community is very likely to transition to State 3.

## **Pathway 2.1a**

### **Community 2.1 to 2.2**

This pathway occurs with prolonged or severe drought.

## **Pathway 2.1b**

### **Community 2.1 to 2.3**

This pathway occurs with moderate to severe fire.

## **Pathway 2.2a**

### **Community 2.2 to 2.1**

This pathway occurs with a return to average or above average precipitation.

## **Pathway 2.2b**

### **Community 2.2 to 2.3**

This pathway occurs with moderate to severe fire, and takes place within three years of a very wet period. At longer than three years of drought, the community is at low risk of burning.

## **Pathway 2.3a**

### **Community 2.3 to 2.1**

This pathway occurs with time and the absence of additional disturbance.

## **State 3**

**Repeated Fire**

This state develops when the fire return interval is less than 20 years, or the fire regeneration community in state 2 is affected by extreme drought or other disturbance (such as off-road vehicle use). This state has been significantly altered from the natural range of variability found in States 1 and 2. Creosote bush is lost, and non-native annual grasses and forbs dominate the community. This state is at high-risk of repeat burning due to high fine fuel cover. This community is also susceptible to wind and water erosion, due to the loss of shrub cover (Bull 1997). This can lead to arroyo development near ephemeral drainage channels. Furthermore, the loss of vegetation structure present in the historic and reference state decreases the suitability of this habitat for wildlife (Brooks et al. 2007, Vamstad 2009). Since rodent seed caching is important for the dispersal and establishment of many desert species this can further inhibit recovery.

**Community 3.1**  
**Repeat burn community**

The community is dominated by Mediterranean grass, red-stem stork’s bill, Asian mustard, and native forbs. Desert marigold, desert trumpet, and burrobrush may be abundant.

**Transition 1**  
**State 1 to 2**

This transition occurred with the naturalization of non-native species in this ecological site. Non-native species were introduced with settlement of the Mojave Desert region in the 1860s. Post-settlement cattle and sheep grazing, as well as dryland farming, helped to spread and facilitate their establishment (Brooks and Pyke 2000, Brooks et al. 2007).

**Transition 2**  
**State 2 to 3**

This transition occurs when the fire return interval is less than 20 years, or when extreme drought or additional disturbance affect the fire regeneration community.

**Restoration pathway 1**  
**State 3 to 2**

Restoration of communities severely altered by repeat fire at the landscape scale is difficult. Methods may include aerial seeding of early native colonizers such as desert globemallow, burrobrush, threeawns (*Aristida* spp.), and desert marigold. Increased native cover may help to reduce non-native plant invasion, helps to stabilize soils, provides a source of food and cover for wildlife, including desert tortoise (*Gopherus agassizii*), and provides microsites that facilitate creosote bush establishment. However, the amount of seed required for success is often prohibitive. Large-scale planting of both early colonizers and community dominants tends to be more successful in terms of plant survival, especially if outplants receive supplemental watering during the first two years. Creosote bush can be successfully propagated and outplanted. Pre-emergent herbicides (Plateau) have been used in the year immediately post-fire to attempt to inhibit or reduce brome invasion. How successful this is on a landscape scale, and the non-target effects have not yet been determined.

**Additional community tables**

Table 6. Community 2.1 plant community composition

| Group                  | Common Name                      | Symbol | Scientific Name              | Annual Production (Kg/Hectare) | Foliar Cover (%) |
|------------------------|----------------------------------|--------|------------------------------|--------------------------------|------------------|
| <b>Shrub/Vine</b>      |                                  |        |                              |                                |                  |
| 1                      | <b>Native Shrubs</b>             |        |                              | 11–82                          |                  |
|                        | creosote bush                    | LATR2  | <i>Larrea tridentata</i>     | 11–78                          | 2–7              |
|                        | burrobush                        | AMDU2  | <i>Ambrosia dumosa</i>       | 0–2                            | 0–1              |
|                        | white ratany                     | KRGR   | <i>Krameria grayi</i>        | 0–1                            | 0–1              |
| <b>Forb</b>            |                                  |        |                              |                                |                  |
| 2                      | <b>Native annual forbs</b>       |        |                              | 0–28                           |                  |
|                        | cryptantha                       | CRYPT  | <i>Cryptantha</i>            | 0–22                           | 0                |
|                        | desert Indianwheat               | PLOV   | <i>Plantago ovata</i>        | 0–7                            | 0–12             |
|                        | smooth desertdandelion           | MAGL3  | <i>Malacothrix glabrata</i>  | 0–6                            | 0–3              |
|                        | devil's spineflower              | CHRI   | <i>Chorizanthe rigida</i>    | 0–4                            | 0                |
|                        | Mojave desertstar                | MOBE2  | <i>Monoptilon bellioides</i> | 0–1                            | 0                |
| 5                      | <b>Non-native annual forbs</b>   |        |                              | 0–67                           |                  |
|                        | Asian mustard                    | BRT0   | <i>Brassica tournefortii</i> | 0–67                           | 0–5              |
| <b>Grass/Grasslike</b> |                                  |        |                              |                                |                  |
| 3                      | <b>Non-native annual grasses</b> |        |                              | 0–2                            |                  |
|                        | Mediterranean grass              | SCHIS  | <i>Schismus</i>              | 0–2                            | 0–1              |
| 4                      | <b>Perennial Grass</b>           |        |                              | 0–1                            |                  |
|                        | big galleta                      | PLRI3  | <i>Pleuraphis rigida</i>     | 0–1                            | 0                |

## Animal community

This ecological site provides critical habitat for the threatened desert tortoise (*Gopherus agassizii agassizii*). Creosote bush shrublands provides a home for an abundance of specialist insect species, for example, creosote bush flowers provide nutrition for over twenty species of bees, and the creosote bush grasshopper (*Boottettix argentatus*) feeds solely on creosote leaves (Pavlik 2008).

## Recreational uses

This site is highly valued for open space and those interested in desert ecology. Uses include mountain biking, hiking, bird watching and botanizing. Desert tortoise and annual wildflowers may also attract visitors during the spring.

## Other products

Creosote bush is an important medicinal plant for Native Americans. It has a very wide range of uses from treatment for consumption, bowl complaints, and menstrual cramps, to induce vomiting, relief for arthritis, rheumatism, aching bones and sprains, congestion and cold, as an antiseptic and disinfectant, dandruff, antispasmodic, to induce urination, gonorrhea, and to cancer treatment. (This list is not exhaustive).

<http://herb.umd.umich.edu/herb/search.pl?searchstring=Larrea+tridentata>.

Creosote bush stems are used to make weapons, digging tools, and basket handles, and creosote gum is used for knife and awl handles. Creosote bush branches are used as thatch in dwelling construction.

<http://herb.umd.umich.edu/herb/search.pl?searchstring=Larrea+tridentata>.

## Inventory data references

The following NRCS plots were used to describe this ecological site:

D1-M (Type location)

1248014701

J5-K

MW-D

PFPR-3

PW-D

## Type locality

|                                  |  |
|----------------------------------|--|
| Location 1: Riverside County, CA |  |
| UTM zone                         | N  |
| UTM northing                     | 3774342  |
| UTM easting                      | 623922   |
| General legal description        | The type location is approximately 1.65 miles east of the intersection of Highway 62 and Ironage Road, 0.12 mile south of Highway 62 on the border of Joshua Tree National Park. |

## Other references

Abella, S. R. 2009. Post-fire plant recovery in the Mojave and Sonoran Deserts of western North America. *Journal of Arid Environments* 73:699-707.

Austin, A. T., L. Yahdjian, J. M. Stark, J. Belnap, A. Porporato, U. Norton, D. A. Ravetta, and S. M. Scheaeffer. 2004. Water pulses and biogeochemical cycles in arid and semiarid ecosystems. *Oecologia* 141:221-235.

Barrows, C. W., E. B. Allen, M. L. Brooks, and M. F. Allen. 2009. Effects of an invasive plant on a desert sand dune landscape. *Biological Invasions* 11:673-686.

Beatley, J. C. 1969. Dependence of desert rodents on winter annuals and precipitation. *Ecology* 50:721-724.

Beatley, J. C. 1974. Effects of rainfall and temperature on the distribution and behavior of *Larrea tridentata* (Creosote-bush) in the Mojave Desert of Nevada. *Ecology* 55:245-261.

Beatley, J. C. 1976. Rainfall and fluctuating plant populations in relation to distributions and numbers of desert rodents in southern Nevada. *Oecologia* 24:21-42.

Bedford, D. R., D. M. Miller, K. M. Schmidt, and G. A. Phelps. 2009. Landscape-scale relationships between surficial geology, soil texture, topography, and creosote bush size and density in the eastern Mojave Desert of California. Pages 252-277 in R. H. Webb, L. F. Fenstermaker, J. S. Heaton, D. L. Hughson, E. V. McDonald, and D. H. Miller, editors. *The Mojave Desert: ecosystem processes and sustainability*. University of Nevada Press, Reno, NV.

Bowers, J. E. 2005. Effects of drought on shrub survival and longevity in the northern Sonoran Desert. *Journal of the Torrey Botanical Society* 132:421-431.

Brooks, M. L. 1999. Habitat invasibility and dominance by alien annual plants in the western Mojave Desert. *Biological Invasions* 1:325-337.

Brooks, M. L. and K. H. Berry. 2006. Dominance and environmental correlates of alien annual plants in the Mojave Desert, USA. *Journal of Arid Environments* 67:100-124.

Brooks, M. L., C. M. D'Antonio, D. M. Richardson, J. B. Grace, J. E. Keeley, J. M. DiTomaso, R. J. Hobbs, M. Pellant, and D. Pyke. 2004. Effects of invasive alien plants on fire regimes. *Bioscience* 54:677-689.

Brooks, M. L., T. C. Esque, and T. Duck. 2007. Creosotebush, blackbrush, and interior chaparral shrublands. *RMRS-GTR-202*.

Brooks, M. L. and D. A. Pyke. 2000. Invasive plants and fire in the deserts of North America. Pages 1-14 in *Fire*

conference 2000: the first national congress on fire ecology, prevention, and management. Tall Timbers Research Station, Tallahassee, FL.

Brown, D. E. and R. A. Minnich. 1986. Fire and Changes in Creosote Bush Scrub of the Western Sonoran Desert, California. *American Midland Naturalist* 116:411-422.

Bull, W. B. 1997. Discontinuous ephemeral streams. *Geomorphology* 19:227-276.

D'Antonio, C. M. and P. M. Vitousek. 1992. Biological invasions by exotic grasses, the grass/fire cycle, and global change. *Annual Review of Ecology and Systematics* 23:63-87.

Engel, E. C. and S. R. Abella. 2011. Vegetation recovery in a desert landscape after wildfires: influences of community type, time since fire and contingency effects. *Journal of Applied Ecology* 48:1401-1410.

Hamerlynk, E. P. and J. R. McAuliffe. 2008. Soil-dependent canopy die-back and plant mortality in two Mojave Desert shrubs. *Journal of Arid Environments* 72:1793-1802.

Hamerlynk, E. P., J. R. McAuliffe, E. V. McDonald, and S. D. Smith. 2002. Ecological responses of two Mojave desert shrubs to soil horizon development and soil water dynamics. *Ecology* 83:768-779.

Hamerlynk, E. P., J. R. McAuliffe, and S. D. Smith. 2000. Effects of surface and sub-surface soil horizons on the seasonal performance of *Larrea tridentata* (creosotebush). *Functional Ecology* 14:596-606.

Hereford, R., R. H. Webb, and C. I. Longpre. 2006. Precipitation history and ecosystem response to multidecadal precipitation variability in the Mojave Desert region, 1893-2001. *Journal of Arid Environments* 67:13-34.

Holland, R. F. 1986. Preliminary descriptions of the terrestrial natural communities of California. State of California Department of Fish and Game, Sacramento, CA.

Martre, P., G. B. North, E. G. Bobich, and P. S. Nobel. 2002. Root deployment and shoot growth for two desert species in response to soil rockiness. *American Journal of Botany* 89:1933-1939.

McAuliffe, J. R. 1994. Landscape evolution, soil formation, and ecological patterns and processes in Sonoran Desert bajadas. *Ecological Monographs* 64:112-148.

Minnich, R. A. 2003. Fire and dynamics of temperate desert woodlands in Joshua Tree National Park. Contract, Joshua Tree National Park.

Miriti, M. N., S. Rodriguez-Buritica, S. J. Wright, and H. F. Howe. 2007. Episodic death across species of desert shrubs. *Ecology* 88:32-36.

Norton, J. B., T. A. Monaco, and U. Norton. 2007. Mediterranean annual grasses in western North America: kids in a candy store. *Plant Soil* 298:1-5.

Noy-Meir, I. 1973. Desert ecosystems: environment and producers. *Annual Review of Ecology and Systematics* 4:25-51.

Pavlik, B. M. 2008. The California Deserts: an ecological rediscovery. University of California Press, Ltd., Berkeley and Los Angeles, California.

Rao, L. E. and E. B. Allen. 2010. Combined effects of precipitation and nitrogen deposition on native and invasive winter annual production in California deserts. *Oecologia* 162:1035-1046.

Rao, L. E., E. B. Allen, and T. M. Meixner. 2010. Risk-based determination of critical nitrogen deposition loads for fire spread in southern California deserts. *Ecological Applications* 20:1320-1335.

Reid, C. R., S. Goodrich, and J. E. Bowns. 2006. Cheatgrass and red brome: history and biology of two invaders. Pages 27-32 in *Shrublands under fire: disturbance and recovery in a changing world*. U.S. Department of

Agriculture, Forest Service, Rocky Mountain Research Station, Cedar City, Utah.

Rickard, W. H. and J. C. Beatley. 1965. Canopy-coverage of the desert shrub vegetation mosaic of the Nevada test site. *Ecology* 46:524-529.

Rundel, P. W. and A. C. Gibson. 1996. *Ecological communities and processes in a Mojave Desert Ecosystem: Rock Valley Nevada*. Cambridge University Press, Cambridge, England.

Sawyer, J. O., T. Keeler-Woelf, and J. M. Evans. 2009. *A manual of California vegetation*. 2nd edition. California Native Plant Society, Sacramento, California.

Steers, R. J. and E. B. Allen. 2011. Fire effects on perennial vegetation in the western Colorado Desert, USA. *Fire Ecology* 7:59-74.

Vamstad, M. S. 2009. Effects of fire on vegetation and small mammal communities in a Mojave Desert Joshua tree woodland. M.S. University of California, Riverside, Riverside, Ca.

Vasek, F. C. 1983. Plant succession in the Mojave Desert. *Crossosoma* 9:1-23.

Webb, R. H., M. B. Muroy, T. C. Esque, D. E. Boyer, L. A. DeFalco, D. F. Haines, D. Oldershaw, S. J. Scoles, K. A. Thomas, J. B. Blainey, and P. A. Medica. 2003. Perennial vegetation data from permanent plots on the Nevada Test Site, Nye County, Nevada. U.S. Geological Society, Tucson, AZ.

## Contributors

Alice Lee Miller

P. Novak-Echenique / H. McAllister

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

|   |                   |
|---|-------------------|
| Author(s)/participant(s)                    |                   |
| Contact for lead author                     |                   |
| Date  |                   |
| Approved by                                 |                   |
| Approval date                               |                   |
| Composition (Indicators 10 and 12) based on | Annual Production |

## Indicators

### 1. Number and extent of rills:

---

### 2. Presence of water flow patterns:

---

3. **Number and height of erosional pedestals or terracettes:**

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

---

5. **Number of gullies and erosion associated with gullies:**

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

---

7. **Amount of litter movement (describe size and distance expected to travel):**

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

---

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

---

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

---

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

---

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

---

14. **Average percent litter cover (%) and depth ( in):**

---

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

---

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

---

17. **Perennial plant reproductive capability:**

---