

Ecological site R032XY244WY Saline Upland (SU) 5-9" Wind River Basin Precipitation Zone

Accessed: 04/30/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

Associated sites

R032XY218WY	Impervious Clay (IC) 5-9" Wind River Basin Precipitation Zone
R032XY222WY	Loamy (Ly) 5-9" Wind River Basin Precipitation Zone
R032XY240WY	Saline Lowland Drained (SLDr) 5-9" Wind River Basin Precipitation Zone

Similar sites

R032XY344WY	Saline Upland (SU) 10-14" East Precipitation Zone Saline Upland 10-14" East P.Z. has higher production.
-------------	---

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

This site occurs on nearly level to moderately sloping land.

Table 2. Representative physiographic features

Landforms	(1) Hill (2) Alluvial fan (3) Stream terrace
Flooding duration	Extremely brief (0.1 to 4 hours) to brief (2 to 7 days)
Flooding frequency	Very rare to rare
Ponding frequency	None
Elevation	1,372–2,012 m
Slope	0–15%
Ponding depth	0 cm
Aspect	Aspect is not a significant factor

Climatic features

Annual precipitation ranges from 5-9 inches per year. The normal precipitation pattern shows peaks in May and June and a secondary peak in September. This amounts to about 50% of the mean annual precipitation. Much of the moisture that falls in the latter part of the summer is lost by evaporation and much of the moisture that falls during the winter is lost by sublimation. Average snowfall is about 20 inches annually. Wide fluctuations may occur in yearly precipitation and result in more dry years than those with more than normal precipitation.

Temperatures show a wide range between summer and winter and between daily maximums and minimums, due to the high elevation and dry air, which permits rapid incoming and outgoing radiation. Cold air outbreaks from Canada in winter move rapidly from northwest to southeast and account for extreme minimum temperatures. Chinook winds may occur in winter and bring rapid rises in temperature. Extreme storms may occur during the winter, but most severely affect ranch operations during late winter and spring.

High winds are generally blocked from the basin by high mountains, but can occur in conjunction with an occasional thunderstorm.

Growth of native cool-season plants begins about April 1 and continues to about July 1. Cool weather and moisture in September may produce some green up of cool season plants that will continue to late October.

The following information is from the "Pavillion" climate station:

Minimum Maximum 5 yrs. out of 10 between
 Frost-free period (days): 95 175 May 19 – September 19
 Freeze-free period (days): 98 185 May 6 – October 3
 Mean Annual Precipitation (inches): 2.50 12.54

Mean annual precipitation: 7.85 inches

Mean annual air temperature: 44.53 F (30.5 F Avg. Min. to 58.5 F Avg. Max.)

For detailed information visit the Natural Resources Conservation Service National Water and Climate Center at <http://www.wcc.nrcs.usda.gov/> website. Other climate station(s) representative of this precipitation zone include "Riverton", "Arminto", and "Lost Cabin".

Table 3. Representative climatic features

Frost-free period (average)	175 days
Freeze-free period (average)	185 days
Precipitation total (average)	229 mm

Influencing water features

Stream Type: None

Soil features

The soils of this site are moderately deep (greater than 20" to bedrock) to very deep well-drained soils formed in alluvium from sodic or alkaline materials. These soils have moderate to slow permeability and are moderately to strongly saline and/or alkaline. The surface soil will vary from 2 to 6 inches in thickness. Some soils may contain more soluble salts in the subsoils than in the surface soils. The soil characteristic having the most influence on the plant community is the high quantity of soluble salts.

Table 4. Representative soil features

Surface texture	(1) Loam (2) Very fine sandy loam (3) Fine sandy loam
Family particle size	(1) Clayey
Drainage class	Well drained
Permeability class	Slow to moderate
Soil depth	51–152 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	3.56–16 cm
Calcium carbonate equivalent (0-101.6cm)	0–15%
Electrical conductivity (0-101.6cm)	4–16 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	8–16
Soil reaction (1:1 water) (0-101.6cm)	7.4–11
Subsurface fragment volume <=3" (Depth not specified)	0%
Subsurface fragment volume >3" (Depth not specified)	0%

Ecological dynamics

Potential vegetation on this site is dominated by salt tolerant plants and drought resistant mid cool-season perennial grasses. The expected potential composition for this site is about 40% grasses, 10% forbs and 50% woody plants. The composition and production will vary naturally due to historical use, fluctuating precipitation and fire frequency.

As this site deteriorates, species such as birdfoot sagebrush and greasewood will increase. Weedy annuals will invade. Cool season grasses such as Indian ricegrass, bottlebrush squirreltail, and rhizomatous wheatgrasses will decrease in frequency and production.

The Historic Climax Plant Community (description follows the plant community diagram) has been determined by study of rangeland relic areas, or areas protected from excessive disturbance. Trends in plant communities going from heavily grazed areas to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.


The following is a State and Transition Model Diagram that illustrates the common plant communities (states) that can occur on the site and the transitions between these communities. The ecological processes will be discussed in

more detail in the plant community narratives following the diagram.

Plant Community Narratives

Following are the narratives for each of the described plant communities. These plant communities may not represent every possibility, but they probably are the most prevalent and repeatable plant communities. The plant composition tables shown above have been developed from the best available knowledge at the time of this revision. As more data is collected, some of these plant communities may be revised or removed, and new ones may be added. None of these plant communities should necessarily be thought of as “Desired Plant Communities”. According to the USDA NRCS National Range and Pasture Handbook, Desired Plant Communities (DPC’s) will be determined by the decision-makers and will meet minimum quality criteria established by the NRCS. The main purpose for including any description of a plant community here is to capture the current knowledge and experience at the time of this revision.

State and transition model



BM - Brush Management (fire, chemical, mechanical)

Freq. & Severe Grazing - Frequent and Severe Utilization of the Cool-season Mid-grasses during the Growing Season

GLMT - Grazing Land Mechanical Treatment

LTPG - Long-term Prescribed Grazing

MCSLG - Moderate, Continuous Season-long Grazing

NU, NF - No Use and No Fire

PG - Prescribed Grazing (proper stocking rates with adequate recovery periods during the growing season)

VLTPG - Very Long-term Prescribed Grazing (could possibly take generations)

WF - Wildfire

State 1

Gardner's saltbush/ Indain Ricegrass/ Bottlebrush Squirreltail

Community 1.1

Gardner's saltbush/ Indain Ricegrass/ Bottlebrush Squirreltail

The interpretive plant community for this site is the Historic Climax Plant Community. This state evolved with grazing by large herbivores and droughty saline and/or alkali soils. This plant community can be found on areas that are properly managed with grazing and on areas receiving short periods of rest. Potential vegetation is about 40% grasses or grass-like plants, 10% forbs, and 50% woody plants. Gardner's saltbush dominates this state. Other salt tolerant shrubs include greasewood and birdfoot sagebrush. The major grasses include Indian ricegrass, bottlebrush squirreltail, and rhizomatous wheatgrasses. Other grasses occurring in this state include Sandberg bluegrass and needleandthread. A variety of forbs also occurs in this state and plant diversity is high (see Plant Composition Table). The total annual production (air-dry weight) of this state is about 300 pounds per acre, but it can range from about 200 lbs./acre in unfavorable years to about 400 lbs./acre in above average years. This state is fragile, but well adapted to the Northern Great Plains climatic conditions. The diversity in plant species allows for high drought tolerance. This is a sustainable plant community, but is difficult to reestablish when damaged. (Site/soil stability, watershed function, and biologic integrity). Transitions or pathways leading to other plant communities are as follows: • Moderate, Continuous Season-Long grazing will convert this plant community to the Gardner Saltbush/Rhizomatous Wheatgrasses Plant Community. Prolonged Drought will exacerbate this transition.

Figure 4. Plant community growth curve (percent production by month). WY0801, 5-9WR upland sites.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
			10	50	25	5		10			

State 2

Gardner's saltbush/ Rhizomatous Wheatgrasses

Community 2.1

Gardner's saltbush/ Rhizomatous Wheatgrasses

Historically, this plant community evolved under grazing by large ungulates. Currently this vegetation state is found under moderate, season-long grazing by livestock. Prolonged drought can also play an important role and will exacerbate these conditions. Gardner's saltbush and rhizomatous wheatgrasses are major components of this plant community. Cool-season grasses make up the majority of the understory with the balance made up of short warm-season grasses, annual cool-season grass, and miscellaneous forbs. Dominant grasses include rhizomatous wheatgrasses, bottlebrush squirreltail, Sandberg bluegrass, and blue grama. Forbs commonly found in this plant community include Smooth woodyaster, Cous biscuitroot, Wild onion, and Leafy wildparsely. Plains pricklypear and winterfat can also occur. When compared to the Historic Climax Plant Community, birdfoot sagebrush and blue grama have increased. Plains pricklypear cactus will also have increased, but occurs only in small patches. Indian ricegrass has decreased and may occur in only trace amounts within the patches of pricklypear. In addition, winterfat may or may not have changed depending on the season of use. The total annual production (air-dry weight) of this state is about 240 pounds per acre, but it can range from about 150 lbs./acre in unfavorable years to about 350 lbs./acre in above average years. This plant community is relatively resistant to change. The herbaceous species present are well adapted to grazing; however, species composition can be altered through long-term overgrazing. The herbaceous component is mostly intact and plant vigor and replacement capabilities are sufficient. Water flow patterns and litter movement may occur, but is not extensive. Incidence of pedestalling is minimal. Soils are mostly stable and the surface shows minimum soil loss. The watershed is functioning and the biotic community is intact. Transitional pathways leading to other plant communities are as follows: • Prescribed grazing will prevent further deterioration and over the long-term may return this state to near Historic Climax Plant Community Vegetation State. • Frequent and severe grazing with the absences of fire on less sodic soils will convert this plant community to a Mixed Shrub/Blue Grama Sod Vegetation State. • Frequent and severe grazing plus no fire on more sodic soils will convert this state to Gardner Saltbush/Bare Ground Vegetative State.

Figure 5. Plant community growth curve (percent production by month). WY0801, 5-9WR upland sites.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
			10	50	25	5		10			

State 3

Mixed Shrub/ Blue Grama sod

Community 3.1

Mixed Shrub/ Blue Grama sod

This plant community evolved under frequent and severe grazing on less sodic soils. Gardner's saltbush, birdfoot sagebrush and blue grama are the dominant species of this plant community. A dense sod of blue grama can be common but mostly occurs in large mosaic patterns. Cool-season grasses have been mostly eliminated and if still present can only be found within patches of plains pricklypear. The dominant grass is blue grama but Sandberg bluegrass can be present. Cactus often has increased. Noxious weeds such as Russian knapweed and halogeton may invade the site. When compared with the HCPC or the Gardner's Saltbush/Rhizomatous Wheatgrass Plant Communities, the annual production is lower as is the production of Gardner's saltbush and the perennial cool-season grasses. The total annual production (air-dry weight) of this state is about 150 pounds per acre, but it can range from about 25 lbs./acre in unfavorable years to about 200 lbs./acre in above average years. This plant community is resistant to change as the stand of blue grama is established and continued frequent and severe grazing or the removal of grazing does not seem to affect the plant composition or structure of the plant community. The biotic integrity of this state is minimally functional and plant diversity is moderate to low. Plant vigor is weakened and replacement capabilities are limited due to the reduced number of cool-season grasses. This state is stable and protected from excessive erosion. The sod formed by these grasses is resistant to water infiltration. While the soil is protected by this sod, excessive runoff may occur off-site. As a result, rills or other more severe erosion can occur on the adjoining sites. The watershed may or may not be functioning, as runoff may affect adjoining sites. The biotic integrity of this plant community is not intact. Transitions or pathways leading to other plant communities are as follows: • Grazing land mechanical treatment (chiseling and seeding, etc.) followed by prescribed grazing (and BM if needed) will return this plant community to near Historic Climax Plant Community.

Figure 6. Plant community growth curve (percent production by month). WY0801, 5-9WR upland sites.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
			10	50	25	5		10			

State 4

Gardners Saltbush/ Bare Ground

Community 4.1

Gardners Saltbush/ Bare Ground

This plant community can occur where sites are subjected to continuous yearlong grazing and where soils are highly sodic. Gardner's saltbush dominates this site and in some cases comprises almost 100% of the plant community. The interspaces between plants have expanded significantly leaving the amount of bare ground prevalent and the soil surface exposed to erosive elements. Cool season grasses have been eliminated or greatly reduced. Noxious weeds such as Russian knapweed and halogeton have likely invaded the large openings and can dominate this site. When compared to the HCPC, plant production is greatly diminished due to the excessive amount of bare ground. The total annual production (air-dry weight) of this state is about 90 pounds per acre, but it can range from about 50 lbs./acre in unfavorable years to about 150 lbs./acre in above average years. This plant community is resistant to change as the stand becomes more decadent. These areas may actually be more resistant to fire as less fine fuels are available and the bare ground between the brush plants is increased. Continued frequent and severe grazing or the removal of grazing does not seem to affect the plant composition or structure of the plant community. Plant diversity is extremely low. The plant vigor is diminished and replacement capabilities are severely reduced due to the decrease in the number of cool-season grasses. Plant litter is noticeably less when compared to the HCPC. Soil erosion is accelerated because of increased bare ground. Water flow patterns and pedestalling are obvious. Infiltration is reduced and runoff is increased. Rill channels may be

noticeable in the interspaces and gullies may be establishing where rills have concentrated down slope. Transitional pathways leading to other plant communities are as follows: • Grazing land mechanical treatment (seeding, etc.) followed by prescribed grazing (and BM if needed), will return this plant community at or near the HCPC.

**Figure 7. Plant community growth curve (percent production by month).
WY0801, 5-9WR upland sites.**

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
			10	50	25	5		10			

Additional community tables

Table 5. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1				34–84	
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	34–84	–
2				34–84	
	squirreltail	ELELE	<i>Elymus elymoides ssp. elymoides</i>	34–84	–
3				34–50	
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	34–50	–
4				17–34	
	Grass, perennial	2GP	<i>Grass, perennial</i>	0–17	–
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	0–17	–
	needle and thread	HECO26	<i>Hesperostipa comata</i>	0–17	–
	Sandberg bluegrass	POSE	<i>Poa secunda</i>	0–17	–
Forb					
5				17–34	
	Forb, perennial	2FP	<i>Forb, perennial</i>	0–17	–
	textile onion	ALTE	<i>Allium textile</i>	0–17	–
	Missouri milkvetch	ASMI10	<i>Astragalus missouriensis</i>	0–17	–
	sulphur-flower buckwheat	ERUM	<i>Eriogonum umbellatum</i>	0–17	–
	cous biscuitroot	LOCO4	<i>Lomatium cous</i>	0–17	–
	leafy wildparsley	MUDI	<i>Musineon divaricatum</i>	0–17	–
	plains pricklypear	OPPO	<i>Opuntia polyacantha</i>	0–17	–
	western aster	SYAS3	<i>Symphyotrichum ascendens</i>	0–17	–
	yellow salsify	TRDU	<i>Tragopogon dubius</i>	0–17	–
	smooth woodyaster	XYGL	<i>Xylorhiza glabriuscula</i>	0–17	–
Shrub/Vine					
6				67–135	
	Gardner's saltbush	ATGA	<i>Atriplex gardneri</i>	67–135	–
7				17–34	
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	0–17	–
	birdfoot sagebrush	ARPE6	<i>Artemisia pedatifida</i>	0–17	–
	Wyoming big sagebrush	ARTRW8	<i>Artemisia tridentata ssp. wyomingensis</i>	0–17	–
	winterfat	KRASC	<i>Krascheninnikovia</i>	0–17	–
	greasewood	SAVE4	<i>Sarcobatus vermiculatus</i>	0–17	–

Animal community

Animal Community – Wildlife Interpretations

Historic Climax Plant Community: The predominance of woody plants in this plant community provides winter grazing for mixed-feeders, such as elk, and antelope. Suitable thermal and escape cover for these animals are limited due to the low quantities of tall woody plants. When found adjacent to sagebrush-dominated states, this plant community may provide lek sites for sage grouse. Other birds that would frequent this plant community include western meadowlarks, horned larks, and golden eagles. Some grassland obligate small mammals would occur here.

Gardner's Saltbush/Rhizomatous Wheatgrasses Plant Community: The combination of shrubs, grasses, and forbs can provide a forage source for large animals. Suitable thermal and escape cover for these animals are limited due to the low quantities of tall woody plants. When found adjacent to sagebrush-dominated states, this plant community may provide lek sites for sage grouse. Other birds that would frequent this plant community include western meadowlarks, horned larks, and golden eagles. Some grassland obligate small mammals would occur here.

Mixed Shrub/Blue Grama Sod Plant Community: These communities provide limited foraging for antelope and other grazers. This community may be used as a foraging site by sage grouse if proximal to woody cover. Generally, these are not target plant communities for wildlife habitat management.

Gardner's Saltbush/Bare Ground Plant Community: This plant community exhibits a low level of plant species diversity. It may provide some forage value for antelope, but in most cases it is not a desirable plant community to select as a wildlife habitat management objective.

Animal Community – Grazing Interpretations

The following table lists suggested stocking rates for cattle under continuous season-long grazing under normal growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of the conservation planning process. Often, the current plant composition does not entirely match any particular plant community (as described in this ecological site description). Because of this, a field visit is recommended, in all cases, to document plant composition and production. More precise carrying capacity estimates should eventually be calculated using this information along with animal preference data, particularly when grazers other than cattle are involved. Under more intensive grazing management, improved harvest efficiencies can result in an increased carrying capacity. If distribution problems occur, stocking rates must be reduced to maintain plant health and vigor.

Plant Community Production Carrying Capacity*

(lb./ac) (AUM/ac)

Historic Climax Plant Community 200-400 .10

Gardner's Saltbush/Rhizomatous Wheatgrasses 150-350 .08

Mixed Shrub/Blue Grama Sod 75-200 .05

Gardner's Saltbush/Bare Ground Plant Community 25-150 .03

* - Continuous, season-long grazing by cattle under average growing conditions.

Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage for cattle, sheep, or horses. During the dormant period, the forage for livestock use needs to be supplemented with protein because the quality does not meet minimum livestock requirements.

Hydrological functions

Water and salinity are the principal factors limiting forage production on this site. This site is dominated by soils in hydrologic group B and C, with localized areas in hydrologic group D. Infiltration ranges from slow to moderate. Runoff potential for this site varies from moderate to high depending on soil hydrologic group and ground cover. In many cases, areas with greater than 75% ground cover have the greatest potential for high infiltration and lower runoff. An example of an exception would be where short-grasses form a strong sod and dominate the site. Areas where ground cover is less than 50% have the greatest potential to have reduced infiltration and higher runoff (refer to Part 630, NRCS National Engineering Handbook for detailed hydrology information).

Rills and gullies should not typically be present. Water flow patterns should be barely distinguishable if at all present. Pedestals are only slightly present in association with bunchgrasses. Litter typically falls in place, and signs of movement are not common. Chemical and physical crusts may be present. Cryptogamic crusts are present, but only cover 1-2% of the soil surface.

Recreational uses

This site provides some hunting opportunities for upland game species.

Wood products

No appreciable wood products are present on the site.

Other products

None noted.

Inventory data references

Information presented here has been derived from NRCS inventory data. Field observations from range trained personnel were also used. Those involved in developing this site include: Chris Krassin, Range Management Specialist, NRCS and Everet Bainter, Range Management Specialist, NRCS. Other sources used as references include USDA NRCS Water and Climate Center, USDA NRCS National Range and Pasture Handbook, USDI and USDA Interpreting Indicators of Rangeland Health Version 3, and USDA NRCS Soil Surveys from various counties.

Contributors

C. Krassin

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	07/01/2005
Approved by	E. Bainter
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** Rills should not be present
-

2. **Presence of water flow patterns:** Barely observable
-

3. **Number and height of erosional pedestals or terracettes:** Essentially non-existent
-

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** Bare ground is 30-40%
-

5. **Number of gullies and erosion associated with gullies:** Active gullies should not be present

-
6. **Extent of wind scoured, blowouts and/or depositional areas:** None
-
7. **Amount of litter movement (describe size and distance expected to travel):** Little to no plant litter movement. Plant litter remains in place and is not moved by erosional forces.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Plant cover and litter is at 50% or greater of soil surface and maintains soil surface integrity. Soil Stability class is anticipated to be 4 or greater.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Use Soil Series description for depth and color of A-horizon
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Grass canopy and basal cover should reduce raindrop impact and slow overland flow providing increased time for infiltration to occur. Healthy deep rooted native grasses enhance infiltration and reduce runoff. Infiltration is slow to moderate.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** No compaction layer or soil surface crusting should be present.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional: Mid stature Grasses > Shrubs > Forbs > Short stature Grasses
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Very Low
-
14. **Average percent litter cover (%) and depth (in):** Average litter cover is 10-15% with depths of 0.1 to 0.25 inches
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 300 lbs/ac

-
16. **Potential invasive (including noxious) species (native and non-native).** List species which **BOTH** characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is **NOT** expected in the reference state for the ecological site: Birdfoot sagebrush, greasewood, Unpalatable forbs, Annuals, Exotics, and Species found on Noxious Weed List
-

17. **Perennial plant reproductive capability:** All species are capable of reproducing
-