

Ecological site R034AA013UT Semi-Wet Fresh Meadow

Last updated: 9/07/2023
Accessed: 05/18/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 034A–Cool Central Desertic Basins and Plateaus

Major Land Resource Area (MLRA) 34A, Cool Central Desertic Basins and Plateaus, consists of approximately 21 million acres in Wyoming, Colorado and Utah, it consists of 11 Land Resource Units (LRU). These units are divisions of the MLRA based on geology, landscape, common soils, water resources and plant community potentials. The elevation spans from approximately 5600 feet (1700 m) along the Green River in UT and CO to approximately 9500 feet (2900 m) near Jeffrey City, WY. Annual precipitation ranges from 7 to 16 inches (177 to 406 mm), with the driest areas in the Green River and Great Divide Basins and the wettest areas in northern Carbon County, southeast Fremont County and Albany County. There is a seasonal weather pattern that trends west to east, with more winter precipitation in the west and more spring and summer in the east, illustrated by diminishing amounts of big sagebrush in the eastern part of the MLRA.

LRU notes

The Bear River Valley LRU is located on the far western side of MLRA 34A between the Bear River Divide and the Monte Cristo Range, from Woodruff, Utah at the southern end to Cokeville, Wyoming at the northern end. The total area of the LRU is approximately 340,000 acres. It shares a boundary with MLRA 47, 43B and 46 (proposed). This LRU differs from the others in its geology, which is comprised mostly of alluvium and colluvium from the Stump Formation. Its weather patterns are such that the soil moisture is xeric, there is a slight peak in winter precipitation in this LRU, with typical yearly precipitation between 9 to 15 inches (230 to 380 mm). The soil temperature regime of this LRU is frigid with mean annual soil temperatures ranging from 44 to 48 degrees Fahrenheit (6.7 to 8.8°C). The elevation range is from 5700 to 7000 feet (1730 to 2130 m). The soils in the Bear River Valley are dominated by young aged very deep soils developed from sandstone and shale parent material re-worked with recent alluvium. Soils are dominated by Alfisols with young argillic horizons and by Fluvents in more recent alluvium. The Bear River runs through this LRU, allowing for ample amounts of irrigation water used in the lowland areas to produce hay. Smaller tributaries originating from the neighboring mountains.

Ecological site concept

The Semi-wet Fresh Meadow ecological site is primarily fed by groundwater from springs. The ecological site has very narrow valley bottoms and a small contributing watershed area. The upland areas adjacent to this site are typically big sagebrush (*Artemisia tridentata*). The streams in this ecological site serve only to convey water away from the spring area. The stream is perennial when near the source (springs), and may become intermittent as move further downstream of the spring. Streams that are perennial where significant springs supply the system, may gradually become intermittent when spring influence ceases. This ecological site in a typical precipitation year does not experience high flows in the spring in association with snowmelt because of the small watershed. The valley and, more specifically, the wet area (green-line) is typically very narrow and confined. Slope is generally between one and two percent. There is one plant community component in the reference condition. The valley is typically narrow and confined.

Associated sites

R034AA011UT	Riparian Complex Perennial Gravelly VIII/E4 (Northwest Territory Sedge)
R034AA239UT	Semi-desert Silt (Basin big sagebrush/ Bluebunch wheatgrass)

Similar sites

R034AA011UT	Riparian Complex Perennial Gravelly VIII/E4 (Northwest Territory Sedge)
-------------	---

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	(1) <i>Carex nebrascensis</i> (2) <i>Juncus arcticus ssp. littoralis</i>

Physiographic features

This site is in the Bear River Basin in the Middle Rocky Mountain Province of the Rocky Mountain System. It typically occurs between 6500 and 6700 feet with valley slopes less than 5 percent. The water table can be found at the surface to over 60 inches below the surface depending on the distance from the stream channel and fluvial surface.

Table 2. Representative physiographic features

Landforms	(1) Drainageway
Flooding duration	Brief (2 to 7 days) to long (7 to 30 days)
Flooding frequency	Frequent to occasional
Ponding frequency	None
Elevation	1,951–2,042 m
Slope	1–5%
Water table depth	51–102 cm
Aspect	Aspect is not a significant factor

Climatic features

The average annual precipitation is 10 to 15 inches. Peak precipitation occurs as snow from October through April. This site has cold winters and short summers. Occasional convective thunderstorms produce small amounts of rain from June through September. The average annual temperature is 25 to 55 degrees Fahrenheit. The freeze-free period ranges from 95 to 115 days.

Table 3. Representative climatic features

Frost-free period (average)	82 days
Freeze-free period (average)	115 days
Precipitation total (average)	356 mm




Figure 1. Monthly precipitation range




Figure 2. Monthly average minimum and maximum temperature

Influencing water features

Slope wetlands are typically found where there is a discharge of groundwater to the surface. Main water sources are groundwater return flow and interflow from surrounding uplands, and precipitation. Water dynamics are dominated by down-slope unidirectional water flow. Slope wetlands lose water primarily by saturation subsurface and surface flows and by evaporation. These wetlands may develop channels, but the channels only convey water away from the wetland. This site may degrade and have a more defined channel.

Soil features

This site occurs in narrow drainageways. The soils on this site are loamy to clay loam and are influenced by a high water table. Soils on this site are poorly to moderately drained depending on distance from water source and distance from bottom of drainageway. The water table fluctuates between 20 and 40 inches. Soil temperature and moisture regime is frigid and xeric, trending ustic.

Table 4. Representative soil features

Parent material	(1) Alluvium
Surface texture	(1) Loam (2) Clay loam
Drainage class	Poorly drained to moderately well drained
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Subsurface fragment volume <=3" (Depth not specified)	0%
Subsurface fragment volume >3" (Depth not specified)	0%


Ecological dynamics

This site is characterized by narrow drainageways that support species tolerant of high water tables. The water source in this site is predominantly ground water. This site may have a small channel in the bottom of the narrow valley, but this channel only serves to move water away from the site and is not typically influenced by fluvial processes. This site is typically found in low order watershed reaches and is typically upstream sites from lotic riparian sites. The plant community components in this site are influenced by depth to groundwater and soils. There is only one plant community components within this site but the species change proportion as depth to ground water increases as the site transitions out to the uplands. Obligate wetland plants are found in the bottom of the drainageway with the highest water table, such as Nebraska sedge or other obligate wetland sedges. Baltic rush is found throughout the site, but typically in higher proportion moving away from the wettest part. Willows and other mesic shrubs may occur on this site in area with unique aeration, but they are not typical of this site.

The hydrology of the site can change with spring development or grazing. A small narrow channel may be present in this site. Under continuous season long grazing over many years, this small channel can widen as the sedges and rushes are eaten or trampled. Some downcutting may occur and the water table deepens and the site becomes narrower than in the reference state. The lowering of the water table also effect the plant species, increasing species more tolerant of dry conditions. Non-native species can be present in this site, particularly Kentucky bluegrass. A second plant community will develop when the water table has dropped on the previously wetter surface.

State and transition model

R034AA013UT Semi-wet Fresh Meadow (Nebraska Sedge)






Figure 3. STM for R034AA013UT

State 1
Reference State

Obligate wetland species dominate the Reference State. The water table is typically within 20 to 40 inches of the ground surface allowing wet species to dominate. This site may or may not have a small channel in the bottom of the drainageway. The channel, when present, serves to move water down valley and typically does not experience high flows during peak runoff.

Community 1.1
Obligate wetland species



Figure 4. Semi-wet Fresh meadow with no channel




Figure 5. Semi-wet Fresh Meadow



Figure 6. Semi-wet fresh meadow with small channel

The water table is high throughout the growing season in the Obligate wetland species Community Phase. Wetland obligate species dominate, with the dominant species being Nebraska sedge and Baltic rush.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	897	1681	2242
Forb	–	56	168
Total	897	1737	2410

Table 6. Soil surface cover

Tree basal cover	0%
Shrub/vine/liana basal cover	0%
Grass/grasslike basal cover	0%
Forb basal cover	0%
Non-vascular plants	0-25%
Biological crusts	0%
Litter	0-5%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0-10%
Bare ground	0-70%

Table 7. Canopy structure (% cover)

Height Above Ground (M)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.15	–	–	0-5%	0-5%
>0.15 <= 0.3	–	–	25-40%	0-5%
>0.3 <= 0.6	–	–	25-40%	–
>0.6 <= 1.4	–	–	–	–
>1.4 <= 4	–	–	–	–
>4 <= 12	–	–	–	–
>12 <= 24	–	–	–	–
>24 <= 37	–	–	–	–
>37	–	–	–	–

State 2

Current Potential State

The Current Potential State is similar to the Reference State with the addition of non-native species, such as Kentucky bluegrass.

Community 2.1

Obligate wetland species/non-native species

Community Phase 2.1 is similar to Community Phase 1.1 with the addition of non-native species, such as Kentucky

bluegrass. The introduction of Kentucky bluegrass and other similar shallow rooted grasses changes the resilience of the system. Kentucky bluegrass will not hold the soil on the site as well as deep rooted sedges. Community Phase is at risk for the incisement of the drainageway, which will channelize the water through the bottom of the drainageway and dry out the adjacent wetlands.

Table 8. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	897	1681	2242
Forb	–	56	168
Total	897	1737	2410

Table 9. Soil surface cover

Tree basal cover	0%
Shrub/vine/liana basal cover	0%
Grass/grasslike basal cover	0%
Forb basal cover	0%
Non-vascular plants	0%
Biological crusts	0-25%
Litter	10-25%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0-10%
Bare ground	50-90%

Table 10. Canopy structure (% cover)

Height Above Ground (M)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.15	–	–	0-5%	0-5%
>0.15 <= 0.3	–	–	25-40%	0-5%
>0.3 <= 0.6	–	–	25-40%	–
>0.6 <= 1.4	–	–	–	–
>1.4 <= 4	–	–	–	–
>4 <= 12	–	–	–	–
>12 <= 24	–	–	–	–
>24 <= 37	–	–	–	–
>37	–	–	–	–

State 3 Degraded State

The Degraded State represents where there is a mix of native and non-native species that dominate the site. It is also typical for the site to have a channel in the bottom of the drainageway that may or may not have been present in reference condition. The channel is typically wider than in reference condition and at a lower elevation. The incision of the channel through the drainageway effectively dries the surrounding meadows. The Degraded State is typically caused by the shift in native vegetation to non-native vegetation and through improper grazing management.

Community 3.1 Non-native species/facultative wet species



Figure 9. Degraded meadow



Figure 10. Hummocks within the small drainageway

Mesic forbs and grasses typically dominate Community Phase 3.1. There is some Nebraska sedge remaining close to the channel edge. This phase has been observed in areas where the grazing pressure has been released and the vegetation, such as sedges, colonizes across the channel, increasing production and stabilizing the site.

Transition T1A State 1 to 2

Non-native species introduced into the system

Transition T2A State 2 to 3

Channel incision occurs and water table lowered.

Additional community tables

Table 11. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1	Rhizomatous graminoids			560–2242	
	Nebraska sedge	CANE2	<i>Carex nebrascensis</i>	560–2242	25–80
	Grass-like, perennial	2GLP	<i>Grass-like, perennial</i>	6–112	1–10
	Grass, perennial	2GP	<i>Grass, perennial</i>	0–56	0–5
	spikerush	ELEOC	<i>Eleocharis</i>	0–56	0–5
	scratchgrass	MUAS	<i>Muhlenbergia asperifolia</i>	0–6	0–5
2	Non-rhizomatous perennial graminoids			0–168	
	Grass, perennial	2GP	<i>Grass, perennial</i>	0–112	0–10
	Grass-like (not a true grass)	2GL	<i>Grass-like (not a true grass)</i>	0–56	0–5
3	Annual graminoids			0–112	
	Grass, annual	2GA	<i>Grass, annual</i>	0–56	0–5
	Grass-like, annual	2GLA	<i>Grass-like, annual</i>	0–56	0–5
Forb					
4	Perennial forb			0–112	
	Forb, perennial	2FP	<i>Forb, perennial</i>	0–56	0–10
	goldenrod	SOLID	<i>Solidago</i>	0–56	0–5
	common yarrow	ACMI2	<i>Achillea millefolium</i>	0–6	0–2
5	Annual forb			0–56	
	Forb, annual	2FA	<i>Forb, annual</i>	0–56	0–10

Table 12. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1	Rhizomatous graminoids			560–2242	
	Nebraska sedge	CANE2	<i>Carex nebrascensis</i>	560–2018	25–75
	creeping bentgrass	AGST2	<i>Agrostis stolonifera</i>	1–168	1–15
	Grass-like, perennial	2GLP	<i>Grass-like, perennial</i>	6–112	1–10
	Kentucky bluegrass	POPR	<i>Poa pratensis</i>	6–67	1–10
	spikerush	ELEOC	<i>Eleocharis</i>	0–56	0–5
	Grass, perennial	2GP	<i>Grass, perennial</i>	0–56	0–5
	scratchgrass	MUAS	<i>Muhlenbergia asperifolia</i>	0–6	0–5
2	Non-rhizomatous graminoids			0–168	
	Grass, perennial	2GP	<i>Grass, perennial</i>	0–112	0–10
	Grass-like (not a true grass)	2GL	<i>Grass-like (not a true grass)</i>	0–56	0–5
	meadow foxtail	ALPR3	<i>Alopecurus pratensis</i>	0–28	0–5
3	Annual graminoids			0–112	
	Grass, annual	2GA	<i>Grass, annual</i>	0–56	0–5
	Grass-like, annual	2GLA	<i>Grass-like, annual</i>	0–56	0–5
Forb					
4	Perennial forb			0–112	
	bull thistle	CIVU	<i>Cirsium vulgare</i>	0–112	0–5
	Forb, perennial	2FP	<i>Forb, perennial</i>	0–56	0–10
	goldenrod	SOLID	<i>Solidago</i>	0–56	0–5
	common dandelion	TAOF	<i>Taraxacum officinale</i>	0–11	0–5
	common yarrow	ACMI2	<i>Achillea millefolium</i>	0–6	0–2
5	Annual forb			0–56	
	Forb, annual	2FA	<i>Forb, annual</i>	0–56	0–10

Hydrological functions

This site is found high in the watershed in 34A, where there is little influence from heavy snow pack and subsequent spring runoff. This site also has spring influence providing ground water at the site through the growing season.

Inventory data references

Data gathered by qualified range professionals within NRCS and cooperating partners.

Other references

Kleinfelder, D., S. Swanson, G. Norris, and W. Clary. 1992. Unconfined compressive strength of some streambank soils with herbaceous roots. *Soil Sci. Am. J.* 56: 1920-1925.

Manning, M.E., S.R. Swanson, T. Svejcar, and J. Trent. 1989. Rooting characteristics of four intermountain meadow community types. *Journal of Range Management* 42(4): 309-312.

Rosgen, D.L., 1996. *Applied River Morphology*. Wildland Hydrology Books, Pagosa Springs, Colorado, and Ft. Collins, CO.

Ryan, G. 1975. *Aquatic habitat inventory of streams on National Resource Land*. Western Interstate Commission for Higher Education, Boulder, Co. 65p.

Smith, D.G. 1976. Effect of vegetation on lateral migration of anastomosed channels of a glacier meltwater river. Geological Society of American Bulletin 87: 857-860.

Steed, J.E. and L.E. DeWald. 2003. Transplanting sedges (*Carex* spp.) in southwestern riparian meadows. Restoration Ecology 11(2): 247-256.

Thorne, C.R. 1990. Effects of vegetation on riverbank erosion and stability. In: Thornes, J.B. (ed.). Vegetation and Erosion. John Wiley and Sons Ltd. pp 125-144.

USDA, NRCS. 2008. (Electronic) Field Office Technical Guide. Available online at: http://efotg.nrcs.usda.gov/efotg_locator.aspx.

USDA, Natural Resource Conservation Service. 2004. National Forestry Handbook. Available online at: <http://soils.usda.gov/technical/nfhandbook/>.

USDA, NRCS. 2007. The PLANTS Database. National Plant Data Center, Baton Rouge, LA 70874-4490 USA. Available online at: <http://plants.usda.gov>.

USDA, NRCS. 2003. National Range and Pasture Handbook. Available online at: <http://www.glti.nrcs.usda.gov/technical/publications/nrph.html>.

USDA, NRCS Soil Survey Manuals for appropriate counties within MLRA 34A.

USDA, NRCS, 2007. Southerland, W. B., Technical Supplement 3E, National Engineering Handbook 654, Rosgen Stream Classification Technique – Supplemental Materials.

USDA, USFS, 2010. Fire Effects Information System Database. Available online at: <http://www.fs.fed.us/database/feis/>.

USDI, USGS, 2010. National Water Information System. Available online at: <http://waterdata.usgs.gov/nwis/>.

Contributors

Sarah Quistberg

Approval

Kirt Walstad, 9/07/2023

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	05/18/2024
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

14. **Average percent litter cover (%) and depth (in):**

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

17. **Perennial plant reproductive capability:**
