

## Ecological site R034BY330UT Upland Stony Loam (Pinyon-Utah Juniper)

Last updated: 3/05/2022  
Accessed: 05/08/2024

### Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

|   |                                 |
|---|---------------------------------|
| Author(s)/participant(s)                    | V. Keith Wadman (NRCS Retired). |
| Contact for lead author                     | shane.green@ut.usda.gov         |
| Date  | 03/10/2012                      |
| Approved by                                 | Kirt Walstad                    |
| Approval date                               |                                 |
| Composition (Indicators 10 and 12) based on | Annual Production               |

### Indicators

- Number and extent of rills:** None to few rills present. Some minor rill development may occur on steeper slopes or on areas located below exposed bedrock, or other water shedding areas, where increased runoff may occur. Any rills present should be <1 inch deep, fairly short (<8 feet) and somewhat widely spaced (6-8 feet). Minor rill development may be observed following major thunderstorm or spring runoff events but should heal during the next growing season.
- Presence of water flow patterns:** A very few sinuous flow patterns wind around perennial plants and surface rock. Evidence of flow patterns is expected to increase somewhat with slopes greater than 15%. Water flow patterns are long (15-20 feet), narrow (<1 foot wide), and spaced widely (10-20 yards) on gentle slopes (<15%) and more closely (<10 yards) on steeper slopes (>15%).
- Number and height of erosional pedestals or terracettes:** Small pedestals may form at the base of plants that occur on the edge of water flow patterns, but should not show any exposed roots. Terracettes are fairly common, forming behind debris dams of small to medium sized litter (up to 2 inches in diameter) in water flow patterns. These debris dams may accumulate smaller litter (leaves, grass and forb stems) and sediment.
- Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 15–20%. (Soil surface is typically covered 35-75% surface fragments). Most bare ground is associated with water flow patterns, rills, and gullies. Poorly developed biological soil crusts that are interpreted as functioning as bare ground should be recorded as bare ground.

- 
5. **Number of gullies and erosion associated with gullies:** None to rare on slopes < 15%. On steeper slopes and on areas below exposed bedrock, a few gullies may occur. Where they do occur, their length often extends from the exposed bedrock to where the gully reaches a stream or other area where water and sediment accumulate. Gullies may show slightly more indication of erosion as slope increases, or as the site occurs adjacent to steep sites/watershed with concentrated flow patterns.
- 
6. **Extent of wind scoured, blowouts and/or depositional areas:** None to very few. Trees break the wind and reduce the potential for wind erosion. The channers on the soil surface help armor it and reduce the potential for wind erosion.
- 
7. **Amount of litter movement (describe size and distance expected to travel):** Most litter resides in place with some redistribution caused by water movement. Minor litter removal may occur in flow channels with deposition occurring within 1 to 2 feet at points of obstruction. The majority of litter accumulates at the base of plants. Some grass leaves and small twigs (grass stems) may accumulate in soil depressions adjacent to plants. Woody stems are not likely to move. However, some litter movement is expected (up to 6 feet) with increases in slopes >15% and/or increased runoff resulting from heavy thunderstorms.
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** This site should have an erosion rating of 4 or 5 under the plant canopies, and a rating of 3 to 4 in the interspaces. The average should be a 4. Vegetation cover, litter, biological soil crusts and surface rock reduce erosion.
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**  
(Whetrock) Soil surface horizon is typically 0 to 2 inches deep. Texture is an extremely channery loam, structure is typically very thin platy parting to moderate very fine subangular blocky. Color is a grayish brownish (10YR5/2). A mollic epipedon ranges to a depth of 8 inches. Use the specific information for the soil you are assessing found in the published soil survey to supplement this description.
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Spatial distribution of well developed biological soil crusts (where present) intercept raindrops reducing splash erosion and provide areas of surface detention to store water allowing additional time for infiltration. Crowns of trees and accumulating litter at base of trees appear to create a micro-topography that may enhance development of water flow patterns below the drip line of the canopy. Significant increases in Pinyon-juniper canopy (beyond the reference state) reduces understory vegetation causing an associated increase in runoff.
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None, There may be dense layers of rock fragments or other naturally occurring hard layers found in the soil subsurface. These should not be considered to be compaction layers.
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Trees (two-needle pinyon/Utah juniper) > = Sprouting shrubs (black sage, alderleaf mountain mahogany)>>

Perennial bunchgrasses (bluebunch wheatgrass, Indian ricegrass).

Sub-dominant: Non-sprouting shrubs (Wyoming big sagebrush) > Perennial bunchgrasses (needle-and-thread, prairie junegrass)>> Forbs (yellow milkvetch, cushion wild buckwheat) > Biological soil crusts.

Other: Functional/structural groups may appropriately contain non-native species if their ecological function is the same as the native species in the reference state. Biological soil crust is variable in its expression where present on this site and is measured as a component of ground cover. Forbs can be expected to vary widely in their expression in the plant community based upon departures from average growing conditions.

Additional: Factors contributing to temporal variability include insects and other pathogens (mistletoe), drought, extreme precipitation events, etc. Factors contributing to spatial variability include slope, amount of rock fragments, aspect, etc. Following recent natural disturbances such as fire, drought or insects that may remove the woody vegetation, forbs and perennial grasses (herbaceous species) may become more dominate in the community. These conditions may reflect a functional community phase within the reference state.

- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** During years with average to above-average precipitation, there should be very little recent mortality or decadence apparent in trees, shrubs, or grasses. During severe (multi-year) drought up to 20% of the pinyons and junipers may die, either from drought, insect damage or pathogens such as mistletoe. There may be partial mortality on individual bunchgrasses and shrubs during drought periods, and complete mortality of individual plants during severe drought periods, particularly on the shallower and coarser soils associated with this site. Because woody stems may persist for many years, both pinyons and junipers (especially older trees) will normally have dead stems within the plant canopy.
- 
14. **Average percent litter cover (%) and depth ( in):** Litter cover (including under plants) nearly all of which should be fine litter. Depth should be 1 leaf thickness in the interspaces, up to 1/2" under shrub canopies, and up to 1" under tree canopies. Litter cover may increase to 30% on some years due to increased production of plants.
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** Annual production in air-dry herbage should be approximately 700 - 800#/acre on an average year, but could range from 400 to 1000#/acre during periods of prolonged drought or above average precipitation.
- 
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Few invasive species are capable of dominating this site. When invasion does occur, cheatgrass, and various mustard species are the most likely to invade.
- 
17. **Perennial plant reproductive capability:** All perennial plants should have the ability to reproduce in all years, except in extreme drought years. There are no restrictions on either seed or vegetative reproduction. Some seedling recruitment of major species is present during average and above average growing years.

---