

Ecological site R034BY342UT

Upland Very Steep Shallow Loam (Pinyon-Utah Juniper)

Last updated: 3/05/2022
Accessed: 05/16/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 034B–Warm Central Desertic Basins and Plateaus

MLRA 34B occurs in is in Utah (70 percent) and Colorado (30 percent). It makes up about 12,850 square miles (33,290 square kilometers). A small part of the area is in the High Plateaus of Utah Section of the Colorado Plateaus Province of the Intermontane Plateaus. The northern part of the MLRA occurs in the Uinta Basin Section, which is bounded by the Uinta Mountains to the north, the Wasatch Range to the west, the Roan Plateau to the south, and the Rabbit Hills to the east. The southern part of the MLRA occurs in the northern third of the Canyon Lands Section. This section is bounded by the Roan Plateau to the north, the Wasatch Plateau to the west, the southern end of the San Rafael Swell to the south, and the western slope of the Rocky Mountains to the east. Elevation ranges from 4,100 feet (1,250 meters) near Green River, Utah, to 7,500 feet (2,285 meters) at the base of the Wasatch Range and the Roan Plateau.

Most of this area is covered by residual basin-floor materials and materials washed in from the surrounding mountains and plateaus. Shale and sandstone are the dominant rock types. The Tertiary-age Green River, Uinta, and Duchesne Formations dominate the northern part of the MLRA. The southern part is dominated by Cretaceous-age materials with lesser amounts of Jurassic and Triassic materials. The dominant Cretaceous formations are Mancos Shale, Dakota Sandstone, and the members of the Mesa Verde Group. The dominant Jurassic formations are the Morrison, Entrada, and Navajo. The dominant Triassic formations are the Chinle and Moenkopi. Quaternary alluvial, eolian, and glacial deposits occur in both parts of the MLRA.

The average annual precipitation in most of this area ranges from 6 to 10 inches (150 to 255 millimeters). A small part of this area receives as much as 24 inches of annual precipitation.

Much of the precipitation occurs as high-intensity, convective thunderstorms during the period July through September. May and June are usually the drier months. Precipitation is more evenly distributed throughout the year in the northern part of the MLRA than in the southern part, where there is a significant peak in late summer. The northern part of the MLRA receives more precipitation as snow during winter than the southern part. The average annual temperature ranges from 41 to 54 degrees F (5 to 12 degrees C). The freeze-free period averages 170 days and ranges from 110 to 235 days.

The dominant soil orders in this MLRA are Aridisols and Entisols. Mollisols occur at the higher elevations, particularly in the northern part of the MLRA. The dominant soil temperature regime is mesic, and the dominant soil moisture regime is aridic. The soils receiving less than 8 inches (205 millimeters) of precipitation annually have an aridic soil moisture regime. The soils receiving 8 to 12 inches (205 to 305 millimeters) have an aridic soil moisture regime that borders on ustic. The soils receiving 12 to 16 inches (305 to 405 millimeters) generally have an ustic soil moisture regime that borders on aridic. The dominant soil mineralogy is mixed and soils are formed in slope alluvium or residuum derived from shale or sandstone. Many of the soils are shallow or moderately deep to shale or sandstone bedrock. The soils at the lower elevations generally have significant amounts of calcium carbonate, salts, and gypsum.

Ecological site concept

The soils of this site formed mostly in colluvium or residuum weathered from sandstone and shale. Surface soils are very channery silt loam, extremely bouldery loam to very bouldery sandy loam in texture. Rock fragments may be present on the soil surface and throughout the profile, but generally make up less than 50 percent of the soil volume. These soils are shallow, well-drained, and have moderately slow to moderately rapid permeability. pH is slightly to moderately alkaline.. Available water-holding capacity ranges from 1 to 3 inches of water in the upper 20 inches of soil. The soil moisture regime is mostly aridic ustic and the soil temperature regime is mesic. Precipitation ranges from 12-16 inches annually.

Associated sites

R034BY230UT	Semidesert Shallow Loam (Salina Wildrye)
R034BY322UT	Upland Shallow Loam (Two-Needle Pinyon / Utah Juniper)
R034BY338UT	Upland Very Steep Loam (Pinyon-Utah Juniper)

Table 1. Dominant plant species

Tree	(1) <i>Pinus edulis</i>
Shrub	(1) <i>Cercocarpus montanus</i>
Herbaceous	Not specified

Physiographic features

This site occurs on hills, canyons, and mountain slopes. Slopes can range from 40 to 80%.

Table 2. Representative physiographic features

Landforms	(1) Mountain slope (2) Hill (3) Canyon
Runoff class	Very high
Flooding frequency	None
Ponding frequency	None
Elevation	1,798–2,621 m
Slope	40–80%
Ponding depth	Not specified
Water table depth	Not specified
Aspect	N, SW

Climatic features

Average annual precipitation is 12 to 16 inches. Approximately 60% occurs as rain from March through October. Much of this summer precipitation occurs as convection thunderstorms. On the average, November through February are the driest months and July through October are the wettest months. The soil temperatures are in the mesic regime. The average freeze-free period is 85 to 125 days. In average years, plants begin growth around March and April and end growth in October. Plants usually remain green until frost in October except in drier than average years. There is usually an active green-up period in the fall. The most rapid growth occurs during April, May and June.

Table 3. Representative climatic features

Frost-free period (average)	
-----------------------------	--

Freeze-free period (average)	125 days
Precipitation total (average)	356 mm

Influencing water features

None

Soil features

The soils of this site formed mostly in colluvium or residuum weathered from sandstone and shale. Surface soils are very channery silt loam, extremely bouldery loam to very bouldery sandy loam in texture. Rock fragments may be present on the soil surface and throughout the profile, but generally make up less than 50 percent of the soil volume. These soils are shallow, well-drained, and have moderately slow to moderately rapid permeability. pH is slightly to moderately alkaline.. Available water-holding capacity ranges from 1 to 3 inches of water in the upper 20 inches of soil. The soil moisture regime is mostly aridic ustic and the soil temperature regime is mesic. Precipitation ranges from 12-16 inches annually.

Modal Soil: Atchee CNV-Sil Moist 50-80% — loamy-skeletal, mixed (calcareous), mesic Lithic Ustic Torriorthents

Table 4. Representative soil features

Parent material	(1) Colluvium—sandstone and shale (2) Residuum—sandstone and shale
Surface texture	(1) Very channery silt loam (2) Extremely bouldery loam (3) Very bouldery sandy loam
Family particle size	(1) Loamy (2) Loamy-skeletal
Drainage class	Well drained
Permeability class	Moderate to moderately rapid
Depth to restrictive layer	10–51 cm
Soil depth	10–51 cm
Surface fragment cover <=3"	10–37%
Surface fragment cover >3"	5–56%
Available water capacity (Depth not specified)	2.54–7.62 cm
Calcium carbonate equivalent (Depth not specified)	0–30%
Electrical conductivity (Depth not specified)	0–2 mmhos/cm
Sodium adsorption ratio (Depth not specified)	0–5
Soil reaction (1:1 water) (Depth not specified)	7.4–8.4
Subsurface fragment volume <=3" (Depth not specified)	5–30%
Subsurface fragment volume >3" (Depth not specified)	0–15%

Ecological dynamics

Ecological Dynamics of the Site

As ecological condition deteriorates due to anthropogenic disturbances and/or the alteration of the natural disturbance regime, native perennial bunch grasses decrease while pinyon and juniper increase. When the potential natural plant community is burned and/or drought conditions persist, pinyon and juniper decrease while native perennial bunch grasses increase.

State 1: Reference State

This state represents the natural variability and dynamics of this site that occurred naturally. This state includes the dominant biotic communities that would have occurred on this ecological site prior to European Settlement. The dominant aspect of this site is Pinyon and Utah Juniper with an understory of shrubs and associated grasses. Fluctuations in species compositions and relative production may change from year to year dependent upon abnormal precipitation or other climatic factors. The primary disturbance mechanisms for this site in reference condition include drought, insects, and infrequent fire. Because catastrophic disturbances like a crown fire or drought happen with long intervals, these communities have long periods of succession, (i.e. long periods of dense Pinyon and Juniper)—300-600 years in upland/foothills ecological site zone and 300 to 1,000 in semi-desert ecological site zone. Typically, fires occurred in late spring through mid-summer following several wet years that allowed the fine fuels to become more contiguous. The higher in elevation and higher precipitation area would burn more frequently as they would have more fine fuels in the understory. The timing of drought, and fire, coupled with surface disturbance can dictate whether the community can stay within the reference state or if the community transitions into another state.

When this site is at or near its potential, pinyon pine and Utah juniper dominate the site and make up over 80 percent of the plant community. Understory production is very limited and provides marginal amounts of forage for livestock and or wildlife. It does provide good escape cover and thermal cover for deer. When the tree canopy cover exceeds 30 percent, diversity, both plant and animal drops to its lowest level.

Community Phase 1.1: Pinyon-Juniper Woodland

A well-developed understory with a canopy of younger pinyon and Utah juniper. At this stage Utah juniper may be dominant over pinyon. Pinyon trees are more susceptible to drought, insects, and disease than Utah juniper trees. In fact, it is difficult to identify methods beside fire that naturally reduce Utah juniper. After long periods of drought weaken the pinyon trees, beetle kills can become quite extensive, especially after the droughts. Drought periods can also weaken and reduce the understory. Plant establishment is mainly limited by the available moisture. Biological crusts can be highly developed and diversified in the large interspaces between trees.

When the tree canopy ranges from 15 to 30 percent, a wide variety of grasses, forbs, and shrubs will also be present in addition to the pinyon pine and Utah juniper. During this tree canopy stage, diversity of plant and animal species will reach its peak.

Community Phase Pathway 1.1A

This pathway occurs when events create a wetter climate cycle, favor pinyon and perennial bunch grass establishment. Following several favorable precipitation years and lack of surface disturbances, native perennial plants will reestablish.

Community Phase Pathway 1.1B

This pathway is very unlikely but can occur when a fire is able to move through the community. Two situations can make this occur: 1) a fire can carry in the understory after several wet years allow fine fuels to accumulate, or 2) as the woodland approaches the later stages of development where canopies become dense and crown sizes have increased, and thus community phase becomes susceptible to crown fires.

Community Phase 1.2: Mature Pinyon-Juniper Woodland

Mature pinyon and Utah juniper woodland characterized this community phase. When weather patterns favor an increase of pinyon and Utah juniper canopy with the associated understory of shrubs, grasses and forbs. Depending on the timing of precipitation, cool season grasses, like Indian ricegrass or warm season grasses like blue grama could be dominant. Interspaces supporting highly developed biological crusts are common.

Community Phase Pathway 1.2A

This pathway occurs during and after events such as drought or insect/pathogen outbreaks. Droughts and insects can kill the trees, increasing nutrient availability in the system. Due to the natural conditions of drought, grasses typically do not take up the extra nutrients in the long term. In the short term, grasses and forbs may increase for a few years until juniper and pinyon recover.

Community Phase Pathway 1.2B

This pathway is very unlikely but can occur when a fire is able to move through the community phase. Two situations can make this occur: 1) a fire can carry in the understory after several wet years allow fine fuels to accumulate, or 2) as the woodland approaches the later stages of development where canopies become dense and crown sizes have increased, and thus community phase becomes susceptible to crown fires.

Community Phase 1.3: Perennial Grassland/Shrubland with scattered PJ

The overall aspect of this community phase is grasses and shrubs with scattered pinyon and Utah juniper. The herbaceous understory has a mix of grasses and forbs. This community phase is a result of a crown fire or sufficiently large and hot ground fire that will kill many of the trees, combined with sufficient seed-banks and moisture for reestablishment of grasses and forbs. It is common that after a crown fire many patches of trees will remain unburned, because of fire's unpredictability and broken topography. This leaves a seed bank for the burned areas. This community phase is very short lived in comparison to the other community phases in this state.

When the tree canopy ranges from 0 - 15 percent; grasses, forbs, and shrubs will produce approximately 80 to 90 percent of the total production. When the tree canopy level is reduced by fire, chaining and/or application of herbicides, forage production will be at its highest level for big game animals as well as domestic livestock.

Community Phase Pathway 1.3A

This pathway occurs when the climate favors the establishment and growth of trees. More energy is taken-up and stored in the trees as the length between fires and droughts increase. In addition, when shrubs establish on the site they can provide safe-sites for tree establishment furthering the presence of trees.

Transition T1A

This transition from the native perennial bunchgrass and shrub understory in the reference state to a state that has been invaded by naturalized species such as crested wheatgrass (blown in or seeded), cheatgrass, annual wheatgrass and other introduced or exotic plants. This transition occurs as natural and/or management actions favor an increase in non-native grasses and forbs, especially annuals. Possible events include the presence of invasive species, improper livestock grazing, extended droughts, and fire combined with an available seed source of non-native species.

State 2: Current Potential State

This state is very similar to the reference state, except that non-native grasses and/or forbs are now present in all community phases. The current potential state may include introduced (seeded) or invasive nonnative species. The invasive plants are present in sparse amounts in this state. Natural disturbance are still drought, insects, and infrequent fires still influence the community shifts. The human caused disturbance drivers (i.e. domestic livestock grazing, vegetation manipulation, and recreational activities (i.e. OHV use)) are now present. This shift in species composition could affect nutrient cycling, hydrology and soil stability. At this time there is no known way to effectively remove the non-native plants from the site once they have become established. State 2 is in jeopardy of moving to State 3 (Pinyon-Juniper Invasive State) when remaining native understory plants are stressed and invasive species have increased till they are dominant.

Community Phase 2.1: Pinyon-Juniper Woodland

A well-developed understory with a canopy of younger Pinyon and Utah juniper. At this stage Utah juniper may be dominant over Pinyon. Pinyon trees are more susceptible to drought, insects, and disease than Utah Juniper trees. In fact, it is difficult to identify methods beside fire that naturally reduce Utah juniper. After long periods of drought weaken the Pinyon trees, beetle kills can become quite extensive, especially after the droughts. Drought periods can also weaken and reduce the understory. Plant establishment is mainly limited by the available moisture. Biological crusts can be highly developed and diversified in the large interspaces between trees. Sparse invasive

introduced plants species would be present in this phase.

Community Phase Pathway 2.1A

This pathway occurs when events create a wetter climate cycle, favor Pinyon and perennial bunch grass establishment. Following several favorable precipitation years and lack of surface disturbances, native perennial bunch grasses and forbs will reestablish.

Community Phase Pathway 2.1B

This pathway is very unlikely but can occur when a fire or vegetation manipulation happens to the trees. Two situations can make this occur: 1) a fire can carry in the understory after several wet years allow fine fuels to accumulate, or 2) as the woodland approaches the later stages of development where canopies become dense and crown sizes have increased, and thus community phase becomes susceptible to crown fires. Seeding after the tree removal may be necessary to help facilitate the return of understory species. Seeding depending on the species may take this community phase into state 4 (Seeded State).

Community Phase 2.2: Mature Pinyon-Juniper Woodland

Mature pinyon and Utah juniper woodland with a well-developed understory would characterized this community phase. This phase supports a diverse understory of grasses, forbs and shrubs. Depending on the timing of precipitation, cool season grasses, like Indian ricegrass or warm season grasses like blue grama could be dominant. Interspaces supporting highly developed biological crusts are common. Sparse invasive introduced plants species would be present in this phase.

Community Phase Pathway 2.2A

This pathway occurs during and after events such as drought or beetle infestations. Droughts and insects can kill pinyon trees, increasing nutrient availability in the system. Due to the natural conditions of drought, grasses typically do not take up the extra nutrients in the long term. In the short term, grasses and forbs may increase for a few years until Juniper recover. Utah Juniper are more able to compete for these nutrients and became the dominant overstory tree over time.

Community Phase Pathway 2.2B

This pathway is very unlikely to occur naturally with fire. But, vegetation manipulation can be used to remove trees. Two situations occur naturally: 1) a fire can carry in the understory after several wet years allow fine fuels to accumulate, or 2) as the woodland approaches the later stages of development where canopies become dense and crown sizes have increased, and thus community phase becomes susceptible to crown fires. Seeding after the tree removal may be necessary to help facilitate the return of understory species. Seeding depending on the species may take this community phase into state 4 (Seeded State).

Community Phase 2.3: Perennial Grassland/Shrubland with scattered PJ

The overall aspect of this community phase is grassland with scattered pinyon and Utah juniper. The herbaceous understory has a mix of grasses and forbs. This community phase is a result of a crown fire or sufficiently large and hot ground fire that will kill many of the trees, combined with sufficient seed-banks and moisture for reestablishment of grasses and forbs. It is common that after a crown fire many patches of trees will remain unburned, because of fire's unpredictability and broken topography. This leaves a seed bank for the burned areas. This community phase is very short lived in comparison to the other community phases in this state. Sparse invasive introduced plants species would be present in this phase.

Community Phase Pathway 2.3A

This pathway occurs when the climate favors the establishment and growth of trees. More energy is taken-up and stored in the trees as the length between fires and droughts increase. In addition, when shrubs establish on the site they can provide safe-sites for tree establishment furthering the presence of trees.

Transition T2A

When this transition to state 3 occurs the site has lost much of its expected resistance and resilience. At this point natural and/or management actions have decreased the understory to a point where erosion increases. Reduced influence from fire, insects, and drought could cause the tree canopy to close, effectively reducing the herbaceous understory thus facilitating the transition. Improper grazing and or increase surface disturbance combined with periods of drought can facilitate this transition because soil stability is lost and susceptibility to soil loss increases.

Transition T2B

This transition is from tree canopy reduction and re-establishment of grasses and forbs. If the community is approaching state 3 (pinyon juniper invasive state), due to a loss of understory and increase in invasive plants this pathway of seeding could be preferable to doing nothing. This pathway may facilitate the recovery of the soils. The infrequent naturally occurring fires could also cause this transition. Reseeding after a fire may be the only way to successfully restore the ecological dynamics to a site. Either way this pathway involves large energy and monetary inputs by man.

State 3: Pinyon-Juniper Invasive State

This state occurs when there is an absence of natural disturbance (i.e. Insects and drought and/or fire) over long time frames. Also, management actions could have allowed trees to become very mature and have effectively closed out the understory. Invasive plants have increased in abundance. This state has the lowest resiliency and resistance of any state in this model. There may be no practicable way back to the Current Potential State (State 2), due to the large amounts of energy and monetary inputs that are needed. Seeding, with either natural disturbance and/or vegetation management to transition it to State 3 (Seeded State) may be the best long term option for this site.

Community Phase 3.1: PJ Woodland with Invasive Plants

A lack of understory with a canopy of older Pinyon and Juniper, where plant interspaces are very large and connected. This community phase occurs when natural or management actions allow for the increase in Pinyon and Utah juniper and a decrease in the grass and forb understory. Invasive introduced plant species would be present in this phase and are increasing.

Community Phase Pathway 3.1A

This pathway occurs when events such as frequent fire or drought remove the trees and shrubs, and facilitate the continued establishment of cheatgrass or other invasive annuals. Cheatgrass will typically invade/increase in tree/shrub interspaces when PJ communities are degraded. Once the cheatgrass establishes the amount and continuity of fine fuels increases. This can reduce the fire return interval and shorten the time between fires. When fire eliminates the tree/shrub/native grass component, it completes the conversion to annual dominant community phase. Cheatgrass and other invasive annuals can persist for long periods of time. Once a fire or a drought removes the trees/shrubs, it is difficult to reestablish because, not only has the fire return interval been shortened to a time that will not allow seedling establishment, the soil and other abiotic factors have been altered.

Community Phase 3.2: Invasive Annuals

This state is characterized by annual grasses like cheatgrass, annual wheatgrass dominating the understory. Also, invasive forbs like storkbill, halogeton and others may be present. This community phase has active erosion under the pinyon and Utah juniper canopy. Utah Juniper has allelopathic effects on some plant (i.e. Sandberg bluegrass, blue grama).

Community Phase Pathway 3.2A

This pathway is when there is a lack of fire and/or disturbance. The fire return interval lengthens. This could be done by having firebreaks and/or fire suppression which will allow the perennial species a chance to establish with natural processes or with vegetation manipulation.

Transition T3A

Vegetation treatment can transition it to a seeded state. Because of the soils (shallow and/or rocky) and the unpredictable precipitation, this pathway should be used cautiously. This pathway involves large energy and monetary inputs by man.

State 4: Seeded State

This state is a result of seeding plant species. Vegetation manipulation may or may not have been done depending on disturbance history of the location. The trees were removed and adapted grasses, forbs and shrubs are established. Plants can be native or introduced depending on the desired management goals. If grazing tolerant species were established these communities can better withstand grazing and other disturbances. Due to the shallow or rocky soils and unpredictable precipitation patterns, it is difficult to establish grasses from seed, so this state may be hard to achieve and require large energy inputs.

Community Phase 4.1: Seeded Grassland/Shrubland

This community phase appears as a grassland with scattered shrubs and trees. The vegetative production is typically higher than in the current potential state, depending on grass species seeded; however the grass is still sparse due to the low water holding capacity of soils associated with pinyon and juniper.

Community Phase Pathway 4.1A

This pathway occurs when events favor the establishment of shrubs and trees, including long periods without disturbances.

Community Phase 4.2: Seeded with PJ

This community phase has a dense under story of introduced grasses and forbs, but a canopy of pinyon and Utah juniper are establishing. Native perennial grasses, forbs, and shrubs may also be starting to establish. Interspaces are filled with biological crusts and herbaceous plants.

Community Phase Pathway 4.2A

This pathway occurs as trees and shrubs are removed from the community, either naturally through insect herbivory or through vegetation manipulation by man.

Transition T4A

This transition occurs when events favor the establishment and dominance of invasive annuals. Events may include an extended drought, surface disturbance such as off road vehicle use, and/or a shortened fire return interval, all of which can stress the native perennial bunchgrasses.

State and transition model




Figure 1. STM

Legend

1.1A, 2.1A, 1.3A, 2.3A – wetter climate period, time without disturbance
 1.1B, 2.1B, 1.2B, 2.2B – Fire
 1.2A, 2.2A – Insect and pathogen outbreaks, drought, small scale fires
 T1A – Establishment of non-native invasive plants
 T2A, T4A – reduced fire return interval, increase in invasive plants in understory, extended drought
 T2B, T3A – Vegetation manipulation
 3.1A – drought, reduced fire return interval
 3.2A, 4.1A – time without disturbance
 4.2A – vegetation manipulation, insect or pathogen outbreaks, drought

Figure 2. Legend

State 1

Reference State

Community 1.1

Reference State

The dominant aspect of the plant community is pinyon Utah juniper. The understory composition by air-dry weight is approximately 35 percent perennial grasses, 15 percent forbs and 50 percent shrubs.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Shrub/Vine	196	280	364
Grass/Grasslike	138	196	256
Forb	59	84	110
Total	393	560	730

Table 6. Ground cover

Tree foliar cover	59-61%
Shrub/vine/liana foliar cover	29-31%
Grass/grasslike foliar cover	19-21%
Forb foliar cover	9-11%
Non-vascular plants	0%
Biological crusts	0%
Litter	0%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0%
Bedrock	0%
Water	0%
Bare ground	0%

Table 7. Canopy structure (% cover)

Height Above Ground (M)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.15	—	—	—	—
>0.15 <= 0.3	—	—	—	9-11%
>0.3 <= 0.6	—	—	19-21%	—
>0.6 <= 1.4	—	29-31%	—	—
>1.4 <= 4	59-61%	—	—	—
>4 <= 12	—	—	—	—
>12 <= 24	—	—	—	—
>24 <= 37	—	—	—	—
>37	—	—	—	—

Additional community tables

Table 8. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Tree					
0	Dominant Trees			93–131	
	twoneedle pinyon	PIED	<i>Pinus edulis</i>	74–93	—
	Utah juniper	JUOS	<i>Juniperus osteosperma</i>	19–31	—
	Rocky Mountain juniper	JUSC2	<i>Juniperus scopulorum</i>	0–7	—
Shrub/Vine					
0	Dominant Shrubs			124–216	
	alderleaf mountain mahogany	CEMO2	<i>Cercocarpus montanus</i>	62–93	—
	Mexican cliffrose	PUME	<i>Purshia mexicana</i>	31–62	—
	Utah serviceberry	AMUT	<i>Amelanchier utahensis</i>	31–62	—
3	Sub-Dominant Shrubs			129–241	
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	62–93	—
	Wyoming big sagebrush	ARTRW8	<i>Artemisia tridentata ssp. wyomingensis</i>	7–12	—
	curl-leaf mountain mahogany	CELE3	<i>Cercocarpus ledifolius</i>	7–12	—
	yellow rabbitbrush	CHVI8	<i>Chrysothamnus viscidiflorus</i>	7–12	—
	Torrey's jointfir	EPTO	<i>Ephedra torreyana</i>	7–12	—
	mormon tea	EPVI	<i>Ephedra viridis</i>	7–12	—
	slender buckwheat	ERMI4	<i>Eriogonum microthecum</i>	7–12	—
	singleleaf ash	FRAN2	<i>Fraxinus anomala</i>	7–12	—
	broom snakeweed	GUSA2	<i>Gutierrezia sarothrae</i>	7–12	—
	antelope bitterbrush	PUTR2	<i>Purshia tridentata</i>	7–12	—
	stemless mock goldenweed	STAC	<i>Stenotus acaulis</i>	7–12	—
	mountain snowberry	SYOR2	<i>Symphoricarpos oreophilus</i>	7–12	—
Grass/Grasslike					
0	Dominant Grasses			94–185	
	Indian ricegrass	ACHV	<i>Achnatherum hymenoides</i>	31–62	

	Indian ricegrass	ACR11	<i>Acriatherium hymenoides</i>	31-62	-
	saline wildrye	LESAS	<i>Leymus salinus ssp. salinus</i>	31-62	-
	bluebunch wheatgrass	PSSP6	<i>Pseudoroegneria spicata</i>	31-62	-
1	Sub-Dominant Grasses			103-197	
	Grass, annual	2GA	<i>Grass, annual</i>	31-62	-
	Grass, perennial	2GP	<i>Grass, perennial</i>	31-62	-
	Geyer's sedge	CAGE2	<i>Carex geyeri</i>	7-12	-
	squirreltail	ELEL5	<i>Elymus elymoides</i>	7-12	-
	needle and thread	HECO26	<i>Hesperostipa comata</i>	7-12	-
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	7-12	-
	muttongrass	POFE	<i>Poa fendleriana</i>	7-12	-
Forb					
0	Dominant Forbs			27-63	
	rock goldenrod	PEPU7	<i>Petroragia pumila</i>	7-19	-
	spiny phlox	PHHO	<i>Phlox hoodii</i>	7-19	-
	rayless tansyaster	MAGR2	<i>Machaeranthera grindelioides</i>	7-12	-
	thickleaf beardtongue	PEPA6	<i>Penstemon pachyphyllus</i>	7-12	-
2	Sub-Dominant Forbs			63-231	
	Forb, annual	2FA	<i>Forb, annual</i>	31-62	-
	Forb, perennial	2FP	<i>Forb, perennial</i>	31-62	-
	littleleaf pussytoes	ANMI3	<i>Antennaria microphylla</i>	0-7	-
	Fendler's sandwort	ARFE3	<i>Arenaria fendleri</i>	0-7	-
	Holboell's rockcress	ARHO2	<i>Arabis holboellii</i>	0-7	-
	woolly locoweed	ASMO7	<i>Astragalus mollissimus</i>	0-7	-
	Brenda's yellow cryptantha	CRFL5	<i>Cryptantha flava</i>	0-7	-
	nodding buckwheat	ERCE2	<i>Eriogonum cernuum</i>	0-7	-
	cushion buckwheat	EROV	<i>Eriogonum ovalifolium</i>	0-7	-
	fineleaf hymenopappus	HYFI	<i>Hymenopappus filifolius</i>	0-7	-
	manybranched ipomopsis	IPPO2	<i>Ipomopsis polycladon</i>	0-7	-
	blue flax	LIPE2	<i>Linum perenne</i>	0-7	-
	juniper biscuitroot	LOJU	<i>Lomatium juniperinum</i>	0-7	-
	tailcup lupine	LUCAC3	<i>Lupinus caudatus ssp. caudatus</i>	0-7	-
	hoary tansyaster	MACA2	<i>Machaeranthera canescens</i>	0-7	-
	heartleaf twistflower	STCO6	<i>Streptanthus cordatus</i>	0-7	-
	Pacific aster	SYCHC	<i>Symphyotrichum chilense var. chilense</i>	0-7	-
	Navajo tea	THSU	<i>Thelesperma subnudum</i>	0-7	-

Animal community

Because of steep slopes, this site is grazed very little by livestock. This site produces food and cover for wildlife. Wildlife using this site include jackrabbit, wood rat, pinyon jay, coyote, mule deer, and elk.

Hydrological functions

The soil is in hydrologic group D. The runoff curve numbers are 80 through 89 depending on the condition of the

watershed.

Recreational uses

This site has aesthetic appeal but limited recreational value.

Wood products

There are approximately 8 to 10 cords of firewood per acre in the potential plant community when the average tree diameter is 5 inches at 1 foot. About 20 posts and 12 Christmas trees are produced per acre.

Contributors

George Cook

Approval

Kirt Walstad, 3/05/2022

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	V. Keith Wadman (NRCS Retired).
Contact for lead author	shane.green@ut.usda.gov
Date	03/22/2012
Approved by	Kirt Walstad
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills:** Rills are common to very common. Their expression may be less defined where coarse fragments (i.e., gravels and/or channers) dominate the soil surface. Rill occurrence may increase slightly on areas located below exposed bedrock or other water shedding areas where increased runoff may occur. Rills should be <1 inches deep, somewhat long (8 to 16 feet) and somewhat widely spaced (8-12 feet). An increase in rill development may be observed immediately following major thunderstorm or spring runoff events.

- 2. Presence of water flow patterns:** Sinuous flow patterns are common and wind around perennial plants and surface rock. Evidence of flow patterns is expected to increase somewhat as slopes approach 80%. Water flow patterns are long (20 to 30 feet), somewhat narrow (1 to 2 feet wide), and spaced widely (5 to 10 yards) and more closely spaced (3 to 6 yards) on slopes nearing 70 to 80%.

- 3. Number and height of erosional pedestals or terracettes:** Small pedestals will form at the base of plants that occur on the edge of water flow patterns, 2 to 4% of plants show minor exposed roots. Terracettes are fairly common, forming

behind debris dams of small to medium sized litter (up to 2 inches in diameter) in water flow patterns. These debris dams may accumulate smaller litter (leaves, grass and forb stems) and sediment.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 15–25%. (Soil surface is typically covered by 25-50% surface fragments). Most bare ground is associated with water flow patterns, rills, and gullies. Bare ground spaces not associated with flow patterns should not be greater than 1 to 2 feet in diameter.
5. **Number of gullies and erosion associated with gullies:** A few gullies may occur. Any gullies present may extend down the length of the site until they reach a stream or other area where water and sediment is diverted or accumulates. Gullies show slightly more indication of erosion as slopes approach 80%, or where the site occurs adjacent to watershed areas with concentrated flow patterns.
6. **Extent of wind scoured, blowouts and/or depositional areas:** None. Trees and shrubs along with surface coarse fragments on this site break the wind and reduce the potential for wind erosion.
7. **Amount of litter movement (describe size and distance expected to travel):** Because of the site's very steep slopes, some litter redistribution downslope caused by water movement is normal. Some litter removal may occur in flow channels with deposition occurring within 3 to 5 feet at points of obstruction. The majority of litter still accumulates at the base of plants. Some grass leaves, stems and small woody twigs may accumulate in soil depressions adjacent to plants. Woody stems are likely to move 1 to 2 feet. A slight increase in litter movement is expected following runoff resulting from heavy spring runoff or thunderstorms.
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** This site should have an erosion rating of 4 or 5 under the plant canopies, and a rating of 2 to 4 in the interspaces. The average should be a 4. Vegetation cover, litter, biological soil crusts and surface rock reduce erosion.
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
(Gompers, 50-80% slopes) Soil surface A horizon is typically 0 to 3 inches deep. Structure is moderate thin platy. Color is brown (10YR5/3). Use the specific information for the soil you are assessing found in the published soil survey to supplement this description.
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Good spatial distribution of well developed biological soil crusts (where present) intercept raindrops, reducing splash erosion and providing areas of increased surface detention to store water, allowing additional time for infiltration. Crowns of trees and accumulating litter at base of trees appear to create a micro-topography that may enhance development of water flow patterns below the drip line of the canopy.
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None. Fractured shale bedrock occurs at about 18 inches.

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Trees (two-needle pinyon/Utah juniper) >> Sprouting shrubs (alderleaf mountain mahogany, Mexican cliffrose) > cool season perennial grasses (Indian ricegrass, Salina wildrye).

Sub-dominant: cool season perennial grasses (bluebunch wheatgrass, muttongrass) = > forbs (grassy rock goldenrod, cushion wild phlox) > biological soil crusts.

Other: Functional/structural groups may appropriately contain non-native species if their ecological function is the same as the native species in the reference state. Biological soil crust is variable in its expression where present on this site and is measured as a component of ground cover. Forbs can be expected to vary widely in their expression in the plant community based upon departures from average growing conditions.

Additional: Factors contributing to temporal variability include insects and other pathogens (mistletoe), drought, extreme precipitation events, etc. Factors contributing to spatial variability include slope, amount of rock fragments, aspect, etc. Following a recent disturbance such as fire, drought or insects that may remove the woody vegetation, forbs and perennial grasses (herbaceous species) may become more dominate in the community. These conditions may reflect a different functional community phase within the reference state.

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** During years with average to above-average precipitation, there should be very little recent mortality or decadence apparent in trees, shrubs, or grasses. During severe (multi-year) drought up to 20% of the pinyons and junipers may die, either from drought, insect damage or pathogens such as mistletoe. There may be partial mortality on individual bunchgrasses and shrubs during drought periods, and complete mortality of individual plants during severe drought periods. Because woody stems persist for many years, both pinyons and junipers (especially older trees) will normally have dead stems within the plant canopy.

14. **Average percent litter cover (%) and depth (in):** Cover should be composed mostly of fine litter. Depth should vary from a 1 or 2 leaf thickness in the interspaces, to up to 1/2" under canopies, and up to 1" under tree canopies. Litter cover may increase to 30% on some years due to increased production of plants.

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** Annual production in air-dry herbage should be approximately 450 - 550#/acre on an average year, but could range from 300 to 700#/acre during periods of prolonged drought or above average precipitation.

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Few invasive species are capable of dominating this site. When invasion does occur, cheatgrass, alysium, and mustard species are the most likely species to invade.

17. **Perennial plant reproductive capability:** All perennial plants should have the ability to reproduce in all years, except in

extreme drought years. There are no restrictions on either seed or vegetative reproduction. Some seedling recruitment of major species is present during average and above average growing years.
