

Ecological site DX035X03B811 **Sandy Loam Upland 17-25" p.z. (PIPO)**

Accessed: 05/16/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 035X–Colorado Plateau

This ecological site occurs in Common Resource Area 35.8 - the Colorado Plateau Ponderosa Pine Forests

The Common Resource Area occurs within the Colorado Plateau Physiographic Province. Elevations range from 6800 to 8500 feet and precipitation averages 17 to 25 inches per year. Vegetation includes ponderosa pine, white fir, aspen, pinyon, juniper, Gambel oak, big sagebrush, ceanothus, blue elderberry, muttongrass, upland sedge, and big wildrye, mountain muhly, Arizona fescue, pine dropseed, and blue grama. The soil temperature regime ranges from mesic to frigid and the soil moisture regime is typic ustic. This unit occurs within the Colorado Plateau Physiographic Province and is characterized by a sequence of flat to gently dipping sedimentary rocks eroded into plateaus, valleys and deep canyons. Sedimentary rock classes dominate the plateau with volcanic fields occurring for the most part near its margin.

Associated sites

DX035X03B826	Sandstone Upland 17-25" p.z. (PIPO)
DX035X03B827	Sandstone Hills 17-25" p.z. (PIPO)
R035XH821AZ	Meadow 17-25" p.z.

Table 1. Dominant plant species

Tree	(1) <i>Pinus ponderosa</i> (2) <i>Quercus gambelii</i>
Shrub	(1) <i>Artemisia tridentata</i> ssp. <i>wyomingensis</i> (2) <i>Ceanothus fendleri</i>
Herbaceous	(1) <i>Poa fendleriana</i> (2) <i>Carex rossii</i>

Legacy ID

F035XH811AZ

Physiographic features

This ecological site occurs on toeslopes, fan terraces, backslopes and footslopes on stable landslides, plateaus and mesas. Soils are moderately deep to deep. Surface textures are sandy loam, fine sandy loam and loam. The site does not benefit from run-on moisture.

Table 2. Representative physiographic features

Landforms	(1) Fan (2) Terrace (3) Landslide
Elevation	2,073–2,591 m
Slope	0–15%
Aspect	Aspect is not a significant factor

Climatic features

Winter-summer moisture ratios are typically 70:30 on the west side of this LRU and shift to 60:40 on the east side. Late spring is usually the driest period, and early fall moisture can be sporadic. Summer rains fall from June through September; moisture originates in the Gulf of Mexico and creates convective, usually brief, intense thunderstorms. Cool season moisture from October through May tends to be frontal; It originates in the Pacific Ocean and the Gulf of California and falls in widespread storms with longer duration and lower intensity. Precipitation generally comes as snow from October into April. Snowpack can persist for 3-4 months, although it may disappear in exposed areas during prolonged dry weather. Summer daytime temperatures are typically 80-90 but can exceed 95 degrees F. Winter temperatures of around 0 degrees F are common and can reach -25 degrees F.

Table 3. Representative climatic features

Frost-free period (average)	100 days
Freeze-free period (average)	130 days
Precipitation total (average)	635 mm

Influencing water features

The soil moisture on this ecological site comes from precipitation. The site does not benefit significantly from run-on moisture. The loamy surface texture of the soil allows the site to capture the majority of the winter storms and the light to moderate summer storms if the site has good vegetative cover. Intense summer thunderstorms will produce runoff, reducing the amount of effective rainfall.

Soil features

Soils are deep to moderately deep. Surface textures are sandy loam to fine sandy loam. Subsoil textures are sandy

loam, sandy, loam, loam and some deep soils will have clay horizon. Parent materials are alluvium and residuum from sandstone, shale and basalt. Available water capacity is moderate to very high. Water erosion hazard potential is slight to moderate; wind erosion potential is moderate to high. Soils are non-sodic, non-saline. pH range is 6.6-7.3.

Soil survey map unit components that have been correlated to this ecological site include:

SSA-699 Hualapai-Havasupai Area 030-Pinntank;

SSA-713 Chinle Area 56-Tuntsa moderately;

SSA-715 Fort Defiance Area NM/AZ MU 52-Kunz, 54 Klizhin family, 130-Manuelito;

SSA-717 Shiprock Area AZ/NM 600, 604 & 614 Yahmore

Table 4. Representative soil features

Parent material	(1) Alluvium–sandstone and shale (2) Residuum–basalt
Surface texture	(1) Sandy loam (2) Fine sandy loam (3) Loam
Family particle size	(1) Loamy
Drainage class	Moderately well drained to well drained
Permeability class	Moderately slow to moderate
Soil depth	51–152 cm
Surface fragment cover <=3"	0–15%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	10.92–24.38 cm
Calcium carbonate equivalent (0-101.6cm)	0%
Electrical conductivity (0-101.6cm)	0–2 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0–5
Soil reaction (1:1 water) (0-101.6cm)	6.6–7.3
Subsurface fragment volume <=3" (Depth not specified)	0–15%
Subsurface fragment volume >3" (Depth not specified)	0%

Ecological dynamics

An ecological site is not a precise assemblage of species for which the proportions are the same from place to place or from year to year. In all plant communities, variability is apparent in productivity and occurrence of individual species. Spatial boundaries of the communities; however, can be recognized by characteristic patterns of species composition, association, and community structure. The historic climax plant community for this ecological site has been described by sampling relict or relatively undisturbed sites and/or reviewing historic records. The historic climax plant community is the plant community that evolved over time with the soil forming process and long term changes in climatic conditions of the area. It is the plant community that was best adapted to the unique combination of environmental factors associated with the site.

Natural disturbances, such as drought, fire, grazing of native fauna, and insects, are inherent in the development and maintenance of these plant communities. The effects of these disturbances are part of the range of characteristics of the ecological site. Fluctuations in plant community structure and function caused by the effects of natural disturbances help establish the boundaries and characteristics of an ecological site. They are accounted for as part of the range of characteristics of the ecological site. Recognizable plant community phases are identified in the reference state of the ecological site. Some sites may have a small range of variation, while others have a large range. Some plant community phases may exist for long periods of time, while others may only occur for a couple of years after a disturbance.

Deterioration of the plant community, hydrology, or soil site stability on an ecological site can result in crossing a threshold or potentially irreversible boundary to another state, or equilibrium. This can occur as a result of the loss of soil surface through erosion, the loss of the stability of the site due to disturbances that cause active erosion on the site, increases in the amounts and/or patterns or runoff from rainstorms, changes in availability of surface and subsurface water, significant changes in plant structural and functional types, or the introduction of non-native species. When these thresholds are crossed, the potential of the ecological site to return to the historic climax plant community can be lost, or restoration will require significant inputs. There may be multiple states possible for an ecological site, determined by the type and or severity of disturbance.

The known states and transition pathways for this ecological site are described in the state and transition model. Within each state, there may be one or more known plant community phases. These community phases describe the different plant community that can be recognized and mapped across this ecological site. The state and transition model is intended to help land users recognize the current plant community on the ecological site, and the management options for improving the plant community to the desired plant community.

Plant production information in this site description is standardized to the annual production on an air-dry weight basis in near normal rainfall years.

State and transition model

F035XH811AZ Sandy Loam Upland 17-25" p.z.




Figure 4. 358 SandyLoamUpland S&T

State 1

Reference State

This state is comprised of mature ponderosa pines and tall Gambel oaks with a tree canopy of 40-55 percent. Other trees present in minor amounts include scattered twoneedle pinyon pine, Rocky Mountain juniper and shorter Gambel oaks. The understory is comprised mainly of grasses with a mix of annual and perennial forbs with scattered shrubs. Common grasses and grasslikes include blue grama, bottlebrush squirreltail, White Mountain sedge, needle and thread, muttongrass and mountain muhly. Shrubs occasionally present in small amounts include broom snakeweed, mountain big sagebrush, spineless horsebrush and rubber rabbitbrush.

Community 1.1

Historic Climax Plant Community



This potential plant community includes a tree canopy cover of 40-55%, with an understory of grasses, forbs, shrubs and small scattered trees.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	224	308	392
Tree	84	140	196
Forb	56	92	129
Shrub/Vine	56	92	129
Total	420	632	846

Figure 6. Plant community growth curve (percent production by month). AZ3581, 35.8 17-25" p.z. all sites. Growth begins in the spring, most growth occurs during the summer rainy season..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	0	4	10	24	21	23	13	5	0	0

Community 1.2 Light PIPO Overstory

Patchy mature PIPO reduced canopy cover, with less cover in understory trees. Dense and abundant grass and forb cover.

Community 1.3 Dense PIPO Overstory

Pole sized PIPO with very dense overstory Dense upper and lower canopies (50-60%) sparse understory.

Pathway 1.1a Community 1.1 to 1.2

Partial removal of mature PIPO canopy through timber harvest, or Bark beetle infestation. This results in higher production on the forest floor of grass, forb and to some extent shrubs.

Conservation practices

Forage Harvest Management
Forest Trails and Landings
Forest Stand Improvement

Woody Residue Treatment
Prescribed Forestry

Pathway 1.1b

Community 1.1 to 1.3

1.1B (1.2) Openings are filled in by natural regeneration of Ponderosa pine, forming an even aged tight canopy of sapling to pole size PIPO

Conservation practices

Tree/Shrub Pruning
Forest Stand Improvement
Prescribed Forestry

Pathway 1.2a

Community 1.2 to 1.3

1.2a Openings are filled in by natural regeneration of Ponderosa pine, forming an even aged tight canopy of sapling to pole size PIPO

Pathway 1.3a

Community 1.3 to 1.1

1.3a Trees mature and are thinned either naturally or through Pre-commercial thinning forming a mature canopy.

Conservation practices

Forest Stand Improvement

Pathway 1.3b

Community 1.3 to 1.2

1.3b Trees mature and are thinned more heavily than pathway 1.3a, through fire or timber harvest forming a patchy and mature canopy.

Conservation practices

Forest Stand Improvement
Prescribed Forestry

State 2

Canopy removed Grass and Forb Dominated

Community 2.1

Canopy removed Grass and Forb Dominated

Tree canopy has been severely reduced (<10%) and site becomes dominated by Grasses Forbs and shrubs. There is not adequate regeneration of Ponderosa Pine on this site.

Transition 1

State 1 to 2

Catastrophic wildfire event removes 90 to 95% of tree overstory.

Restoration pathway 1

State 2 to 1

Tree planting and other Forest land improvements return site to a forested condition.

Conservation practices

Forest Stand Improvement
Agroforestry Planting

Additional community tables

Table 6. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					
1	Grasses/Grasslikes			224–392	
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	93–155	–
	White Mountain sedge	CAGE	<i>Carex geophila</i>	62–123	–
	squirreltail	ELEL5	<i>Elymus elymoides</i>	62–93	–
	needle and thread	HECO26	<i>Hesperostipa comata</i>	7–31	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	7–31	–
	mountain muhly	MUMO	<i>Muhlenbergia montana</i>	7–31	–
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	0–31	–
	muttongrass	POFE	<i>Poa fendleriana</i>	7–31	–
	Grass, perennial	2GP	<i>Grass, perennial</i>	0–31	–
	sleepygrass	ACRO7	<i>Achnatherum robustum</i>	0–12	–
	pine dropseed	BLTR	<i>Blepharoneuron tricholepis</i>	0–12	–
	little bluestem	SCSC	<i>Schizachyrium scoparium</i>	0–12	–
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	0–12	–
	Grass, annual	2GA	<i>Grass, annual</i>	0–12	–
Forb					
2	Forbs			56–129	
	pingue rubberweed	HYRI	<i>Hymenoxys richardsonii</i>	12–62	–
	silvery lupine	LUAR3	<i>Lupinus argenteus</i>	7–31	–
	white sagebrush	ARLU	<i>Artemisia ludoviciana</i>	0–19	–
	spreading fleabane	ERDI4	<i>Erigeron divergens</i>	0–19	–
	trailing fleabane	ERFL	<i>Erigeron flagellaris</i>	0–19	–
	redroot buckwheat	ERRA3	<i>Eriogonum racemosum</i>	0–12	–
	pineywoods geranium	GECA3	<i>Geranium caespitosum</i>	0–12	–
	hairy false goldenaster	HEVI4	<i>Heterotheca villosa</i>	0–12	–
	Forb, perennial	2FP	<i>Forb, perennial</i>	0–12	–
	western yarrow	ACMIO	<i>Achillea millefolium</i> var. <i>occidentalis</i>	0–12	–
	nodding onion	ALCE2	<i>Allium cernuum</i>	0–12	–
	whitemargin pussytoes	ANMA5	<i>Antennaria marginata</i>	0–12	–
	small-leaf pussytoes	ANPA4	<i>Antennaria parvifolia</i>	0–12	–
	Carruth's sagewort	ARCA14	<i>Artemisia carruthii</i>	0–12	–

	prairie sagewort	ARFR4	<i>Artemisia frigida</i>	0–12	–
	upright blue beardtongue	PEVI4	<i>Penstemon virgatus</i>	0–12	–
	broom-like ragwort	SESPM	<i>Senecio spartioides</i> var. <i>multicapitatus</i>	0–12	–
	globemallow	SPHAE	<i>Sphaeralcea</i>	0–12	–
	Forb, annual	2FA	<i>Forb, annual</i>	0–7	–
	woolly plantain	PLPA2	<i>Plantago patagonica</i>	0–7	–
	flatspine stickseed	LAOC3	<i>Lappula occidentalis</i>	0–7	–
Shrub/Vine					
3	Shrubs			56–129	
	mountain big sagebrush	ARTRV	<i>Artemisia tridentata</i> ssp. <i>vaseyana</i>	7–62	–
	broom snakeweed	GUSA2	<i>Gutierrezia sarothrae</i>	7–31	–
	spineless horsebrush	TECA2	<i>Tetradymia canescens</i>	0–25	–
	rubber rabbitbrush	ERNA10	<i>Ericameria nauseosa</i>	0–25	–
	toadflax penstemon	PELI2	<i>Penstemon linarioides</i>	0–12	–
	Navajo yucca	YUBA2	<i>Yucca baileyi</i>	0–12	–
	creeping barberry	MARE11	<i>Mahonia repens</i>	0–12	–
	Fendler's ceanothus	CEFE	<i>Ceanothus fendleri</i>	0–12	–
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	0–12	–
	plains pricklypear	OPPO	<i>Opuntia polyacantha</i>	0–7	–
	pincushion cactus	PEDIO	<i>Pediocactus</i>	0–7	–
Tree					
4	Trees (<4.5')			84–196	
	Gambel oak	QUGA	<i>Quercus gambelii</i>	93–155	–
	ponderosa pine	PIPO	<i>Pinus ponderosa</i>	7–62	–
	Rocky Mountain juniper	JUSC2	<i>Juniperus scopulorum</i>	7–31	–
	twoneedle pinyon	PIED	<i>Pinus edulis</i>	7–31	–

Animal community

Site is fairly suitable for grazing in the summer and early fall by cattle, sheep, goats and horses prior to tree canopy exceeding 45%.

Management considerations are grazing systems, proper grazing, and water developments to aid in livestock distribution. Grazing should not damage young trees.

Water is frequently found in scattered natural wetlands and springs. Cover is good for most species. Food has good potential except where grazing is unmanaged. Topography provides escape habitat.

Recreational uses

Recreational activities include hiking, horseback riding, camping, wildlife observations and hunting. Landscape quality is good.

Wood products

All kinds of equipment is suitable for harvesting, site preparation, tree planting and precommercial thinning. Slopes

are non-restrictive. Unsurfaced roads limit equipment when wet. Water erosion is slight to moderate in bare areas; wind erosion is moderate to high.

Compaction potential is good, but soils will rut when wet. Revegetation potential is good.

Harvest fully grown trees on a sustained yield basis. Final removal cuts and intermediate cuts are desirable. Thick stands need thinned to improve growth rates of trees left. Periodic ground fires will reduce dense understory vegetation competing for moisture. Slopes are gentle and soils are deep, so mechanical tree removal is not limited. Control unwanted pests to limit tree damage and loss. Fire hazard becomes extreme as understory fuel load builds. Site is good for replanting. There is slight limitation to seedling mortality; expected mortality rate is less than 25%. Natural regeneration is slightly limited; low end of productive pine sites. Seedlings need protection from grazing. Plant competition is moderately limited due to dense grass cover and grazing that may impede regeneration. Windthrow hazard: slight limitation, rooting depth greater than 30".

Table 7. Representative site productivity

Common Name	Symbol	Site Index Low	Site Index High	CMAI Low	CMAI High	Age Of CMAI	Site Index Curve Code	Site Index Curve Basis	Citation
ponderosa pine	<i>PIPO</i>	55	60	270	276	—	—	—	

Type locality

Location 1: San Juan County, NM	
Township/Range/Section	T22N R19W S25
General legal description	Washington Pass Quad, Chuska Mountains, Navajo Reservation
Location 2: Apache County, AZ	
UTM zone	N
UTM northing	665157
UTM easting	3980735
General legal description	Sawmill Quad, Defiance Plateau area, Navajo Reservation. About 5.5 miles due north of Sawmill on route 9702

Other references

Updates and revisions for this ESD were conducted as part of a 2007-2012 Interagency Technical Assistance Agreement between the Bureau of Indian Affairs–Navajo Region and the NRCS-Arizona.

Contributors

Dan Carroll
Dean Schlichting
HSH
Kenneth Gishi
Larry D. Ellicott
Steve Barker

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. Number and extent of rills:

2. Presence of water flow patterns:

3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

5. Number of gullies and erosion associated with gullies:

6. Extent of wind scoured, blowouts and/or depositional areas:

7. Amount of litter movement (describe size and distance expected to travel):

8. Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):

9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):

10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:

11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):

12. **Functional/Structural Groups** (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence** (include which functional groups are expected to show mortality or decadence):
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production** (this is TOTAL above-ground annual-production, not just forage annual-production):
-

16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
-

17. **Perennial plant reproductive capability:**
-