

Ecological site DX035X03B812 Sandy Upland 17-25" p.z. (PIPO)

Accessed: 06/30/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 035X–Colorado Plateau

This ecological site occurs in Common Resource Area 35.8 - the Colorado Plateau Ponderosa Pine Forests

The Common Resource Area occurs within the Colorado Plateau Physiographic Province. Elevations range from 6800 to 8500 feet and precipitation averages 17 to 25 inches per year. Vegetation includes ponderosa pine, white fir, aspen, pinyon, juniper, Gambel oak, big sagebrush, ceanothus, blue elderberry, muttongrass, upland sedge, and big wildrye, mountain muhly, Arizona fescue, pine dropseed, and blue grama. The soil temperature regime ranges from mesic to frigid and the soil moisture regime is typic ustic. This unit occurs within the Colorado Plateau Physiographic Province and is characterized by a sequence of flat to gently dipping sedimentary rocks eroded into plateaus, valleys and deep canyons. Sedimentary rock classes dominate the plateau with volcanic fields occurring for the most part near its margin.

Table 1. Dominant plant species

Tree	(1) <i>Pinus ponderosa</i> (2) <i>Pseudotsuga menziesii</i>
Shrub	(1) <i>Quercus gambelii</i> (2) <i>Ceanothus fendleri</i>

Herbaceous	(1) <i>Carex geophila</i> (2) <i>Lupinus argenteus</i>
------------	---

Legacy ID

F035XH812AZ

Physiographic features

This site occurs on fan terraces and toeslopes of stable landslides, mountains and plateaus. Slopes are generally 0 to 15 percent, but may be slightly steeper in spots. The soils are deep. Surface textures range from loamy fine sand to fine sandy loam. The site does not benefit significantly from run-on moisture from other sites.

Table 2. Representative physiographic features

Landforms	(1) Fan (2) Plateau (3) Structural bench
Flooding frequency	None
Ponding frequency	None
Elevation	7,600–8,500 ft
Slope	0–15%
Aspect	Aspect is not a significant factor

Climatic features

Winter-Summer moisture ratios are typically 70:30 on the west side of this CRA and shift to 60:40 on the east side. Late spring is usually the driest period and early fall moisture can be sporadic. Summer rains fall from June through September; moisture originates in the Gulf of Mexico and creates convective, usually brief, intense thunderstorms. Cool season moisture from October through May tends to be frontal; it originates in the Pacific and the Gulf of California and falls in widespread storms with longer duration and lower intensity. Precipitation generally comes as snow from October into April. Snowpack can persist for 3-4 months, although it may disappear in exposed areas during prolonged dry weather. Summer daytime temperatures are typically 80-90 F but can exceed 95 F. Winter temperatures around 0 F are common and can reach -25 F.

Table 3. Representative climatic features

Frost-free period (average)	100 days
Freeze-free period (average)	130 days
Precipitation total (average)	25 in

Influencing water features

Soil features

Soils are moderately deep to very deep. Surface textures range from loamy fine sand to fine sandy loam. Subsurface textures range from loamy fine sand to fine sandy loam. Parent material is alluvium from sandstone and basalt.

Water erosion hazard is slight to moderate; wind erosion is very high to high. Soils are non-saline, non-sodic. pH range is 5.6-7.3. Available water capacity is very low to very high. Soil moisture regime is typic ustic; temperature regime is frigid.

Soil survey map unit components that have been correlated to this ecological site include;

SSA-713 Chinle Area MU 59-Typic Ustipsammments;

SSA-715 Ft. Defiance Area NM/AZ 51-Sandark family;

SSA-717 Shiprock Area AZ/NM MU's 604 & 606 Sandark.

Table 4. Representative soil features

Parent material	(1) Alluvium–sandstone (2) Residuum–basalt
Surface texture	(1) Loamy fine sand (2) Sand (3) Fine sandy loam
Family particle size	(1) Sandy
Drainage class	Somewhat excessively drained to excessively drained
Permeability class	Moderately slow to rapid
Soil depth	40–80 in
Surface fragment cover <=3"	0–10%
Available water capacity (0-40in)	2–10 in
Soil reaction (1:1 water) (0-40in)	5.6–7.3
Subsurface fragment volume <=3" (Depth not specified)	0–10%

Ecological dynamics

An ecological site is not a precise assemblage of species for which the proportions are the same from place to place or from year to year. In all plant communities, variability is apparent in productivity and occurrence of individual species. Spatial boundaries of the communities; however, can be recognized by characteristic patterns of species composition, association, and community structure. The historic climax plant community for this ecological site has been described by sampling relict or relatively undisturbed sites and/or reviewing historic records. The historic climax plant community is the plant community that evolved over time with the soil forming process and long term changes in climatic conditions of the area. It is the plant community that was best adapted to the unique combination of environmental factors associated with the site.

Natural disturbances, such as drought, fire, grazing of native fauna, and insects, are inherent in the development and maintenance of these plant communities. The effects of these disturbances are part of the range of characteristics of the ecological site. Fluctuations in plant community structure and function caused by the effects of natural disturbances help establish the boundaries and characteristics of an ecological site. They are accounted for as part of the range of characteristics of the ecological site. Recognizable plant community phases are identified in the reference state of the ecological site. Some sites may have a small range of variation, while others have a large range. Some plant community phases may exist for long periods of time, while others may only occur for a couple of years after a disturbance.

Deterioration of the plant community, hydrology, or soil site stability on an ecological site can result in crossing a threshold or potentially irreversible boundary to another state, or equilibrium. This can occur as a result of the loss of soil surface through erosion, the loss of the stability of the site due to disturbances that cause active erosion on the site, increases in the amounts and/or patterns or runoff from rainstorms, changes in availability of surface and subsurface water, significant changes in plant structural and functional types, or the introduction of non-native species. When these thresholds are crossed, the potential of the ecological site to return to the historic climax plant community can be lost, or restoration will require significant inputs. There may be multiple states possible for an ecological site, determined by the type and or severity of disturbance.

The known states and transition pathways for this ecological site are described in the state and transition model.

Within each state, there may be one or more known plant community phases. These community phases describe the different plant community that can be recognized and mapped across this ecological site. The state and transition model is intended to help land users recognize the current plant community on the ecological site, and the management options for improving the plant community to the desired plant community.

Plant production information in this site description is standardized to the annual production on an air-dry weight basis in near normal rainfall years.

State and transition model




Figure 4. 358 SandyUpland S&T

State 1

Reference State

This site has a potential tree canopy cover of 65-75% dominated by ponderosa pine with a mix of Douglas fir and quaking aspen. Other tree species present in minor amounts include Gambel oak, Rocky Mountain juniper and Colorado pinyon pine. Common herbaceous species include White Mountain sedge, nodding brome, squirreltail, needle and thread, silvery lupine, western brackenfern and geranium. Low woody species present in small amounts include creeping barberry, Navajo yucca, kinnikinnick, Fendler ceanothus and short Gambel oaks. At higher elevations quaking aspen and Douglas fir will increase in canopy cover along with kinnikinnick and common juniper.

Community 1.1

Historic Cllimax Plant Community

This plant community has a tree canopy cover of 65-75%, with an understory comprised mainly of grasses and forbs, with a few scattered shrubs and short trees. The main tree stand is dominated by ponderosa pine with the secondary stand of Douglas fir and quaking aspen.

Table 5. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	150	220	310
Forb	120	175	245
Tree	105	155	215
Shrub/Vine	30	65	95
Total	405	615	865

Figure 6. Plant community growth curve (percent production by month).
 AZ3581, 35.8 17-25" p.z. all sites. Growth begins in the spring, most growth occurs during the summer rainy season..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	0	4	10	24	21	23	13	5	0	0

Community 1.2
Severely reduced PIPO overstory

1 to 10% canopy, with 10 to 20% cover in shorter trees. Dense and abundant grass and forb cover.

Community 1.3
Short dense PIPO Overstory

Pole sized PIPO with very dense overstory Dense upper and lower canopies (50-60%) sparse understory

Pathway 1.1
Community 1.1 to 1.2

Partial removal of mature PIPO canopy through timber harvest, or bark beetle infestation with drought. The reduced canopy results in higher production on the forest floor of grass, forb and to some extent shrubs.

Conservation practices

Forage Harvest Management
Forest Land Management

Pathway 1.2
Community 1.1 to 1.3

Openings are filled in by natural regeneration of ponderosa pine through favorable wet climate, forming an even aged tight canopy of sapling to pole size ponderosa pine.

Pathway 1.2
Community 1.2 to 1.3

Openings are filled in by natural regeneration of ponderosa pine, forming an even aged tight canopy of sapling to pole size ponderosa pine

Pathway 1.3
Community 1.3 to 1.1

Trees mature and are thinned either naturally or through pre-commercial forest thinning forming a mature canopy.

Conservation practices

Forest Stand Improvement
Forest Land Management

**Pathway 1.4
Community 1.3 to 1.2**

Trees mature and are thinned more heavily than 1.3 plant community, through nature fire or timber harvest (thinning) forming a patchy and mature canopy.

Conservation practices

Forest Stand Improvement
Forest Land Management

**State 2
Canopy removed Grass and Forb Dominated**

**Community 2.1
Canopy removed Grass and Forb Dominated**

Less than 5% canopy remains site is dominated by grasses and forbs with inadequate tree regeneration.

**Transition 1
State 1 to 2**

Catastrophic wildfire event removes the majority of overstory.

**Restoration pathway 1
State 2 to 1**

Tree planting and forest management

Conservation practices

Agroforestry Planting
Forest Land Management

Additional community tables

Table 6. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
Grass/Grasslike					
1	Common Perennial Grasses/Grasslikes			123–185	
	nodding brome	BRAN	<i>Bromus anomalus</i>	31–92	–
	White Mountain sedge	CAGE	<i>Carex geophila</i>	31–92	–
	squirreltail	ELEL5	<i>Elymus elymoides</i>	31–62	–
	Arizona fescue	FEAR2	<i>Festuca arizonica</i>	6–62	–
	needle and thread	HECO26	<i>Hesperostipa comata</i>	6–62	–

2	Ocassional Perennial Grasses			31-62	
	pine dropseed	BLTR	<i>Blepharoneuron tricholepis</i>	0-31	-
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	0-31	-
	mountain muhly	MUMO	<i>Muhlenbergia montana</i>	0-31	-
	muttongrass	POFE	<i>Poa fendleriana</i>	0-31	-
	western wheatgrass	PASM	<i>Pascopyrum smithii</i>	0-18	-
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	0-18	-
3	Others Grasses			0-18	
	Grass, perennial	2GP	<i>Grass, perennial</i>	0-18	-
	Grass, annual	2GA	<i>Grass, annual</i>	0-12	-
Forb					
4	Common Forbs			125-185	
	silvery lupine	LUAR3	<i>Lupinus argenteus</i>	31-62	-
	hairy brackenfern	PTAQP2	<i>Pteridium aquilinum var. pubescens</i>	31-62	-
	New Mexico groundsel	PANE7	<i>Packera neomexicana</i>	0-31	-
	woodland strawberry	FRVEB2	<i>Fragaria vesca ssp. bracteata</i>	0-31	-
	pineywoods geranium	GECA3	<i>Geranium caespitosum</i>	0-31	-
5	Occasional Forbs			31-62	
	western yarrow	ACMIO	<i>Achillea millefolium var. occidentalis</i>	0-18	-
	pussytoes	ANTEN	<i>Antennaria</i>	0-18	-
	ragleaf bahia	BADI	<i>Bahia dissecta</i>	0-18	-
	winged buckwheat	ERAL4	<i>Eriogonum alatum</i>	0-18	-
	pingue rubberweed	HYRI	<i>Hymenoxys richardsonii</i>	0-18	-
	scarlet gilia	IPAG	<i>Ipomopsis aggregata</i>	0-12	-
	Wright's deervetch	LOWR	<i>Lotus wrightii</i>	0-12	-
	wild bergamot	MOFI	<i>Monarda fistulosa</i>	0-12	-
	catchfly	SILEN	<i>Silene</i>	0-12	-
	Mt. Albert goldenrod	SOSI3	<i>Solidago simplex</i>	0-12	-
	Forb, perennial	2FP	<i>Forb, perennial</i>	0-12	-
	sanddune wallflower	ERCA14	<i>Erysimum capitatum</i>	0-12	-
	trailing fleabane	ERFL	<i>Erigeron flagellaris</i>	0-12	-
	aspen fleabane	ERSP4	<i>Erigeron speciosus</i>	0-12	-
	yellow hawkweed	HIFE	<i>Hieracium fendleri</i>	0-12	-
	Mexican woollywhite	HYME	<i>Hymenopappus mexicanus</i>	0-12	-
	James' cryptantha	CRCI3	<i>Cryptantha cinerea</i>	0-12	-
	Carruth's sagewort	ARCA14	<i>Artemisia carruthii</i>	0-12	-
	tarragon	ARDR4	<i>Artemisia dracunculus</i>	0-12	-
	Fendler's sandwort	ARFE3	<i>Arenaria fendleri</i>	0-12	-
	Forb, annual	2FA	<i>Forb, annual</i>	0-6	-
Shrub/Vine					
6	Shrubs			31-92	
	kinnikinnick	ARUV	<i>Arctostaphylos uva-ursi</i>	0-31	-
	Fendler's ceanothus	CEFE	<i>Ceanothus fendleri</i>	0-31	-

	creeping barberry	MARE11	<i>Mahonia repens</i>	0–31	–
	alderleaf mountain mahogany	CEMO2	<i>Cercocarpus montanus</i>	0–18	–
	rubber rabbitbrush	ERNA10	<i>Ericameria nauseosa</i>	0–18	–
	hairy false goldenaster	HEVI4	<i>Heterotheca villosa</i>	0–18	–
	common juniper	JUCO6	<i>Juniperus communis</i>	0–18	–
	Forb, perennial	2FP	<i>Forb, perennial</i>	0–18	–
	Shrub (>.5m)	2SHRUB	<i>Shrub (>.5m)</i>	0–12	–
	Subshrub (<.5m)	2SUBS	<i>Subshrub (<.5m)</i>	0–12	–
	Forb, annual	2FA	<i>Forb, annual</i>	0–12	–
	Navajo yucca	YUBA2	<i>Yucca baileyi</i>	0–12	–
Tree					
7	Trees (<4.5')			123–185	
	Gambel oak	QUGA	<i>Quercus gambelii</i>	92–154	–
	twoneedle pinyon	PIED	<i>Pinus edulis</i>	0–31	–
	ponderosa pine	PIPO	<i>Pinus ponderosa</i>	0–31	–
	quaking aspen	POTR5	<i>Populus tremuloides</i>	0–31	–
	Douglas-fir	PSME	<i>Pseudotsuga menziesii</i>	0–31	–

Animal community

Site is suitable for grazing by cattle, sheep, goats and horses during summer or early fall before the tree canopy exceeds 50%.

Management considerations include grazing systems, proper grazing, and avoiding livestock concentrations that would denude the vegetation exposing the soil to erosion. Grazing should not damage young trees.

Water is found in scattered springs or man-made watering facilities. Cover for wildlife is good for most species. Food has good potential except where grazing is unmanaged. Topography provides escape habitat.

Recreational uses

Recreational activities include hiking, horseback riding, camping, hunting, wildlife observation and photography. Landscape quality is good.

Wood products

Rubber tired equipment is best on these sandy soils for harvesting, site preparation, tree planting and precommercial thinning. Caution should be used on loose sands. Slopes are non-restrictive. Equipment could get stuck in sand with continuous heavy use.

Water erosion potential on cutover areas/bare ground/roads/trails/landings is slight to moderate. Wind erosion potential in this type of area is very high to high.

Compaction potential is poor. Soils are too coarse to rut. Revegetation potential is fair to good due to water holding capacity.

Harvest mature trees on a sustained yield basis. Final removal cuts and intermediate cuts are desirable. Thick stands need thinned to improve growth rate of trees left. Periodic ground fires will reduce dense understory vegetation competing for moisture. Loose sands are the only possible limitation to mechanical tree removal. Control unwanted pests to limit tree damage and loss. Fire hazard could become extreme if understory fuel load builds up. Replanting suitability is fair to good, drought sandy soils could be a problem. Seedling mortality is moderate to

severe. Natural regeneration is slightly limited because of competition for moisture. Seedlings need protection from grazing. Dense grass and forb cover may impede regeneration. Windthrow hazard is slight; rooting depth is greater than 30".

Table 7. Representative site productivity

Common Name	Symbol	Site Index Low	Site Index High	CMAI Low	CMAI High	Age Of CMAI	Site Index Curve Code	Site Index Curve Basis	Citation
ponderosa pine	<i>PIPO</i>	80	85	450	462	–	–	–	
quaking aspen	<i>POTR5</i>	80	87	325	330	–	–	–	
Rocky Mountain Douglas-fir	<i>PSMEG</i>	60	65	235	240	–	–	–	

Type locality

Location 1: San Juan County, NM	
Township/Range/Section	T23N R20W S11
General legal description	Toadlena Quad, Chuska Mountains, Navajo Reservation

Other references

Updates and revisions for this ESD were conducted as part of a 2007-2012 Interagency Technical Assistance Agreement between the Bureau of Indian Affairs–Navajo Region and the NRCS-Arizona.

Contributors

Dan Carroll
 Dean Schlichting
 HSH
 Karlynn Huling
 Kenneth Gishi
 Larry D. Ellicott
 Steve Barker

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

-
2. **Presence of water flow patterns:**
-
3. **Number and height of erosional pedestals or terracettes:**
-
4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**
-
5. **Number of gullies and erosion associated with gullies:**
-
6. **Extent of wind scoured, blowouts and/or depositional areas:**
-
7. **Amount of litter movement (describe size and distance expected to travel):**
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

14. **Average percent litter cover (%) and depth (in):**

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

17. **Perennial plant reproductive capability:**
