

# Ecological site R035XB202AZ Clayey Wash 6-10" p.z.

Accessed: 05/20/2024

#### **General information**

**Provisional**. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

### **MLRA** notes

Major Land Resource Area (MLRA): 035X–Colorado Plateau

This ecological site occurs in Common Resource Area 35.2 - the Colorado Plateau Shrub – Grasslands

Elevations range from 3800-5800 feet and precipitation averages 6 to 10 inches per year. Vegetation includes shadscale, fourwing saltbush, Mormon tea, blackbrush, Indian ricegrass, galleta, blue grama, and black grama. The soil temperature regime is mesic and the soil moisture regime is typic aridic. This unit occurs within the Colorado Plateau Physiographic Province and is characterized by a sequence of flat to gently dipping sedimentary rocks eroded into plateaus, valleys and deep canyons. Sedimentary rock classes dominate the plateau with volcanic fields occurring for the most part near its margin.

Table 1. Dominant plant species

Tree	Not specified			
Shrub	(1) Atriplex canescens			
Herbaceous	<ul><li>(1) Sporobolus airoides</li><li>(2) Bouteloua gracilis</li></ul>			

## Physiographic features

This site occurs in a bottom position with gentle slopes ranging from 0-5%. It benefits significantly from run-in moisture.

Landforms	<ul><li>(1) Wash</li><li>(2) Alluvial fan</li><li>(3) Flood plain</li></ul>
Flooding duration	Brief (2 to 7 days)
Flooding frequency	Occasional
Elevation	1,158–1,768 m
Slope	0–5%
Aspect	Aspect is not a significant factor

#### Table 2. Representative physiographic features

#### **Climatic features**

About 50% of annual precipitation is received as rain from July to September. With the least amount from May and June. Most of the moisture that comes from November to February is in the form of snow. Strong winds are common during the winter and spring.

#### Table 3. Representative climatic features

Frost-free period (average)	181 days
Freeze-free period (average)	207 days
Precipitation total (average)	254 mm

#### Influencing water features

This ecological site is located in the low areas along the drainages where runoff water flows following rainfall events. The soil moisture on this site is from both rainfall, and run-on moisture from the watershed above it. This additional water makes this site much more productive than the adjacent uplands that only receive moisture from rainfall.

#### **Soil features**

Soils in this site are deep and well drained. The surface horizons have textures ranging from clay loam to vertic clays. The subsurface textures are clay, silty clay, clay loam, or silty clay loam and contain very little gravel.

Typical taxonomic units include:

Coconino County Central Part (AZ631) Soil Map Units- 26-Navajo Clay; 51-Tours Silty Clay Loam; 52-Tours. Navajo County Central Part (AZ633) Soil Map Unit- 67-Tours.

Little Colorado River Area (AZ707) Soil Map Unit- 61-Tours.

Navajo Mountain(AZ711) Soil Map Units- 7-Berryhill family; 47-Sanfeco.

Fort Defiance Area (NM - AZ)(AZ715) Soil Map Units- 133-Wepo, 134-Wepo.

Shiprock Area (NM-AZ) (AZ717)Soil Map Unit- 527-Outpost.

#### Table 4. Representative soil features

Surface texture	(1) Clay loam (2) Clay
Family particle size	(1) Clayey
Drainage class	Moderately well drained to well drained
Permeability class	Moderately slow to very slow

Soil depth	102–152 cm
Surface fragment cover <=3"	0–5%
Available water capacity (0-101.6cm)	15.24–33.02 cm
Calcium carbonate equivalent (0-101.6cm)	0–15%
Electrical conductivity (0-101.6cm)	16–30 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0–2
Soil reaction (1:1 water) (0-101.6cm)	7.4–8.4
Subsurface fragment volume <=3" (Depth not specified)	0–5%

## **Ecological dynamics**

An ecological site is not a precise assemblage of species for which the proportions are the same from place to place or from year to year. In all plant communities, variability is apparent in productivity and occurrence of individual species. Spatial boundaries of the communities; however, can be recognized by characteristic patterns of species composition, association, and community structure. The historic climax plant community for this ecological site has been described by sampling relict or relatively undisturbed sites and/or reviewing historic records. The historic climax plant community is the plant community that evolved over time with the soil forming process and long term changes in climatic conditions of the area. It is the plant community that was best adapted to the unique combination of environmental factors associated with the site.

Natural disturbances, such as drought, fire, grazing of native fauna, and insects, are inherent in the development and maintenance of these plant communities. The effects of these disturbances are part of the range of characteristics of the ecological site. Fluctuations in plant community structure and function caused by the effects of natural disturbances help establish the boundaries and characteristics of an ecological site. They are accounted for as part of the range of characteristics of the ecological site. Recognizable plant community phases are identified in the reference state of the ecological site. Some sites may have a small range of variation, while others have a large range. Some plant community phases may exist for long periods of time, while others may only occur for a couple of years after a disturbance.

Deterioration of the plant community, hydrology, or soil site stability on an ecological site can result in crossing a threshold or potentially irreversible boundary to another state, or equilibrium. This can occur as a result of the loss of soil surface through erosion, the loss of the stability of the site due to disturbances that cause active erosion on the site, increases in the amounts and/or patterns or runoff from rainstorms, changes in availability of surface and subsurface water, significant changes in plant structural and functional types, or the introduction of non-native species. When these thresholds are crossed, the potential of the ecological site to return to the historic climax plant community can be lost, or restoration will require significant inputs . There may be multiple states possible for an ecological site, determined by the type and or severity of disturbance.

The known states and transition pathways for this ecological site are described in the state and transition model. Within each state, there may be one or more known plant community phases. These community phases describe the different plant community that can be recognized and mapped across this ecological site. The state and transition model is intended to help land users recognize the current plant community on the ecological site, and the management options for improving the plant community to the desired plant community.

Plant production data provided in this site description is standardized to the air-dry weight of one year's growth. The plant communities described in this site description are based on near normal rainfall years.

#### State and transition model

35.2 Clayey Wash 6-10" p.z.



Figure 4. 35.2 Clayey Wash S&T

### State 1 Reference State

This plant community is a perennial grassland with some shrubs and a small percentage of forbs.

#### Community 1.1 Historic Climax Plant Community

This plant community is dominated by warm and cool season perennial grasses with fourwing saltbush and other shrubs and a small percentage of forbs. Plants most likely to increase or invade on this site due to disturbance are rabbitbrush, broom snakeweed, wooly groundsel, annuals and succulents.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	504	616	785
Shrub/Vine	280	392	448
Forb	28	67	112
Total	812	1075	1345

#### Table 6. Ground cover

Tree foliar cover	0%	
Shrub/vine/liana foliar cover	4-12%	
Grass/grasslike foliar cover	6-25%	
Forb foliar cover	1-5%	
Non-vascular plants	0%	
Biological crusts	0%	
Litter	10-40%	
Surface fragments >0.25" and <=3"	0%	
Surface fragments >3"	0%	
Bedrock	0%	
Water	0%	
Bare ground	10-30%	

Figure 6. Plant community growth curve (percent production by month). AZ3521, 35.2 6-10" p.z. all sites. Growth begins in the spring and continues through the summer. Most growth in this CRA occurs in the spring using stored winter moisture..

Jan	Feb	Mar	Apr	Мау	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	1	9	20	27	14	10	11	5	3	0	0

Figure 7. Plant community growth curve (percent production by month). AZ5203, 35.2 6-10" p.z. alkali sacaton. Growth begins in the spring, most growth occurs in the summer, goes dormant in the fall.

Jan	Feb	Mar	Apr	Мау	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	5	10	20	10	20	20	10	5	0	0

Figure 8. Plant community growth curve (percent production by month). AZ5211, 35.2 6-10" p.z. fourwing saltbush. Growth begins in spring and continues through the summer. Seed stalk extension occurs in summer with seed set in the fall.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	5	10	15	20	20	15	10	5	0	0

## State 2 Natives with Introduced Annuals State

This site is a grassland with increased native shrubs along with the introduction of non-native invasive species.

#### Community 2.1 Native/Non-native Grass and Shrubland

This site is a grassland with warm and cool season grasses and increased native shrubs along with the introduction

of non-native invasive species. The amount of non-native invasive species does not change the overall function of this site. The amount of bare ground has increased on this site. Trace amounts of tamarisk may be present.

### State 3 Natives with Introduced Tree State

This site is dominated by an overstory of tamarisk with an understory of primarily annuals and few native shrubs and perennial grasses.

## Community 3.1 Introduced Tree Community



Figure 9. 35.3 Clayey Wash Community Phase 3.1

This site contains an overstory of tamarisk leaving little ground cover. Native shrubs and grasses are being replaced by non-native annual species, such as Russian thistle, in the understory.

### Transition T1A State 1 to 2

Unmanaged grazing/drought/introduction of non-native invasive species.

#### Transition T2A State 2 to 3

Unmanaged grazing, persistent drought, increase in invasive species.

# Restoration pathway R3A State 3 to 2

Managed grazing, woody species control, invasive weed control, range seeding.

### Additional community tables

 Table 7. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)						
Grass	Grass/Grasslike										
1	Dominant Warm Seas	on Grasse	280–448								
	alkali sacaton	SPAI	Sporobolus airoides	224–336	-						
	James' galleta	PLJA	Pleuraphis jamesii	56–168	-						
	blue grama	BOGR2	Bouteloua gracilis	112–168	-						
	sideoats grama	BOCU	Bouteloua curtipendula	28–140	-						

	vine mesquite	PAOB	Panicum obtusum	56–112	-
2	Cool Season Grasses			224–280	
	western wheatgrass	PASM	Pascopyrum smithii	168–280	_
	Indian ricegrass	ACHY	Achnatherum hymenoides	28–140	-
	squirreltail	ELELE	Elymus elymoides ssp. elymoides	28–140	-
3	Other Grasses			84–168	
	threeawn	ARIST	Aristida	6–34	-
	mat muhly	MURI	Muhlenbergia richardsonis	6–34	Ι
	ring muhly	MUTO2	Muhlenbergia torreyi	6–34	-
	burrograss	SCBR2	Scleropogon brevifolius	6–34	-
	tumblegrass	SCPA	Schedonnardus paniculatus	6–34	-
	spike dropseed	SPCO4	Sporobolus contractus	6–34	-
	sand dropseed	SPCR	Sporobolus cryptandrus	6–34	Ι
	Grass, annual	2GA	Grass, annual	0–22	-
	Grass, perennial	2GP	Grass, perennial	0–22	Ι
Forb					
4	Forbs			28–112	
	Forb, perennial	2FP	Forb, perennial	16–73	-
	Forb, annual	2FA	Forb, annual	12–67	Ι
	sideoats grama	BOCU	Bouteloua curtipendula	28–56	-
	burrograss	SCBR2	Scleropogon brevifolius	11–45	Ι
	tumblegrass	SCPA	Schedonnardus paniculatus	11–45	Ι
	threeawn	ARIST	Aristida	6–45	-
	spike dropseed	SPCO4	Sporobolus contractus	6–34	_
	sand dropseed	SPCR	Sporobolus cryptandrus	6–34	_
	mat muhly	MURI	Muhlenbergia richardsonis	6–34	_
	ring muhly	MUTO2	Muhlenbergia torreyi	6–34	_
	Grass, perennial	2GP	Grass, perennial	0–11	_
Shrub	/Vine				
5	Shrubs			280–448	
	fourwing saltbush	ATCA2	Atriplex canescens	280–392	_
	shadscale saltbush	ATCO	Atriplex confertifolia	11–45	_
	winterfat	KRLA2	Krascheninnikovia lanata	11–45	_
	menodora	MENOD	Menodora	11–45	_
	New Mexico groundsel	PANE7	Packera neomexicana	6–22	_
	pricklypear	OPUNT	Opuntia	6–11	_
	rabbitbrush	CHRYS9	Chrysothamnus	6–11	
	broom snakeweed	GUSA2	Gutierrezia sarothrae	6–11	_

### **Animal community**

This site is suitable for grazing during any period of the year by cows and calves, stocker cattle, sheep and horses. Prescribed grazing systems can benefit this site by allowing rest periods for the cool season species.

This site is dominated by grassland wildlife species. However, the site is transitory to almost all species because of

the proximity of water.

#### **Recreational uses**

Since this site is located on non-entrenched swales and bottoms it has good aesthetic appeal from being mostly grasslands interspersed with brushy thickets. Winters are cold and springtime is windy, however, mild summers make this site attractive for horseback riding, wildlife observation, and hunting.

#### **Type locality**

Location 1: Navajo County, AZ		
General legal description	Typical example of this site is located on the Turquoise ranch, two miles northwest of Winslow, AZ.	

#### **Other references**

Information and updates collected during 2009-2010 for this ESD was conducted as part of an Interagency Technical Assistance Agreement between the Bureau of Indian Affairs–Navajo Region and the NRCS-Arizona.

#### Contributors

Karlynn Huling Ken Gishi Larry D. Ellicott

#### Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Kenneth Gishi
Contact for lead author	State Rangeland Management Specialist - NRCS State Office - Phoenix, AZ
Date	08/27/2012
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

#### Indicators

- 1. **Number and extent of rills:** Few. Any rills present will continually form and reshape periodically due to occasional flooding on the site.
- 2. **Presence of water flow patterns:** Some water flow patterns may be present, but are highly variable due to position on landscape and dependant on run-in moisture from uplands. The high shrink/swell characteristics of the soil and occasional flooding (overland flow) will continually reshape the surface.

- Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground): This site has moderate to high potential to produce herbaceous cover and low amounts of bare ground. Bare ground ranges from 15-35%.
- 5. **Number of gullies and erosion associated with gullies:** Uncommon, some gullies may form due to location on floodplains. Any gullies present should be stable with vegetation and no signs of active erosion.
- 6. Extent of wind scoured, blowouts and/or depositional areas: Some deposition areas (from sediments) may occur due to overland flow and occasional flooding.
- 7. Amount of litter movement (describe size and distance expected to travel): Most herbaceous and woody litter will be transported throughout the site during intense overland flow/flood events. Only minor amounts of fine litter will be transported by wind.
- Soil surface (top few mm) resistance to erosion (stability values are averages most sites will show a range of values): Expected values of 1-3 in interspaces and 4-6 under the plant canopies. This site is fairly resistance to erosion when well vegetated.
- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Surface textures typically range from clay loam to clay with a thickness of 2-14 inches. Typically soil structure is moderately (thin, fine to medium) platy structure. Surface colors range from light reddish brown to reddish brown.
- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: This site is characterized as a native grassland with scattered large and half shrubs with few forbs. This site is dominated by a mix of bunch and colonizing grasses providing a diverse root structure and vegetation pattern. Perennial grasses reduce raindrop impact and slow run-in moisture (overland flow) to allow for increased infiltration.
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): None. These soils have a naturally platy structure.
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: Warm season grasses > Cool season grasses >

Sub-dominant: Large shrubs > Half shrubs > Forbs > Cacti

Additional:

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): All plants functional groups are adapted to survival in all but the most severe droughts. Severe winter droughts affect shrubs the most. Severe summer droughts affect grasses the most.
- 14. Average percent litter cover (%) and depth ( in):
- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annualproduction): Average annual production on this site is expected to be 900 to 1100 lbs/ac. in a year of average annual precipitation.
- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Tamarisk, cheatgrass, foxtail barley, silverleaf nightshade, Russian thistle, and camelthorn.
- 17. **Perennial plant reproductive capability:** All plants native to this site are adapted and are capable of producing seeds, stolons and rhizomes in all but the most severe drought.