

Ecological site R035XB270AZ Porcelanite Hills 6-10" p.z.

Accessed: 05/03/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Steve Barker
Contact for lead author	NRCS State Rangeland Management Specialist, Phoenix AZ
Date	09/20/2012
Approved by	Steve Barker
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- Number and extent of rills: A few rills occur throughout site (1-10% cover) at infrequent intervals, mostly in exposed areas. Rills may be 8 or more feet in length and are likely to form below adjacent exposed bedrock or areas where surface rock fragments are less than 15%. The number of rills and extent will increase on slopes greater than 35%, or sites with a decrease of herbaceous cover and/or immediately following high intensity storm events.
- Presence of water flow patterns: The occurrence of water flow patterns is frequent (5-10% cover) and occur throughout the site interspersed throughout the larger rock fragments. These water flow patterns are typically less than 6 feet long. As slope increase (>15%) water flow pattern occurrence and length increases. A temporary increase in water flow patterns in also expected following high intensity storm events.
- 3. Number and height of erosional pedestals or terracettes: Minor pedestalling (can occur at the base of plants and rocks as a result of natural wind and water erosion in the reference state; however, terracettes are uncommon and occur only in flow paths.
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground): 10 to 20% bare ground with 40-60% cover of porcelenite rock and gravel cover. Bare areas are small in size, and are rarely connected.
- 5. Number of gullies and erosion associated with gullies: No gullies are present.

- 6. Extent of wind scoured, blowouts and/or depositional areas: No wind scoured areas are present.
- 7. Amount of litter movement (describe size and distance expected to travel): Most herbaceous litter will be transported and concentration by wind and water in flow pathways and around obstructions, while a very small percentage stays in place. Coarse woody litter (>1/4" diameter) and duff will accumulate under shrub canopies.
- 8. Soil surface (top few mm) resistance to erosion (stability values are averages most sites will show a range of values): This site should have an average soil stability rating of 4 throughout the site.
- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness): Structure is generally weak thin platy. The A horizon will show minimal difference in structure and depth between interspaces and under plant canopies.
- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: This site is characterized by a relatively even distribution of mostly perennial grasses and low shrubs across the landscape. Canopy and basal cover are dominated by warm season grasses and evergreen shrubs. This type of plant community along with surface rock cover and slopes are somewhat effective at capturing and storing precipitation.
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): No compaction layers.
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant: Evergreen shrubs (25-35%)> Warm season colonizing grasses (15-20%) = Cool season bunch grasses (15-20%)

Sub-dominant: Deciduous shrubs (5-15%)> Warm season bunch grasses (5-10%), forbs (5-10%)

Other: Cacti (0-3%).

Additional:

13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): In a normal year up to 10 to 15% of grasses and shrubs die off. During and after drought years there can be from 10 to 25% die off of shrubs and grasses. Severe winter droughts affect shrubs, and cool season grasses the most. Severe summer droughts affect the warm season grasses the most.

- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annualproduction): Average annual production on this site is expected to be 415 lbs/ac. in a year of average annual precipitation.
- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site: Non-native species that can invade and establish on this site are cheatgrass and Russian thistle. Native species such as James' galleta, broom snakeweed, rabbitbrush and Mormon tea are native to the site, but can increase with disturbance.
- 17. **Perennial plant reproductive capability:** All plants native to this site are adapted to the climate and are capable of producing seeds, stolons and rhizomes except during the most severe droughts.