

## **Ecological site R035XY103UT Desert Clay (Castle Valley Saltbush)**

Accessed: 04/25/2024

### Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Jacob Owens (NRCS), V. Keith Wadman (NRCS, Ret.)
Contact for lead author	shane.green@ut.usda.gov
Date	12/11/2009
Approved by	Shane A. Green
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### Indicators

1. **Number and extent of rills:** Very common. Rills present could be 15 or more feet long. They could be 2-3 inches deep. As surface coarse fragments increase, rills become shorter and may not be as deep.

---

2. **Presence of water flow patterns:** Very common throughout the site. They are expected to be long and connected into drainage networks. Evidence of flow will increase with slope. As surface coarse fragments increase, flow patterns become less evident and may not be as deep.

---

3. **Number and height of erosional pedestals or terracettes:** Plants may show some pedestalling (up to .5 inch) on their down slope side. Terracettes should be few and stable. Interspaces between well developed biological soil crusts may resemble pedestals but they are actually a characteristic of the crust formation.

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 20 – 55% depending on surface rock or pararock fragments. Ground cover is based on the first raindrop impact, and bare ground is the inverse of ground cover. Ground cover + bare ground = 100%. Poorly developed biological soil crusts that are interpreted as functioning as bare ground (therefore they would be susceptible to raindrop splash erosion) should be recorded as bare ground.

---

5. **Number of gullies and erosion associated with gullies:** Present. May be found where adjacent sites/watershed provides concentrated flows into the site. Gullies should show only minor signs of active erosion and should be

somewhat stabilized with perennial vegetation. Gullies may show slightly more indication of erosion as slope steepens, or as the site occurs adjacent to sites where runoff accumulation occurs.

---

6. **Extent of wind scoured, blowouts and/or depositional areas:** No evidence of wind generated soil movement. Wind caused blowouts and deposition are not expected to be present.
- 
7. **Amount of litter movement (describe size and distance expected to travel):** Some down slope redistribution caused by water. Some litter removal may occur in flow patterns and rills with deposition occurring at points of obstruction, especially following major storm events. Litter movement will increase with slope.
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** This site should have a soil stability rating of 3 to 4 under plant canopies and a rating of 2 to 3 in the interspaces using the soil stability kit test. The average should be a 3. Surface texture is silty clay loam. Vegetation cover, litter, biological soil crusts and surface rock reduce erosion.
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Structure is weak fine subangular blocky. Surface color is light gray. Use the specific information for the soil you are assessing found in the published soil survey to supplement this description.
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Vascular plants are expected to break raindrop impact and splash erosion reducing splash erosion but not eliminating it. Spatial distribution of vascular plants slows runoff somewhat by obstructing surface flows to help create sinuous flow patterns that dissipate energy and allow time for some infiltration. Natural erosion would be expected in most storms and spring runoff.
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Castle valley saltbush.

Sub-dominant: Perennial grasses and mat saltbush.

Other: Dominance by average annual production: Non-sprouting shrubs > Warm season perennial grasses > Cool season perennial bunchgrasses > perennial and annual native forbs. Biological soil crust is variable in its expression where present on this site and is measured as a component of ground cover.

Additional: Perennial and annual forbs can be expected to vary widely in their expression in the plant community based upon departures from average growing conditions. Functional/structural groups may appropriately contain non-native species if their ecological function is the same as the native species in the reference state (e.g. Siberian Wheatgrass, Forage kochia etc.)

- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** All age classes of perennial grasses should be present on an average to above average precipitation year with age class expression likely subdued during below average years, or on sites with high (usually greater than 65%) similarity index (late seral to historic climax). In general, a mix of age classes may be expected with some dead and decadent plants present.
- 
14. **Average percent litter cover (%) and depth ( in):** Variability may occur due to weather.
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 150-200 #/acre on an average year.
- 
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Russian thistle, halogeton, kochia, common sunflower, and annual mustards.
- 
17. **Perennial plant reproductive capability:** All perennial plants should have the ability to reproduce sexually or asexually in most years, except in drought years.
-