

## **Ecological site R035XY209UT Semidesert Loam (Wyoming Big Sagebrush)**

Accessed: 05/04/2024

### **Rangeland health reference sheet**

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	V. Keith Wadman (NRCS Ret.), Paul Curtis (BLM), Dana Truman (NRCS), Robert Stager (BLM), Shane A. Green (NRCS)
Contact for lead author	shane.green@ut.usda.gov
Date	09/11/2008
Approved by	Shane A. Green
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### **Indicators**

1. **Number and extent of rills:** None to rare. Any rill formation may be seen after recent large storms on adjacent landscape settings where increased runoff may accumulate (such as areas below exposed bedrock). Such rill development should usually be limited to slopes exceeding 20%. Any rills present should be somewhat short in length (less than 6 feet long), and heal rapidly (within a couple of winters) through frost heaving.

---

2. **Presence of water flow patterns:** Flow patterns wind around perennial plant bases and show no evidence of erosion. They are short and stable and there is minor evidence of deposition. On gently sloping (< 15 % slopes) locations within the site, water flow patterns are infrequent and usually less than 3 feet. Longer water flow patterns may be found on steeper slopes (> 20 %). Water flow patterns are narrow (> 1 foot wide) and widely spaced (10-20 yards apart)

---

3. **Number and height of erosional pedestals or terracettes:** Plants should show no pedestalling. Small (>1 sq ft) terracettes should be very few. A few rare pedestals may occur on steeper slopes (> 20%) and are usually associated with water flow patterns, and should not have any exposed roots. Interspaces between well developed biological soil crusts may resemble pedestals but they are actually a characteristic of the crust formation.

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 3-20% bare ground. Ground cover is based on the first raindrop impact, and bare ground is the opposite of ground cover. Well developed biological soil crusts should not be recorded as bare ground. Poorly developed biological soil crusts that are interpreted as functioning as bare ground (therefore they would be susceptible to raindrop

splash erosion) should be recorded as bare ground.

---

5. **Number of gullies and erosion associated with gullies:** None to rare. Rare gullies may be present in landscape settings where increased runoff may accumulate (such as areas below exposed bedrock). Such gully development is expected to be limited to slopes exceeding 15% and adjacent to sites where runoff accumulation occurs (i.e. small watersheds, step sites, exposed bedrock, etc.). Any gullies present should show little sign of accelerated erosion and should be stabilized with perennial vegetation.

---

6. **Extent of wind scoured, blowouts and/or depositional areas:** Very minor evidence of wind generated soil movement. Wind scoured (blowouts) and depositional areas are rarely present.

---

7. **Amount of litter movement (describe size and distance expected to travel):** Most litter resides in place with some redistribution caused by water movement. Minor fine litter removal may occur in flow patterns and rills with deposition occurring at points of obstruction. The majority of litter accumulates at the base of plants. Some herbaceous litter may accumulate in soil depressions adjacent to plants. Woody litter is not likely to move.

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** This site should have a soil stability rating of 5 to 6 under vegetation canopies and a rating of 4 in the interspaces. The average should be a 5. Vegetation cover, litter, biological soil crusts and surface rock reduce erosion.

---

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Soil surface thickness typically varies from 2 to 3 inches. Structure varies from fine granular to thin platy. Color is typically light brown (10YR6/2 to yellowish red 5YR5/6). Organic matter is > 2%. An ochric horizon typically extends to a depth of 2 to 10 inches. The ochric horizon is a surface horizon lacking fine stratification and which is either light colored, or thin, or has a low organic carbon content, or is massive and (very) hard when dry. The A horizon would be expected to be more strongly developed under plant canopies. It is important if you are sampling to observe the A horizon under plant canopies as well as the interspaces. Refer to soil survey for more detailed information about your specific site.

---

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Vascular plants and well developed biological soil crusts will break raindrop impact and reduce splash erosion. Spatial distribution of vascular plants and interspaces between well developed biological soil crusts (where present) provide detention storage and surface roughness that slows runoff allowing time for infiltration. Interspaces between plants and well developed biological soil crusts (where present) may serve as water flow patterns during episodic runoff events, with natural erosion expected in severe storms. When perennial grasses decrease, reducing ground cover and increasing bare ground, runoff is expected to increase and any associated infiltration reduced.

---

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None. A few soils have bedrock at about 30+ inches. Naturally occurring soil horizons may be harder than the surface because of an accumulation of clay or calcium carbonate and should not be considered as compaction layers.

---

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: non-sprouting shrubs (Wyoming big sagebrush) > cool season perennial grasses (Indian ricegrass, Squirreltail) > warm season perennial grasses (Galleta, Blue grama)

Sub-dominant: sprouting shrubs (rabbitbrush) > forbs > biological soil crusts

Other: Functional/structural groups may appropriately contain non-native species if their ecological function is the same as the native species in the reference state (e.g. Crested wheatgrass and Russian wildrye etc.) Biological soil crust is variable in its expression where present on this site and is measured as a component of ground cover. Forbs can be expected to vary widely in their expression in the plant community based upon departures from average growing conditions.

Additional: Factors contributing to temporal variability include drought, insects, and fire. Assumed fire cycle of 50-100+ years. Spatial variability factors include slope, aspect, etc. Following a recent disturbance such as fire or drought that removes the woody vegetation, forbs and perennial grasses (herbaceous species) may dominate the community. If a disturbance has not occurred for an extended period of time, woody species may continue to increase crowding out the perennial herbaceous understory species. In either case, these conditions could reflect a functional community phase within the reference state.

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** During years with average to above-average precipitation, there should be very little recent mortality or decadence apparent in trees, shrubs, or grasses. During severe (multi-year) drought some sagebrush may die. There may be partial mortality of individual bunchgrasses and other shrubs during drought. Some bunchgrass and shrub mortality may occur during severe droughts, particularly on the shallower and coarser soils associated with this site.

---

14. **Average percent litter cover (%) and depth ( in):** Litter cover Average litter cover (including under plants) Depth should be 1 leaf thickness in the interspaces, up to ¼” under shrub. Litter redistribution following natural extreme runoff events can reduce litter cover by concentrating it in low-lying areas. Litter cover may increase by 5 to 10% followings seasons with high production of annuals.

---

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 210 - 430 #/acre on an average year

---

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Green rabbitbrush, Cheatgrass, Purple Threeawn, Broom snakeweed & introduced annual forbs (Filarie, Russian thistle).

---

17. **Perennial plant reproductive capability:** All perennial plants should have the ability to reproduce sexually or asexually in most years, except in drought years. Low green rabbitbrush sprouts vigorously following fire.

