

Ecological site R035XY221UT Semidesert Shallow Loam (Utah Juniper-Pinyon)

Accessed: 05/04/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Robert Stager (BLM), F.E. Busby (USU), Dana Truman (NRCS), Paul Curtis (BLM), Shane A. Green (NRCS)
Contact for lead author	shane.green@ut.usda.gov
Date	09/12/2008
Approved by	Shane A. Green
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- Number and extent of rills:** A. On more gentle slopes (< 15 %): Very few. Rills may be 10 or more feet in length. Rills are most likely to form below adjacent exposed bedrock or water flow patterns where sufficient water accumulates to cause erosion. B. On steep slopes (> 15 %): Few rills are present. Where they occur, rills may extend down entire slope.
- Presence of water flow patterns:** Some sinuous flow patterns wind around perennial plants and surface rock. Evidence of flow patterns is expected to increase somewhat with slopes greater than 15%. Water flow patterns are long (15-20 feet), narrow (<1 foot wide), and spaced widely (5-10 yards) on gentle slopes (<15%) and more closely (<5 yards) on steeper slopes (>15%).
- Number and height of erosional pedestals or terracettes:** Rare small pedestals may form at the base of plants that occur on the edge of water flow patterns, but should not show any exposed roots. Terracettes are fairly common, forming behind debris dams of small to medium sized litter (up to 2 inches in diameter) in water flow patterns. These debris dams may accumulate smaller litter (leaves, grass and forb stems) and sediment.
- Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 20 – 60 %. (Soil surface is typically covered 10 to 50 percent surface fragments). Most bare ground is associated with water flow patterns, rills, and gullies. Poorly developed biological soil crusts that are interpreted as functioning as bare ground (therefore they would be susceptible to raindrop splash erosion) should be recorded as bare ground. Ground cover is based on first raindrop impact, and bare ground is the opposite of ground cover. Ground cover

+ bare ground = 100%.

5. **Number of gullies and erosion associated with gullies:** None to rare on gentle slopes (< 15%). On steep slopes and areas below adjacent exposed bedrock, gullies may occur. Length often extends from exposed bedrock until gully reaches a stream or an area where water and sediment accumulate. Gullies may remove soil from base of shrubs and/or trees exposing roots. Gullies may show slightly more indication of erosion as slope increases, or as the site occurs adjacent to steep sites/watershed with concentrated flow patterns.
-
6. **Extent of wind scoured, blowouts and/or depositional areas:** None to very few. Trees break the wind and reduce the potential for wind erosion. The stones on the soil surface armor and reduces the potential for wind erosion.
-
7. **Amount of litter movement (describe size and distance expected to travel):** Most litter accumulates at the base of plants. Woody litter is usually not moved unless present in water flow patterns, rills, or gullies.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** This site should have an erosion rating of 4 or 5 under the plant canopies, and a rating of 4 in the interspaces. The average should be a 4. Vegetation cover, litter, biological soil crusts and surface rock reduce erosion.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Soil surface typically varies from 2 to 4 inches. Structure typically varies from fine granular to subangular blocky. Color typically varies from brown (10YR5/3) to pale brown (10YR6/3). The A horizon characteristics are the same under plant canopies or in the interspaces. Refer to soil survey for more detailed information about your specific site.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Spatial distribution of well developed biological soil crusts (where present) intercept raindrops reducing splash erosion and provide areas of surface detention to store water allowing additional time for infiltration. Crowns of trees and accumulating litter at base of trees appear to create a micro-topography that may enhance development of water flow patterns below the drip line of the canopy. Significant increases in Pinyon-juniper canopy (beyond the reference state) reduces understory vegetation causing an associated increase in runoff.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None, although bedrock is found from 4 to 20 inches of soil surface. In addition, there may be layers of calcium carbonate or other naturally occurring hard layers found in the soil subsurface. These should not be considered to be compaction layers.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: trees (Juniper > Pinion) > sprouting shrubs (Buffaloberry, Mormontea, Rabbitbrush)

Sub-dominant: warm season perennial grasses (Galleta) = cool season perennial grasses (Indian ricegrass) = non sprouting shrubs (Cliffrose, Snakeweed) > forbs > Biological soil crusts

Other: Functional/structural groups may appropriately contain non-native species if their ecological function is the same as the native species in the reference state (e.g. Crested wheatgrass, Smooth brome, Intermediate wheatgrass, Siberian wheatgrass and/or forage kochia etc.) Biological soil crust is variable in its expression where present on this site and is measured as a component of ground cover. Forbs can be expected to vary widely in their expression in the plant community based upon departures from average growing conditions.

Additional: Factors contributing to temporal variability include insects and other pathogens (mistletoe), drought, extreme precipitation events, etc. Factors contributing to spatial variability include slope, amount of rock fragments, aspect, etc. Following a recent disturbance such as fire, drought or insects that may remove the woody vegetation, forbs and perennial grasses (herbaceous species) may become more dominant in the community. These conditions may reflect a functional community phase within the reference state.

-
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** During years with average to above-average precipitation, there should be very little recent mortality or decadence apparent in trees, shrubs, or grasses. During severe (multi-year) drought up to 20% of the junipers may die, either from drought or pathogens such as mistletoe. There may be partial mortality of individual bunchgrasses and other shrubs during drought. Some bunchgrass and shrub mortality may occur during severe droughts, particularly on the shallower and coarser soils associated with this site. Because woody stems may persist for many years, juniper (especially older trees) will normally have dead stems within the plant canopy.
-
14. **Average percent litter cover (%) and depth (in):** Litter cover (including under plants) nearly all of which should be fine litter. Depth should be 1 leaf thickness in the interspaces and up to ¼" under canopies, and up to ¾" under tree canopies. Litter cover may increase to 30% on some years due to increased production of plants.
-
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 240-580 #/acre on an average year
-
16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Few invasives capable of dominating this site. Cheatgrass, Broom snakeweed, and Mustard may invade the community.
-
17. **Perennial plant reproductive capability:** All perennial plants should have the ability to reproduce sexually or asexually in most years, except in drought years.
-