

## Ecological site R035XY260UT Semidesert Very Steep Stony Loam (Salina Wildrye)

Accessed: 05/03/2024

### Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Robert Stager (BLM), Dana Truman (NRCS), Paul Curtis (BLM), Shane A. Green (NRCS), Randy Beckstrand (BLM)
Contact for lead author	shane.green@ut.usda.gov
Date	01/30/2007
Approved by	Shane A. Green
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

### Indicators

- Number and extent of rills:** Few. Due to the surface rock fragments on this site, traditional rill identification is reduced. The overall gravelly to channery surface is expected to be resistant to rill formation and accelerated erosion in general. Where rills occur, they may extend down entire slope.
- Presence of water flow patterns:** Common- flow patterns can be very sinuous and wind around rock fragments and perennial plant bases. They show some evidence of erosion with fines and litter depositing against the uphill side of gravel, rocks and plants. During precipitation events e.g. thunderstorms, these sites are expected to shed large volumes of water to adjacent ecological sites.
- Number and height of erosional pedestals or terracettes:** Pedestals are common. Without protection from surface gravels, plants will become slightly pedestaled. Pedestals may form at the base of plants that occur on the edge of primary flow patterns and rills, but there should be no exposed roots. Interspaces between any well developed biological soil crusts resemble pedestals and may be up to 2 inches high. Terracettes are common due to slope, terracettes form throughout the site. Debris dams of small to medium sized litter (up to 2 inches in diameter) may form in water flow patterns, rills, and gullies. These debris dams may accumulate herbaceous litter.
- Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 10 – 25 %. Most bare ground is associated with water flow patterns, rills, and gullies. Soil is covered by 40 to 80 percent rock fragments. Any areas with well developed biological soil crusts should not be counted as bare

ground. Poorly developed biological soil crusts that are interpreted as functioning as bare ground (therefore they would be susceptible to raindrop splash erosion) should be recorded as bare ground. Ground cover is based on first raindrop impact, and bare ground is the opposite of ground cover.

---

5. **Number of gullies and erosion associated with gullies:** Few - Length often extends from exposed bedrock until gully reaches a stream or an area where water and sediment accumulate, but they may be wide and shallow and armored with very large rocks.
6. **Extent of wind scoured, blowouts and/or depositional areas:** None to very few.
7. **Amount of litter movement (describe size and distance expected to travel):** Down slope redistribution of any incident litter caused by water is expected. Deposition would likely occur at points of obstruction such as the uphill side of gravel, rocks and plants, especially following major storm events. Fine litter is moved with even moderate precipitation events and spring runoff. Woody litter may be washed from site. Gullies may remove accumulated litter from under trees.
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** 80 to 90% of this site should have an erosion rating of 4 or 5. 10 to 20% may have a rating of 3 to 4. The average should be a 4. Surface texture is gravelly loam to very channery sandy loam. Vegetation cover, litter, biological soil crusts and surface rock reduce erosion.
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Soil surface is typically 3 inches deep. Structure is typically weak thick platy parting to weak fine subangular blocky. Color is typically strong brown (7.5YR5/6). The A horizon would be expected to be more strongly developed under plant canopies. It is important if you are sampling to observe the A horizon under plant canopies as well as the interspaces. Use the specific information for the soil you are assessing found in the published soil survey to supplement this description.
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Spatial distribution of plants and/or well developed biological soil crusts (where present) intercept raindrops, somewhat reducing splash erosion and provide areas of surface detention to store water allowing additional time for infiltration. (surface rocks influence the runoff and water retention).
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None. There may be layers of calcium carbonate, gravel, cobbles or other naturally occurring hard layers found in the soil subsurface. These should not be considered to be compaction layers.
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: examples: Indian ricegrass, Salina Wildrye, Shadscale.

Sub-dominant: examples: Mormontea, Bigelow sagebrush, forbs

Other: Perennial and annual forbs can be expected to vary widely in their expression in the plant community based upon departures from average growing conditions.

Additional: Disturbance regime includes infrequent fire, drought, and insects.

Dominance by average annual production: Cool season perennial (bunch and rhizomatous) grasses = Non-sprouting shrubs > Sprouting shrubs > forbs > Warm season perennial grasses. Functional/structural groups may appropriately contain non-native species if their ecological function is the same as the native species in the reference state (e.g. Crested wheatgrass and Russian wildrye etc.)

Biological soil crust is variable in its expression where present on this site and is measured as a component of ground cover.

Following a recent disturbance such as fire or drought that removes the woody vegetation, forbs and perennial grasses (herbaceous species) may dominate the community. These conditions would reflect a functional community phase within the reference state.

- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** In general, a mix of age classes may be expected with some dead and decadent plants present.
- 

14. **Average percent litter cover (%) and depth ( in):** Litter cover 1-5%. Variability may occur due to weather. Litter cover is generally low and only accumulates beneath mature plants at depths of less than 1 inch.
- 

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 150-530 lbs./ac.
- 

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Possible invaders or increasers on this site are cheatgrass, halogeton, locoweed, and broom snakeweed.
- 

17. **Perennial plant reproductive capability:** All perennial plants should have the ability to reproduce sexually or asexually in most years, except in drought years.
-