

Ecological site R035XY263UT Semidesert Very Steep Stony Loam (Two-Needle Pinyon, Utah Juniper)

Accessed: 06/03/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Jacob Owens (NRCS), Shane Green (NRCS)
Contact for lead author	shane.green@ut.usda.gov
Date	12/07/2009
Approved by	Shane A. Green
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills:** Very few. Due to the surface rock fragments on this site, traditional rill formation is reduced. The overall gravelly to channery surface is expected to be resistant to rill formation and accelerated erosion in general. Where rills do occur, they may extend down entire slope.
- 2. Presence of water flow patterns:** Due to the steep slopes, flow patterns are present and tend to be very sinuous and wind around rock fragments and perennial plant bases. They show some evidence of erosion with fines and litter depositing against the uphill side of gravel, rocks and plants. During episodic precipitation events e.g. thunderstorms, these sites are expected to shed large volumes of water to adjacent ecological sites.
- 3. Number and height of erosional pedestals or terracettes:** Pedestals may form at the base of plants that occur on the edge of primary flow patterns and rills. Interspaces between any well developed biological soil crusts resemble pedestals and may be up to 2 inches high. Terracettes are present. Debris dams of small to medium sized litter (up to 2 inches in diameter) may form in water flow patterns, rills, and gullies. These debris dams may accumulate smaller litter (leaves, grass and forb stems).
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 8 – 30 %. Most bare ground is associated with water flow patterns, rills, and gullies. Soil is covered by up to 60 percent rock fragments. Any areas with well developed biological soil crusts should not be counted as bare ground. Poorly developed biological soil crusts that are interpreted as functioning as bare ground (therefore they would be susceptible to raindrop splash erosion) should be recorded as bare ground. Ground cover is based on first raindrop

impact, and bare ground is the opposite of ground cover.

5. **Number of gullies and erosion associated with gullies:** Few gullies may be present. Length often extends from exposed bedrock until gully reaches a stream or an area where water and sediment accumulate, but they may be wide and shallow and armored with very large rocks.

6. **Extent of wind scoured, blowouts and/or depositional areas:** None.

7. **Amount of litter movement (describe size and distance expected to travel):** Due to the steepness of slope being between 30 to 70 percent, down slope redistribution of any incident litter caused by water is expected. Deposition would likely occur at points of obstruction such as the uphill side of gravel, rocks and plants, especially following major storm events. Fine litter is moved with even moderate precipitation events and spring runoff. Woody stems may be washed from site. Gullies may remove accumulated litter from under trees.

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** 80 to 90% of this site should have an erosion rating of 4 or 5. 10 to 20% may have a rating of 3 to 4. The average should be a 4. Surface texture is gravelly loam to very channery sandy loam. Vegetation cover, litter, biological soil crusts and surface rock reduce erosion.

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Soil surface is 3 inches deep. Structure is weak fine platy to moderate fine and medium granular. Color is reddish brown (7.5YR5/4). The A horizon would be expected to be more strongly developed under plant canopies. It is important if you are sampling to observe the A horizon under plant canopies as well as the interspaces. Use the specific information for the soil you are assessing found in the published soil survey to supplement this description.

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Spatial distribution of plants and/or well developed biological soil crusts (where present) intercept raindrops reducing splash erosion and provide areas of surface detention to store water allowing additional time for infiltration. Crowns of trees and accumulating litter at base of trees appear to create a micro-topography that may enhance development of water flow patterns below the drip line of the canopy. Significant increases in Pinyon-juniper canopy (beyond the reference state) reduces understory vegetation causing an associated increase in runoff.

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None. There may be layers of calcium carbonate, gravel, cobbles or other naturally occurring hard layers found in the soil subsurface. These should not be considered to be compaction layers.

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Dominance by average annual production: Trees (Juniper > Pinion) > Sprouting shrubs > Cool season perennial grasses > Warm season perennial grasses > forbs. Functional/structural groups may appropriately contain

non-native species if their ecological function is the same as the native species in the reference state (e.g. Crested wheatgrass, Intermediate wheatgrass, etc.)

Sub-dominant:

Other: Biological soil crust is variable in it's expression where present on this site and is measured as a component of ground cover.

Additional: Following a recent disturbance such as fire, drought, or insects that removes the woody vegetation, forbs and perennial grasses (herbaceous species) may dominate the community. If a disturbance has not occurred for an extended period of time, woody species may continue to increase crowding out the perennial herbaceous understory species. In either case, these conditions would reflect a functional community phase within the reference state.

Dominants— Utah juniper, two-needle pinion, Bigelow's sagebrush, mormon tea, and Indian ricegrass. Sub Dominants— Broom snakeweed, Single leaf ash, roundleaf buffalo berry, James galleta. Perennial and annual forbs can be expected to vary widely in their expression in the plant community based upon departures from average growing conditions.

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** In general, a mix of age classes may be expected with a few dead and decadent plants present.
-

14. **Average percent litter cover (%) and depth (in):** Litter cover 2-5%. Variability may occur due to weather.
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 150-320 #/acre on an average year.
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Possible invaders or increasers on this site are cheatgrass, sunflower, and halogeton, locoweed, broom snakeweed and juniper.
-

17. **Perennial plant reproductive capability:** All perennial plants should have the ability to reproduce sexually or asexually in most years, except in drought years.
-