

## Ecological site R035XY264UT Semidesert Gypsum (Torrey's Jointfir)

Accessed: 05/19/2024

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

### MLRA notes

Major Land Resource Area (MLRA): 035X—Colorado Plateau

This site occurs on moderately deep to deep, gypsum-affected soils in the semidesert zone of MLRA D35 (Colorado and Green River Plateaus). It receives 7 to 10 inches of annual precipitation, most notably as convective thunderstorms from July through October. It most commonly occurs on dipslopes ofuestas hogbacks. Slope is usually less than 5-60% and elevations range from 3800-5400 feet. The reference plant community is sparsely vegetated and dominated by Torrey's jointfir, shadscale, Bigelow sagebrush, and diverse perennial grasses. The reference plant community is highly resistant to change due to its inability to carry fire, along with a harsh soil environment that resists invasion and dominance by other species. Lichen crusts can dominate the shrub interspaces in lieu of grass.

### Classification relationships

Modal Soil: Raplee VFSL — coarse-loamy, gypsic, mesic Typic Torriorthents

Type Location: West of Comb Wash between HWY. 163 and the Mormon Trail.

### Similar sites

R035XY237UT	<b>Semidesert Shallow Gypsum (Mormontea)</b> This site is also shallow, but it receives greater than 9 inches of annual precipitation. Plant community composition is similar, but the higher precipitation results in 20-40% more production.
R035XY142UT	<b>Desert Very Shallow Gypsum (Torrey's Jointfir)</b> This site is very shallow and has very high amounts of gypsum. Composition is similar, but production is about half.
R035XY106UT	<b>Desert Gypsum Loam (Torrey's Jointfir)</b> This site has soils greater than 20 inches deep and therefore supports more perennial grass. It is more resilient against drought due to higher water holding capacity.
R035XY126UT	<b>Desert Shallow Gypsum (Torrey's Jointfir)</b> This site has shallow soils and an aridic soil moisture regime. It produces a similar plant community, but with about 40% less production.

**Table 1. Dominant plant species**

Tree	Not specified
Shrub	(1) <i>Ephedra torreyana</i> (2) <i>Atriplex confertifolia</i>
Herbaceous	(1) <i>Pleuraphis jamesii</i> (2) <i>Achnatherum hymenoides</i>

## Physiographic features

This site occurs on dipslopes of cuestras and hogbacks, and can occur in flatter areas where soils are deep and gypsum-affected. Slopes are 5-65% and elevations range from 3800-5400 feet.

**Table 2. Representative physiographic features**

Landforms	(1) Dip slope
Flooding frequency	None
Ponding frequency	None
Elevation	1,158–1,646 m
Slope	5–65%
Aspect	Aspect is not a significant factor

## Climatic features

This site is hot in the summer and cool in the winter. Average annual precipitation is 7 to 9 inches, with about half of the precipitation occurring as convection thunderstorms from July through October. June is typically the driest month during the growing season. Precipitation is extremely variable from month to month and from year to year. Large fluctuations in daily temperatures are also common.

Modeled climate data (PRISM) was used to describe the climate for map units correlated to this site.

**Table 3. Representative climatic features**

Frost-free period (average)	160 days
Freeze-free period (average)	190 days
Precipitation total (average)	229 mm

## Influencing water features

Due to its landscape position, this site is not typically influenced by streams or wetlands. Ephemeral washes may

cross this site, but these washes only carry water during intense storms. As a result, production may increase and composition may differ near washes, but they do not support riparian-obligate vegetation.

## Soil features

Soils for this site are moderately deep to deep and gypsum-affected. They are typically well-drained, fine sandy loams with moderately rapid permeability. These soils formed in residuum or slope alluvium derived from sandstone material high in gypsum. Rock fragments are not abundant on the soil surface or throughout the profile. The soil moisture regime is ustic aridic and the soil temperature regime is mesic. Water-holding capacity ranges from 2 to 5 inches of water in the upper 40 inches of soil.

This site has been correlated to soils in the following soil surveys:

UT685 - Capitol Reef - Lemrac;

**Table 4. Representative soil features**

Parent material	(1) Residuum—rock gypsum
Surface texture	(1) Very fine sandy loam
Family particle size	(1) Loamy
Drainage class	Well drained
Permeability class	Moderately rapid
Soil depth	51–152 cm
Surface fragment cover ≤3"	0%
Surface fragment cover >3"	0–5%
Available water capacity (0–101.6cm)	5.08–12.7 cm
Calcium carbonate equivalent (0–101.6cm)	1–10%
Electrical conductivity (0–101.6cm)	0–4 mmhos/cm
Sodium adsorption ratio (0–101.6cm)	0
Soil reaction (1:1 water) (0–101.6cm)	7.9–8.4
Subsurface fragment volume ≤3" (Depth not specified)	0–16%
Subsurface fragment volume >3" (Depth not specified)	0–6%

## Ecological dynamics

This site developed under Colorado Plateau ecological conditions and the natural influences of herbivory and climate. This site's plant species composition is generally dominated by Torrey's jointfir, shadscale, Bigelow sagebrush, and diverse perennial grasses. In some areas, lichen crusts stabilize the soil and preclude grass production.

There is no evidence that this site historically burned on a regular basis due to very large and persistent gaps between plants. Invasive species, particularly Russian thistle, are capable of establishing on this site, but they have not been documented as gaining dominance.

This ecological site has been grazed by domestic livestock since they were first introduced into the area around 1860. The site is highly resistant to grazing due to the low palatability of Torrey's jointfir, and the general lack of

forage plants. The introduction of domestic livestock and the use of fencing and reliable water sources have therefore only minimally influenced the historic disturbance regime associated with this ecological site. Still, livestock-related disturbance may facilitate establishment of non-native species.

Suitability for rangeland seeding is very poor because of low annual precipitation, harsh soil conditions, and low available water capacity.

The following State and Transition diagram shows the reference plant community. No other plant communities have been documented on this site to date. As more data are collected, new plant communities or states may be added. This model was developed using range data collected over the last 40 years in MLRA D35 in southeastern Utah. Both ocular and measured data was collected and utilized.

## State and transition model

### R035XY264UT Semidesert Gypsum (Torrey's Jointfir)




Figure 4. State-and-Transition Model

## State 1 Reference State

The reference state consists of a diverse shrub canopy dominated by Torrey's jointfir. The understory is usually dominated by a diverse perennial grass community, but can sometimes be dominated by lichen crusts. Lichen crusts tend to dominate in areas with relatively harsh soil conditions (less deep, less infiltration, more gypsum, etc) that are less likely to be disturbed (more isolated, more steep, etc). The exact dynamics between grass and lichen communities are poorly understood. This state is susceptible to non-native invasive species establishment. Disturbances such as livestock grazing and recreation can increase the likelihood of invasion by promoting germination sites and/or seed sources for non-native species. However, Russian thistle is capable of establishing on this site in the absence of disturbance.

## Community 1.1

Torrey's jointfir Shrubland with perennial grasses.



R035XY264UT—Semidesert Gypsum (Torrey's jointfir) community 1.1—Torrey's jointfir Shrubland with

Figure 5. Phase 1.1

This plant community phase is dominated by Torrey's jointfir, shadscale, Bigelow Sagebrush, and diverse perennial grasses. Composition by air-dry weight is 5-15% forbs, 40-60% Grasses, and 35-55% shrubs.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	56	140	196
Shrub/Vine	84	140	196
Forb	11	28	45
Total	151	308	437

Table 6. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	8-15%
Grass/grasslike foliar cover	5-15%
Forb foliar cover	0-5%
Non-vascular plants	0%
Biological crusts	0-30%
Litter	5-15%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0-5%
Bedrock	0%
Water	0%
Bare ground	30-60%

Table 7. Canopy structure (% cover)

Height Above Ground (M)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.15	—	0-5%	0-5%	0-4%
>0.15 <= 0.3	—	0-10%	0-10%	0-5%
>0.3 <= 0.6	—	5-15%	0-10%	0-2%
>0.6 <= 1.4	—	0-5%	—	—
>1.4 <= 4	—	—	—	—
>4 <= 12	—	—	—	—
>12 <= 24	—	—	—	—
>24 <= 37	—	—	—	—
>37	—	—	—	—

## Community 1.2 Torrey's jointfir Shrubland.



R035XY264UT—Semidesert Gypsum (Torrey's jointfir) community 1.2—Torrey's jointfir Shrubland. Cover is

Figure 7. 1.2

This plant community phase is dominated by Torrey's jointfir, and other shrubs. Grasses are limited or absent from the community, while lichen crusts dominate the soil surface.

Table 8. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Shrub/Vine	84	140	196
Forb	—	11	28
Grass/Grasslike	—	11	28
<b>Total</b>	<b>84</b>	<b>162</b>	<b>252</b>

Table 9. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	12-20%
Grass/grasslike foliar cover	0-2%
Forb foliar cover	0-2%
Non-vascular plants	0%
Biological crusts	40-60%
Litter	5-10%

Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0-5%
Bedrock	0%
Water	0%
Bare ground	10-30%

**Table 10. Canopy structure (% cover)**

Height Above Ground (M)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.15	—	0-5%	0-1%	0-2%
>0.15 <= 0.3	—	0-10%	0-2%	0-2%
>0.3 <= 0.6	—	8-20%	0-2%	—
>0.6 <= 1.4	—	0-5%	—	—
>1.4 <= 4	—	—	—	—
>4 <= 12	—	—	—	—
>12 <= 24	—	—	—	—
>24 <= 37	—	—	—	—
>37	—	—	—	—

## Pathway 1.1a Community 1.1 to 1.2



**Torrey's jointfir shrubland with perennial grasses.**



**Torrey's jointfir shrubland.**

This pathway occurs when climatic events, such as extended drought reduce the basal cover of perennial grasses for a long periods of time. When coupled with a lack of surface disturbance, it is expected that lichen crust will increase and preclude the re-establishment of perennial grasses.

## Pathway 1.2A Community 1.2 to 1.1



**Torrey's jointfir shrubland.**



**Torrey's jointfir shrubland with perennial grasses.**

Wet periods combined with soil surface disturbance facilitate this community pathway. Seed source for perennial grasses must also be available.

## State 2 Invaded State

The invaded state resembles the reference state in both community structure and function, but non-native species, notably Russian thistle, are present. As a result, the resilience of the state is somewhat reduced and the possibility of further degradation is greater.



## Community 2.1

### Torrey's jointfir / Perennial grasses



R035XY264UT—Semidesert Gypsum (Torrey's jointfir) community 2.1—Invaded Torrey's jointfir Shrubland

**Figure 9. Phase 2.1**

This plant community is similar to Reference State Community 1.1. except that invasive species are now present. Dominant species are Torrey's jointfir, shadscale, Bigelow sagebrush, James' galleta and Indian ricegrass.

**Table 11. Annual production by plant type**

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	56	140	196
Shrub/Vine	84	140	196
Forb	11	28	45
<b>Total</b>	<b>151</b>	<b>308</b>	<b>437</b>

**Table 12. Ground cover**

Tree foliar cover	0%
Shrub/vine/liana foliar cover	8-15%
Grass/grasslike foliar cover	5-15%
Forb foliar cover	0-5%
Non-vascular plants	0%
Biological crusts	0-30%
Litter	5-15%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0-5%
Bedrock	0%
Water	0%
Bare ground	30-60%

**Table 13. Canopy structure (% cover)**



Height Above Ground (M)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.15	—	0-5%	0-5%	0-4%
>0.15 <= 0.3	—	0-10%	0-10%	0-5%
>0.3 <= 0.6	—	5-15%	0-10%	0-2%
>0.6 <= 1.4	—	0-5%	—	—
>1.4 <= 4	—	—	—	—
>4 <= 12	—	—	—	—
>12 <= 24	—	—	—	—
>24 <= 37	—	—	—	—
>37	—	—	—	—

## Community 2.2

### Torrey's jointfir with invasives

This plant community is similar to Reference State Community 1.2 except that invasive species are now present. Perennial grasses are greatly reduced, and Russian thistle or other invasive annuals are present. This phase may produce annuals, but it is still dominated by lichen crusts and Torrey's jointfir.

**Table 14. Annual production by plant type**

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Shrub/Vine	84	140	196
Forb	6	17	28
Grass/Grasslike	—	11	28
<b>Total</b>	<b>90</b>	<b>168</b>	<b>252</b>

**Table 15. Ground cover**

Tree foliar cover	0%
Shrub/vine/liana foliar cover	12-20%
Grass/grasslike foliar cover	0-2%
Forb foliar cover	0-5%
Non-vascular plants	0%
Biological crusts	40-60%
Litter	5-10%
Surface fragments >0.25" and <=3"	0%
Surface fragments >3"	0-5%
Bedrock	0%
Water	0%
Bare ground	10-30%

**Table 16. Canopy structure (% cover)**

Height Above Ground (M)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.15	—	0-5%	0-1%	0-2%
>0.15 <= 0.3	—	0-10%	0-5%	0-2%
>0.3 <= 0.6	—	8-20%	0-2%	—
>0.6 <= 1.4	—	0-5%	—	—
>1.4 <= 4	—	—	—	—
>4 <= 12	—	—	—	—
>12 <= 24	—	—	—	—
>24 <= 37	—	—	—	—
>37	—	—	—	—

## Pathway 2.1a

### Community 2.1 to 2.2

This pathway occurs when climatic events, such as extended drought reduce the basal cover of perennial grasses for a long periods of time. When coupled with a lack of surface disturbance, it is expected that lichen crust will increase and preclude the re-establishment of perennial grasses.

## Pathway 2.2A

### Community 2.2 to 2.1

Wet periods combined with soil surface disturbance facilitate this community pathway. Seed source for perennial grasses must also be available.

## Transition T1A

### State 1 to 2

This transition occurs with the establishment of non-native invasive species. Disturbances that promote this transition include season long continuous grazing of perennial grasses, prolonged drought, recreation or other surface disturbances. However, invasive species such as Russian thistle can invade intact perennial plant communities with little to no disturbance. Once invasive plants are found in the plant community, a return to the reference state is not likely.

## Additional community tables

Table 17. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
<b>Shrub/Vine</b>					
0	<b>Dominant Shrubs</b>			84–168	
	Torrey's jointfir	EPTO	<i>Ephedra torreyana</i>	34–78	3–8
	shadscale saltbush	ATCO	<i>Atriplex confertifolia</i>	17–56	1–5
	Bigelow sage	ARBI3	<i>Artemisia bigelovii</i>	17–45	1–3
3	<b>Sub-dominant Shrubs</b>			0–56	
	broom snakeweed	GUSA2	<i>Gutierrezia sarothrae</i>	0–28	0–2
	Shrub (>.5m)	2SHRUB	<i>Shrub (&gt;.5m)</i>	0–22	0–2
	crispleaf buckwheat	ERCOA	<i>Eriogonum corymbosum var. aureum</i>	0–22	0–2
	fourwing saltbush	ATCA2	<i>Atriplex canescens</i>	0–11	0–1
	yellow rabbitbrush	CHVI8	<i>Chrysothamnus viscidiflorus</i>	0–6	0–1

	Utah juniper	JUOS	<i>Juniperus osteosperma</i>	0–6	0–1
	winterfat	KRLA2	<i>Krascheninnikovia lanata</i>	0–6	0–1
	narrowleaf yucca	YUAN2	<i>Yucca angustissima</i>	0–6	0–1
	plains pricklypear	OPPO	<i>Opuntia polyacantha</i>	0–3	0–1
<b>Grass/Grasslike</b>					
0	<b>Dominant Grasses</b>			34–112	
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	17–56	1–5
	James' galleta	PLJA	<i>Pleuraphis jamesii</i>	17–56	1–5
1	<b>Sub-dominant Grasses</b>			22–112	
	needle and thread	HECO26	<i>Hesperostipa comata</i>	0–56	0–5
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	0–45	0–4
	alkali sacaton	SPAI	<i>Sporobolus airoides</i>	0–22	0–2
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	0–22	0–2
	desert needlegrass	ACSP12	<i>Achnatherum speciosum</i>	0–11	0–1
	Grass, annual	2GA	<i>Grass, annual</i>	0–6	0–1
	Grass, perennial	2GP	<i>Grass, perennial</i>	0–6	0–1
	sandhill muhly	MUPU2	<i>Muhlenbergia pungens</i>	0–3	0–1
<b>Forb</b>					
2	<b>Forbs</b>			11–45	
	rayless tansyaster	MAGR2	<i>Machaeranthera grindelioides</i>	0–22	0–2
	Forb, perennial	2FP	<i>Forb, perennial</i>	0–22	0–2
	roughseed cryptantha	CRFL6	<i>Cryptantha flavoculata</i>	0–11	0–1
	Forb, annual	2FA	<i>Forb, annual</i>	0–11	0–1
	Rocky Mountain stickweed	CLPA3	<i>Cleomella palmeriana</i>	0–6	0–1
	cleftleaf wildheliotrope	PHCR	<i>Phacelia crenulata</i>	0–6	0–1
	scarlet globemallow	SPCO	<i>Sphaeralcea coccinea</i>	0–6	0–1
	stemless four-nerve daisy	TEACA2	<i>Tetrameuris acaulis</i> var. <i>acaulis</i>	0–3	0–1
	Jones' pepperweed	LEMOJ	<i>Lepidium montanum</i> var. <i>jonesii</i>	0–3	0–1
	Wright's bird's beak	COWR2	<i>Cordylanthus wrightii</i>	0–2	0–1
	tufted evening primrose	OECA10	<i>Oenothera caespitosa</i>	0–1	0–1

Table 18. Community 1.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
<b>Shrub/Vine</b>					
0	<b>Dominant Shrubs</b>			84–168	
	Torrey's jointfir	EPTO	<i>Ephedra torreyana</i>	34–78	3–8
	shadscale saltbush	ATCO	<i>Atriplex confertifolia</i>	11–56	1–5
	Bigelow sage	ARBI3	<i>Artemisia bigelovii</i>	11–45	1–4
	James' galleta	PLJA	<i>Pleuraphis jamesii</i>	0–22	–
	matted crinklemat	TILA6	<i>Tiquilia latior</i>	3–11	–
	fanleaf hawthorn	CRFL	<i>Crataegus flabellata</i>	1–6	–
	scarlet globemallow	SPCO	<i>Sphaeralcea coccinea</i>	1–6	–
3	<b>Sub-dominant Shrubs</b>			0–56	
	Shrub (>.5m)	2SHRUB	<i>Shrub (&gt;.5m)</i>	6–22	1–2
	yellow rabbitbrush	CHVI8	<i>Chrysothamnus viscidiflorus</i>	0–17	0–2
	fourwing saltbush	ATCA2	<i>Atriplex canescens</i>	0–11	0–1
	crispleaf buckwheat	ERCOA	<i>Eriogonum corymbosum</i> var. <i>aureum</i>	0–6	0–1
	broom snakeweed	GUSA2	<i>Gutierrezia sarothrae</i>	0–6	0–1
	winterfat	KRLA2	<i>Krascheninnikovia lanata</i>	0–6	0–1
	plains pricklypear	OPPO	<i>Opuntia polyacantha</i>	0–3	0–1
	narrowleaf yucca	YUAN2	<i>Yucca angustissima</i>	0–2	0–1
<b>Grass/Grasslike</b>					
1	<b>Grasses</b>			0–28	
	James' galleta	PLJA	<i>Pleuraphis jamesii</i>	2–22	0–2
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	0–11	0–1
	desert needlegrass	ACSP12	<i>Achnatherum speciosum</i>	0–6	0–1
	Grass, annual	2GA	<i>Grass, annual</i>	0–6	0–1
	Grass, perennial	2GP	<i>Grass, perennial</i>	0–6	0–1
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	0–6	0–1
	alkali sacaton	SPAI	<i>Sporobolus airoides</i>	0–3	0–1
	needle and thread	HECO26	<i>Hesperostipa comata</i>	0–3	0–1
	sandhill muhly	MUPU2	<i>Muhlenbergia pungens</i>	0–3	0–1
<b>Forb</b>					
2	<b>Forbs</b>			6–28	
	scarlet globemallow	SPCO	<i>Sphaeralcea coccinea</i>	0–6	0–1
	rayless tansyaster	MAGR2	<i>Machaeranthera grindelioides</i>	0–6	0–1
	Forb, annual	2FA	<i>Forb, annual</i>	0–6	0–1
	Forb, perennial	2FP	<i>Forb, perennial</i>	0–6	0–1
	Wright's bird's beak	COWR2	<i>Cordylanthus wrightii</i>	0–6	0–1
	roughseed cryptantha	CRFL6	<i>Cryptantha flavoculata</i>	0–6	0–1
	Jones' pepperweed	LEMOJ	<i>Lepidium montanum</i> var. <i>jonesii</i>	0–3	0–1
	stemless four-nerve daisy	TEACA2	<i>Tetraneuris acaulis</i> var. <i>acaulis</i>	0–3	0–1
	tufted evening primrose	OECA10	<i>Oenothera caespitosa</i>	0–1	0–1

Table 19. Community 2.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
<b>Shrub/Vine</b>					
0	<b>Dominant Shrubs</b>			84–168	
	Torrey's jointfir	EPTO	<i>Ephedra torreyana</i>	34–78	3–8
	shadscale saltbush	ATCO	<i>Atriplex confertifolia</i>	11–56	1–5
	Bigelow sage	ARBI3	<i>Artemisia bigelovii</i>	11–45	1–4
3	<b>Sub-dominant Shrubs</b>			0–56	
	broom snakeweed	GUSA2	<i>Gutierrezia sarothrae</i>	0–28	0–2
	Shrub (>.5m)	2SHRUB	<i>Shrub (&gt;.5m)</i>	0–22	0–2
	crispleaf buckwheat	ERCO14	<i>Eriogonum corymbosum</i>	0–22	0–2
	yellow rabbitbrush	CHVI8	<i>Chrysothamnus viscidiflorus</i>	0–17	0–2
	fourwing saltbush	ATCA2	<i>Atriplex canescens</i>	0–11	0–1
	plains pricklypear	OPPO	<i>Opuntia polyacantha</i>	0–3	0–1
	narrowleaf yucca	YUAN2	<i>Yucca angustissima</i>	0–2	0–1
<b>Grass/Grasslike</b>					
0	<b>Dominant Grasses</b>			34–112	
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	17–56	1–5
	James' galleta	PLJA	<i>Pleuraphis jamesii</i>	17–56	1–5
1	<b>Subdominant Grasses</b>			22–112	
	needle and thread	HECOC8	<i>Hesperostipa comata ssp. comata</i>	0–56	0–5
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	0–45	0–4
	alkali sacaton	SPAI	<i>Sporobolus airoides</i>	0–22	0–2
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	0–22	0–2
	desert needlegrass	ACSP12	<i>Achnatherum speciosum</i>	0–11	0–1
	Grass, annual	2GA	<i>Grass, annual</i>	0–6	0–1
	Grass, perennial	2GP	<i>Grass, perennial</i>	0–6	0–1
	sandhill muhly	MUPU2	<i>Muhlenbergia pungens</i>	0–3	0–1
<b>Forb</b>					
2	<b>Forbs</b>			11–45	
	rayless tansyaster	MAGR2	<i>Machaeranthera grindelioides</i>	0–22	0–2
	prickly Russian thistle	SATR12	<i>Salsola tragus</i>	0–17	0–1
	scarlet globemallow	SPCO	<i>Sphaeralcea coccinea</i>	0–6	0–1
	desert princesplume	STPI	<i>Stanleya pinnata</i>	0–6	0–1
	Forb, annual	2FA	<i>Forb, annual</i>	0–6	0–1
	Forb, perennial	2FP	<i>Forb, perennial</i>	0–6	0–1
	cleftleaf wildheliotrope	PHCR	<i>Phacelia crenulata</i>	0–6	0–1
	Wright's bird's beak	COWR2	<i>Cordylanthus wrightii</i>	0–3	0–1
	roughseed cryptantha	CRFL6	<i>Cryptantha flavoculata</i>	0–3	0–1
	Jones' pepperweed	LEMOJ	<i>Lepidium montanum var. jonesii</i>	0–3	0–1
	stemless four-nerve daisy	TEAC	<i>Tetraneuris acaulis</i>	0–3	0–1
	tufted evening primrose	OECA10	<i>Oenothera caespitosa</i>	0–1	0–1

Table 20. Community 2.2 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
<b>Shrub/Vine</b>					
0	<b>Dominant Shrubs</b>			84–140	
	Torrey's jointfir	EPTO	<i>Ephedra torreyana</i>	34–78	3–8
	shadscale saltbush	ATCO	<i>Atriplex confertifolia</i>	11–56	1–5
	Bigelow sage	ARBI3	<i>Artemisia bigelovii</i>	11–45	0–4
	cheatgrass	BRTE	<i>Bromus tectorum</i>	6–17	–
	James' galleta	PLJA	<i>Pleuraphis jamesii</i>	0–17	–
3	<b>Subdominant Shrubs</b>			0–56	
	Bigelow sage	ARBI3	<i>Artemisia bigelovii</i>	0–17	0–1
	yellow rabbitbrush	CHVI8	<i>Chrysothamnus viscidiflorus</i>	0–17	0–1
	Stansbury cliffrose	PUST	<i>Purshia stansburiana</i>	0–6	0–1
	fourwing saltbush	ATCA2	<i>Atriplex canescens</i>	0–6	0–1
	crispleaf buckwheat	ERCO14	<i>Eriogonum corymbosum</i>	0–3	0–1
	broom snakeweed	GUSA2	<i>Gutierrezia sarothrae</i>	0–3	0–1
	plains pricklypear	OPPO	<i>Opuntia polyacantha</i>	0–3	0–1
	narrowleaf yucca	YUAN2	<i>Yucca angustissima</i>	0–2	0–1
<b>Grass/Grasslike</b>					
1	<b>Grasses</b>			0–28	
	James' galleta	PLJA	<i>Pleuraphis jamesii</i>	6–17	0–2
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	0–11	0–1
	needle and thread	HECO26	<i>Hesperostipa comata</i>	0–6	0–1
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	0–6	0–1
	Grass, annual	2GA	<i>Grass, annual</i>	0–6	0–1
	Grass, perennial	2GP	<i>Grass, perennial</i>	0–6	0–1
	sandhill muhly	MUPU2	<i>Muhlenbergia pungens</i>	0–3	0–1
	alkali sacaton	SPAI	<i>Sporobolus airoides</i>	0–3	0–1
<b>Forb</b>					
2	<b>Forbs</b>			0–28	
	prickly Russian thistle	SATR12	<i>Salsola tragus</i>	6–17	1–2
	scarlet globemallow	SPCO	<i>Sphaeralcea coccinea</i>	0–6	0–1
	desert princesplume	STPI	<i>Stanleya pinnata</i>	0–6	0–1
	Forb, annual	2FA	<i>Forb, annual</i>	0–6	0–1
	Forb, perennial	2FP	<i>Forb, perennial</i>	0–6	0–1
	rayless tansyaster	MAGR2	<i>Machaeranthera grindelioides</i>	0–6	0–1
	cleftleaf wildheliotrope	PHCR	<i>Phacelia crenulata</i>	0–6	0–1
	Wright's bird's beak	COWR2	<i>Cordylanthus wrightii</i>	0–3	0–1
	roughseed cryptantha	CRFL6	<i>Cryptantha flavoculata</i>	0–3	0–1
	Jones' pepperweed	LEMOJ	<i>Lepidium montanum</i> var. <i>jonesii</i>	0–3	0–1
	stemless four-nerve daisy	TEAC	<i>Tetraneuris acaulis</i>	0–3	0–1
	tufted evening primrose	OECA10	<i>Oenothera caespitosa</i>	0–1	0–1



## **Animal community**

### **--Wildlife Interpretation--**

Small herds of mule deer and pronghorn antelope can be seen grazing/browsing on these sites especially when near water sources and in the winter. Desert bighorn sheep may utilize this site when occurring on steeper slopes. The hot climate and lack of water favors small mammals, which have an easier time finding shelter, food, and water to live. Many species of rats, mice, squirrels, bats, and chipmunks can be observed, along with coyotes and foxes. Lizards are the most visible and can be observed during the day. Species may include the northern whiptail, desert spiny, and the colorful western collard lizard. (NPS.gov, 2008)

### **--Grazing Interpretations--**

This site provides very limited grazing for livestock. Torrey's jointfir provides fair forage for cattle and sheep on winter range. When present, grasses, primarily Indian ricegrass and James galleta, provide good forage for livestock, however, these species are often not abundant enough to support many livestock. Forage composition and annual production depend largely on yearly precipitation amounts and thus provide challenges for those making livestock grazing management decisions. Regardless of class of livestock, this site's carrying capacity is always low. A lack of available drinking water, can also influence its suitability for livestock grazing. Care should be taken to maintain the native perennial grasses and shrubs present on this site because they are hard to restore once gone.

This site may serve a dual purpose by also being important habitat for wildlife species such as pronghorn antelope, mule deer, and desert bighorn sheep, and can be important for wintering areas for bighorn sheep. In many places, however, wildlife populations are small and thus have little grazing/browsing impact on the site.

## **Recreational uses**

Resources that have special aesthetic and landscape values are wildflowers. Some recreation uses of this site include hiking, hunting, and horseback riding.

## **Wood products**

None

## **Other products**

None

## **Other information**

### **--Poisonous/Toxic Plant Communities--**

Toxic plants associated with this site include broom snakeweed. Broom snakeweed contains steroids, terpenoids, saponins, and flavones that can cause abortions or reproductive failure in sheep and cattle, however cattle are most susceptible. These toxins are most abundant during active growth and leafing stage. Cattle and sheep will generally only graze broom snakeweed when other forage is unavailable, typically in winter when toxicity levels are at their lowest.

Other potentially toxic plants associated with this site can include some buckwheat species, which may accumulate selenium, but only when growing on selenium enriched soils. These plants, when consumed will cause alkali disease or chronic selenosis, which affects all classes of livestock (excluding goats). Typically, animals consuming 5-50 ppm selenium will develop chronic selenosis and animals consuming greater than 50 ppm selenium will develop acute selenosis. Clinical signs include lameness, souging of the hoof, hair loss, blindness, and aimless wondering. Horses tend to develop what is called a "bob" tail or "roached" mane due to breakage of the long hairs.

Although not found on this site yet, Russian thistle is a toxic plant of concern in arid environments that could become established on this site. It can cause nitrate and to a lesser extent oxalate poisoning which affects all classes of livestock. The buildup of nitrates in these plants is highly dependent upon environmental factors, such as after a rain storm during a drought, during a period with cool/cloudy days, and on soils high in nitrogen and low in

sulfur and phosphorus. Nitrate collects in the stems and can persist throughout the growing season. Clinical signs of nitrate poisoning include drowsiness, weakness, muscular tremors, increased heart and respiratory rates, staggering gait, and death. Conversely, oxalate poisoning causes kidney failure; clinical signs include muscle tremors, tetany, weakness, and depression. Poisoning generally occurs when livestock consume and are not accustomed to grazing oxalate-containing plants. Animals with prior exposure to oxalates have increased numbers of oxalate-degrading rumen microflora and thus are able to degrade the toxin before clinical poisoning can occur (Knight and Walter, 2001).

#### --Invasive Plant Communities--

When ecological conditions deteriorate and perennial vegetation decreases due to disturbance (i.e., fire, over grazing, drought, off road vehicle overuse, erosion, etc.) annual forbs and grasses may invade this site. Of particular concern in semi-arid environments are non-native annual invaders including cheatgrass, Russian thistle, kochia, halogeton, and annual mustards. The presence of these species will depend on soil properties and moisture availability; however, these invaders are highly adaptive and can flourish in many locations. Once established, complete removal is difficult, but suppression may be possible. Due to Torrey jointfir's slow growth rate it does not compete well with invading plants after a disturbance and thus restoration efforts could be hindered.

#### --Fire Ecology--

The ability for an ecological site to carry fire depends primarily on its present fuel load and plant moisture content. Sites with small fuel loads will burn more slowly and less intensely than sites with large fuel loads. Many semi-desert plant communities in the Colorado Plateau may have evolved without a significant influence of fire. However, a year with exceptionally heavy winter rains can generate sufficient fuels for fire by producing heavy stands of annual forbs and grasses. When fires do occur, the effect on plant communities may be extreme due to the sites harsh environment and slow rate of recovery.

Fires on Torrey jointfir ecological sites are relatively uncommon due to sparse vegetation and insufficient fuels. Its fire regime depends on the adjacent plant communities and has a wide range of return intervals. This plant generally sprouts from the roots or woody crown after fire, but it also has the capability of reestablishing through seed. While it establishes quickly after fire, its slow growth rate inhibits vigorous competition with invading annuals which could change the fire regime due to an increase in fine fuels.

### **Inventory data references**

The data collected in 2010-2012 were in conjunction with the soil survey update for Capitol Reef National Park. The vegetation data was collected in association with a soil pit and geo-referenced. All the data is stored as hard copy files and in electronic format in the NRCS Utah State Office.

### **Other references**

Knight, A.P. and R.G. Walter. 2001. A guide to plant poisoning of animals in North America. Teton NewMedia. Jackson, WY.

NRCS Grazing Lands Technology Institute. 2003. National Range and Pasture Handbook. Fort Worth, TX, USA: US Department of Agriculture, Natural Resources Conservation Service, 190-VI-NRPH.

### **Contributors**

George Cook  
Jamin Johanson

### **Rangeland health reference sheet**

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community

cannot be used to identify the ecological site.

Author(s)/participant(s)	Shane A. Green (NRCS), Robert D. Stager (BLM), Dana Truman (NRCS), Paul Curtis (BLM) and Randy Beckstrand (BLM).
Contact for lead author	shane.green@ut.usda.gov
Date	03/21/2007
Approved by	Shane A. Green
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

- 1. Number and extent of rills:** A. On more gentle slopes (< 10 %): Few, but may occur on the site. Rills may be 6 to 10 feet in length. Sides of rills may be up to 2 inches high. Rills are most likely to form below adjacent exposed bedrock or water flow patterns where sufficient water accumulates to cause erosion. B. On steeper slopes (> 10 %): Common. Occur throughout the site. Rills may extend down entire slope.

---
- 2. Presence of water flow patterns:** Frequent and occur throughout area. They are expected to be long and connected into drainage networks. Evidence of flow will increase somewhat with slope.

---
- 3. Number and height of erosional pedestals or terracettes:** Pedestals form at the base of plants. On steeper slopes (>20%), gullies may remove soil from the base of shrubs and/or trees exposing roots that resemble pedestals. Terracettes may be present. Some debris dams of small to medium sized litter (up to 1 inch in diameter) may form in water flow patterns, rills, and gullies. These debris dams may accumulate smaller litter (leaves, grass and forb stems).

---
- 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 30 – 60%. Soil surface may have 0 to 5 percent rock fragments. Ground cover is based on first raindrop impact, and bare ground is the opposite of ground cover. Ground cover + bare ground = 100%. Poorly developed biological soil crusts that are interpreted as functioning as bare ground (therefore they would be susceptible to raindrop splash erosion) should be recorded as bare ground.

---
- 5. Number of gullies and erosion associated with gullies:** Very few to few on gentle slopes (< 10 %). On steeper slopes and areas below adjacent exposed bedrock, gullies may be more numerous. Length often extends the entire slope until it reaches an area where water and sediment accumulate. Gullies may show slightly more indication of erosion as slope steepens, or as the site occurs adjacent to steep areas where runoff accumulation occurs.

---
- 6. Extent of wind scoured, blowouts and/or depositional areas:** Very few. While there may be some evidence of wind generated soil movement, wind caused blowouts would not be expected. Some depositional areas may exist.

---
- 7. Amount of litter movement (describe size and distance expected to travel):** On gentle slopes (< 10 %) most litter accumulates at base of plants. Some down slope redistribution caused by water. Some litter removal may occur in flow

patterns and rills with deposition occurring at points of obstruction, especially following major storm events. Litter movement will increase with slopes > 10%.

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** This site should have a soil stability rating of 3 or 4 under the plant canopies and a rating of 2 to 3 in the interspaces using the soil stability kit test. The average should be a 3. Surface texture is loam to very fine sandy loam. Vegetation cover, litter, biological soil crusts and surface rock reduce erosion.
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** 0 to 0.5 inch; brown (7.5YR 5/4) loam, brown (7.5YR 4/4) moist; strong thin platy structure; slightly hard, friable, nonsticky and nonplastic; few very fine and fine roots; strongly effervescent, moderately alkaline (pH 8.4); abrupt smooth boundary. (0 to 2 inches thick)  
The A horizon would be expected to be more strongly developed under plant canopies. It is important if you are sampling to observe the A horizon under plant canopies as well as the interspaces. Use the specific information for the soil you are assessing found in the published soil survey to supplement this description.
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Distribution of vascular plants are expected to intercept raindrops reducing splash erosion but not eliminating it. Plants are usually distributed to slow runoff a little to allow time for some infiltration. With the physiographic location of the site being on remnant hillsides, rolling hills, pediment surfaces, alluvial fans, dissected benches and upland valley plains infiltration is somewhat reduced by slope and less plant cover. Natural erosion would be expected and especially in severe thunder storms or heavy spring runoff. When perennial grasses and shrubs decrease, reducing ground cover and increasing bare ground, runoff is expected to increase and any associated infiltration reduced.
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None. The associated structure is strong to moderate thin platy. This should not be considered to be compaction layers.
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Sprouting and nonsprouting shrubs = perennial grass > perennial and native annual forbs
- Sub-dominant:
- Other: Functional/structural groups may appropriately contain non-native species if their ecological function is the same as the native species in the reference state (e.g. Siberian Wheatgrass, Forage kochia etc.)  
Biological soil crust is variable in it's expression where present on this site and is measured as a component of ground cover.
- Additional: Disturbance regime includes erosion events, drought and insects. Following a recent disturbance such as drought or insects that removes the woody vegetation, forbs and perennial grasses (herbaceous species) may dominate the community. These conditions would reflect a functional community phase within the reference state.
-

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** In general, a mix of age classes for shrubs and grasses may occur with some dead and decadent plants present but the absence some age classes should not be a concern.
- 

14. **Average percent litter cover (%) and depth ( in):**
- 

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 135-390 Lbs/ac
- 

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** The only species expected to invade this site are those that can tolerate high gypsum and arid conditions. Cheatgrass and Russian thistle are noted invaders.
- 

17. **Perennial plant reproductive capability:** All perennial plants should have the ability to reproduce sexually or asexually in most years, except in drought years. The gypsic and arid nature of this site would be expected to inhibit some reproduction on site.
-