

Ecological site R035XY328UT Upland Very Steep Stony Loam (Pinyon-Utah Juniper)

Accessed: 04/23/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Robert Stager (BLM), Dana Truman (NRCS), Paul Curtis (BLM), Shane A. Green (NRCS), Randy Beckstrand (BLM)
Contact for lead author	
Date	01/30/2007
Approved by	Shane A. Green
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:** Common. Occur throughout the site. Rills may extend down entire slope or to obstructions such as large boulders.

2. **Presence of water flow patterns:** Frequent and occur throughout area. Flow patterns wind around the surface rocks and perennial plant bases.

3. **Number and height of erosional pedestals or terracettes:** Common-Pedestals form at the base of plants that occur on the edge of water flow patterns, rills and gullies. Gullies may remove soil from the base of trees exposing roots that resemble pedestals. Interspaces between well developed biological soil crusts resemble pedestals and may be up to 2 inches high. Terracettes are common. Debris dams of small to medium sized litter (up to 2 inches in diameter) may form in water flow patterns, rills, and gullies. These debris dams may accumulate smaller litter (leaves, grass and forb stems).

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 20 – 30 %. Most bare ground is associated with water flow patterns, rills, and gullies. Soil is covered by 25-50% rock fragments. Areas with well developed biological soil crusts should not be counted as bare ground. Poorly developed biological soil crusts that are interpreted as functioning as bare ground (therefore they would be susceptible to raindrop splash erosion) should be recorded as bare ground. Ground cover is based on first raindrop impact, and bare ground is the opposite of ground cover.

-
5. **Number of gullies and erosion associated with gullies:** Gullies are few to common. Length often extends from exposed bedrock until gully reaches a stream or an area where water and sediment accumulate, but they may be wide and shallow and armored with very large rocks. Gullies may remove soil from base of trees exposing roots.
-
6. **Extent of wind scoured, blowouts and/or depositional areas:** None to very few. Trees break the wind and reduce the potential for wind erosion.
-
7. **Amount of litter movement (describe size and distance expected to travel):** Fine litter is moved with even moderate precipitation events and spring runoff, accumulating down slope behind plants and rock features in the site. Woody stems may be washed from site. Gullies may remove accumulated litter from under trees.
-
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** This site should have a soil stability rating of 4 or 5 under the plant canopies using the soil stability kit test, and a rating of 3 to 4 in the interspaces. The average should be a 4. Surface texture is stony fine sandy loam. Vegetation cover, litter, biological soil crusts and surface rock reduce erosion.
-
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Soil surface is 1 inch deep. Structure is weak thin platy. Color is yellowish brown (10YR5/4). There is little if any difference under canopy or in interspaces and a recognizable A horizon is expected to be present throughout. Use the specific information for the soil you are assessing found in the published soil survey to supplement this description.
-
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Spatial distribution of plants and/or well developed biological soil crusts (where present) intercept raindrops reducing splash erosion and provide areas of surface detention to store water allowing additional time for infiltration.
-
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**): None. There may be layers of calcium carbonate or other naturally occurring hard layers found in the soil subsurface. These should not be considered to be compaction layers.
-
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant: Perennial bunchgrasses > Shrubs > Trees (Pinion > Juniper)
- Sub-dominant: forbs
- Other: Functional/structural groups may appropriately contain non-native species if their ecological function is the same as the native species in the reference state (e.g. Crested wheatgrass, Intermediate wheatgrass, etc.)
- Biological soil crust is variable in its expression where present on this site and is measured as a component of ground cover.

Additional: Disturbance regime includes parasites, insects, and drought. Dominants— Salina Wildrye, Utah Juniper, Pinyon Pine, Utah serviceberry. Sub Dominants— Indian ricegrass, forbs. Perennial and annual forbs can be expected to vary widely in their expression in the plant community based upon departures from average growing conditions.

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Several standing dead trees may be present on the site and approximately 20 % of the trees can show evidence of decadence. In drought tree mortality may increase with the first sign being a yellowish to reddish leaf color.
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 340-575lbs/acre annually
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** Cheatgrass is most likely to invade this site.
-

17. **Perennial plant reproductive capability:** All perennial plants should have the ability to reproduce sexually or asexually in most years, except in drought years.
-