

Ecological site R038XA114AZ Schist Hills 12-16" p.z.

Accessed: 02/17/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.




Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 038X–Mogollon Transition South

AZ 38.1 – Lower Mogollon Transition

Elevations range from 3000 to 4500 feet and precipitation averages 12 to 16 inches per year. Vegetation includes canotia, one-seed juniper, mesquite, catclaw acacia, jojoba, turbinella oak, ratany, shrubby buckwheat, algerita, skunkbush, tobosa, vine mesquite, bottlebrush squirreltail, grama species, curly mesquite, desert needlegrass and New Mexico feathergrass. The soil temperature regime is thermic and the soil moisture regime is ustic aridic. This unit occurs within the Transition Zone Physiographic Province and is characterized by canyons and structural troughs or valleys. Igneous, metamorphic and sedimentary rock classes occur on rough mountainous terrain in association with less extensive sediment filled valleys exhibiting little integrated drainage.

Classification relationships

This site is similar to TE Sites #250 and #275 found on the Prescott National Forest.

Associated sites

R038XA104AZ	Granitic Hills 12-16" p.z.
-------------	----------------------------

R038XA105AZ	Limestone Hills 12-16" p.z.
R038XA133AZ	Volcanic/Metamorphic Hills 12-16" p.z.
R038XA135AZ	Diabase Hills 12-16" p.z.

Similar sites

R040XA119AZ	Schist Hills 10"-13" p.z.
R038XB212AZ	Schist Hills 16-20" p.z.

Table 1. Dominant plant species

Tree	(1) <i>parkinsonia</i>
Shrub	(1) <i>Eriogonum fasciculatum</i> (2) <i>Simmondsia chinensis</i>
Herbaceous	(1) <i>aristida</i> (2) <i>selaginella</i>

Physiographic features

This site occurs at the lowest elevations of the interior chaparral zone in the Mogollon Transition area. This site occurs in an upland position. It occurs on hill-slopes, ridge-tops and mountains.

Table 2. Representative physiographic features

Landforms	(1) Hill (2) Mountain slope (3) Ridge
Flooding frequency	None
Elevation	3,100–4,600 ft
Slope	15–65%
Aspect	N, E, S

Climatic features

Precipitation in this common resource area averages 12 to 16 inches annually. The winter-summer rainfall ratio ranges from about 60/40% in the northwest part of the area to 50/50% in the southeast part. Summer rains fall July through September; are from high-intensity, convective thunderstorms. This moisture originates primarily from the Gulf of Mexico, but can come from the remnants of Pacific hurricanes in September. Winter moisture is frontal, originates in the north Pacific, and falls as rain or snow in widespread storms of low intensity and long duration. Snowfall ranges from a trace to 10 inches per year and can occur from November through March. Snow seldom persists for more than a day except on north aspects. May and June are the driest months of the year. Humidity is generally low all year. Average annual air temperatures range from 59 to 70 degrees F (thermic temperature regime). Daytime temperatures in the summer are commonly in the high 90's. Freezing temperatures are common from October through April, usually during the night or early morning hours. The actual precipitation, available moisture and temperature vary, depending on, region, elevation, rain shadow effect and aspect.

Table 3. Representative climatic features

Frost-free period (average)	230 days
Freeze-free period (average)	285 days
Precipitation total (average)	16 in

Influencing water features

There are no water features associated with this site.

Soil features

These soils are shallow (10 to 20 inches) and dark colored. They are loamy textured, non-calcareous and well drained. These soils range from lacking any development to having weak argillic horizons. They have formed in residuum and colluvium from schist and related metamorphic parent materials. Soil surfaces are well covered by brownish to silvery gravels, channers and stones. The effective rooting depth is limited by slightly weathered bedrock at 10 to 20 inches. Runoff is moderate to high on moist soils. The erosion hazard is slight due to gravel, cobble and rock covers. Rock outcrop and can be as high as 10%.

Typical taxonomic units where this site is mapped include: SSA-697 Mohave County Central part MU Birdsbeak-12.

Table 4. Representative soil features


Parent material	(1) Residuum–schist
Surface texture	(1) Very gravelly loam (2) Very gravelly sandy loam (3) Very channery sandy loam
Family particle size	(1) Loamy
Drainage class	Well drained
Permeability class	Moderately rapid to moderate
Soil depth	10–20 in
Surface fragment cover ≤3"	35–50%
Surface fragment cover >3"	1–10%
Available water capacity (0-40in)	0.6–2.2 in
Calcium carbonate equivalent (0-40in)	0–5%
Electrical conductivity (0-40in)	0–2 mmhos/cm
Sodium adsorption ratio (0-40in)	0–2
Soil reaction (1:1 water) (0-40in)	6.8–7.6
Subsurface fragment volume ≤3" (Depth not specified)	10–55%
Subsurface fragment volume >3" (Depth not specified)	0–10%

Ecological dynamics

The historic native plant community is a diverse mixture of desert trees, shrubs, succulents, forbs and grasses. This includes a diverse flora of native annual grasses and forbs of both the winter and summer seasons. Periodic wildfires occurred at moderate intervals (15 to 30 years) and helped maintain a balance between herbs and shrubs. In the absence of fire for longer periods, shrubby species and cacti can become dominant. The interactions of drought, fire and continuous livestock grazing can, over time, result in the loss of palatable grasses, half shrubs and suffrutescent forbs. In some situations non-native annuals can dominate the site. These species can, over time, diminish the soil seed-bank of native annual species. Non-native annuals can act to increase the fire frequency of areas of the site near roads and urban areas, where the incidence of man-made fires is high.

State and transition model

MLRA 38-1 (12-16''), Schist Hills



- 1a. Lack of fire for long periods, CHG and drought that reduced fuel loads.
- 1b. Unknown, possible herbicide followed by prescribed fire as maintenance.
- 2a. Introduction of seed source of exotic annuals like red brome, wild oats plus increased fire frequency (every 5-10 years)
- 2b. Unknown
- 3. Introduction of seed source of exotic annuals, El Nino type event, catastrophic fire.
- 4, 5. Accelerated soil erosion may occur where vegetation is absent. Repeated fires may remove most perennial vegetation. Slopes are trailed, soils are compacted and fill erosion occurs.

* Annual grasses include natives and non-natives

Figure 4. State and Transition, Schist Hills 12-16" p.z.

State 1 Mixed Shrub/Herbaceous State

Community 1.1 Historic Native Plant Community

The historic native plant community is a diverse mixture of perennial grasses, suffrutescent forbs, shrubs, succulents and desert trees. A rich flora of native annual forbs and grasses, of both the winter and summer seasons, exist in the plant community. Periodic, naturally occurring, wildfires were important in maintaining the potential plant community. North slopes have a chaparral of evergreen shrubs like jojoba, turbinella oak and flattop buckwheat. Southern exposures will have a higher percentage of desert shrubs, trees and succulents in the plant community. More xeric grasses will dominate southern exposures (aristida, tanglehead). Grasses on cooler aspects include desert stipa and sideoats grama. Club moss (selaginella) is especially common on very shallow soil areas.

Table 5. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Shrub/Vine	270	400	650
Grass/Grasslike	75	200	520
Forb	25	75	250
Tree	5	30	100
Total	375	705	1520

Table 6. Soil surface cover

Tree basal cover	0-1%
Shrub/vine/liana basal cover	1-2%
Grass/grasslike basal cover	1-2%
Forb basal cover	1-2%
Non-vascular plants	0%
Biological crusts	0-1%
Litter	15-50%
Surface fragments >0.25" and <=3"	35-60%
Surface fragments >3"	1-10%
Bedrock	0-10%
Water	0%
Bare ground	10-50%

Table 7. Canopy structure (% cover)

Height Above Ground (Ft)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.5	–	0-5%	0-10%	1-10%
>0.5 <= 1	–	1-5%	1-5%	1-5%
>1 <= 2	–	5-10%	1-5%	1-5%
>2 <= 4.5	–	5-15%	0-2%	0-1%
>4.5 <= 13	1-10%	10-15%	–	–
>13 <= 40	0-5%	–	–	–
>40 <= 80	–	–	–	–
>80 <= 120	–	–	–	–
>120	–	–	–	–

Figure 6. Plant community growth curve (percent production by month). AZ3811, 38.1 12-16" p.z. all sites. Growth begins in the spring, most growth occurs in the summer..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	1	7	15	20	22	20	10	5	0	0

State 2 Shrub Dominated State

Community 2.1 Shrub Dominated Plant Community

Perennial grass canopy cover is reduced due to the interactions of drought, grazing and fire. Desert shrubs and cacti dominate the plant community. Shrub cover exceeds 30%. Major species include jojoba, flattop buckwheat and paloverde. Annuals, both native and non-native, dominate the under-story. Fire frequency is reduced but the site can still burn, especially after "El Nino" years produce heavy fuel loads of annual grasses and forbs.

State 3 Exotic Invaded State

Community 3.1 Exotic Invaded Plant Community

Non-native annual grasses and forbs like; red brome, cheatgrass, and wild oats, can invade and dominate areas of the site. These species can, over time, reduce the seed-bank of native annual grasses and forbs. Their presence can increase the fire frequency (of man made fires) especially where roads and urban areas are adjacent to areas of the site. Repeated fires tend to remove fire sensitive species like paloverde, cacti and buckwheat, and leave fire tolerant species like turbinella oak, mesquite, whitethorn and jojoba.

State 4 Shrub Dominated and Eroded State

Community 4.1 Shrub Dominated, Eroded Plant Community

Shrubs like jojoba, mesquite, turbinella oak and flattop buckwheat, and succulents like prickly pear and cholla can increase to dominate the site in the absence of fire for very long periods of time. Native and non-native annual forbs and grasses dominate the under-story. In "El Nino" years, herbaceous fuels can be sufficient to carry fire through the heavy canopy of shrubs. The major woody shrubs are, however, fire resistant once established. Such fires would remove less tolerant species like cacti, buckwheat and paloverde and leave intact the sprouting woody plants to become more and more dominant. Extreme rainfall events coupled with; the fire, drought and grazing interaction,

can lead to rilling of steep slopes. Compaction of soils can occur with heavy trailing from continuous livestock use. Loss of plant cover after repeated fire can lead to accelerated rill erosion under these circumstances.

Additional community tables

Table 8. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
Grass/Grasslike					
1	Dominant perennial grasses			50–200	
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	10–100	–
	slender grama	BORE2	<i>Bouteloua repens</i>	5–50	–
	purple threeawn	ARPU9	<i>Aristida purpurea</i>	5–50	–
	slim tridens	TRMU	<i>Tridens muticus</i>	1–50	–
	tanglehead	HECO10	<i>Heteropogon contortus</i>	1–50	–
	Parish's threeawn	ARPUP5	<i>Aristida purpurea var. parishii</i>	0–30	–
	spidergrass	ARTE3	<i>Aristida ternipes</i>	1–30	–
	cane bluestem	BOBA3	<i>Bothriochloa barbinodis</i>	1–30	–
	black grama	BOER4	<i>Bouteloua eriopoda</i>	0–30	–
	Arizona cottontop	DICA8	<i>Digitaria californica</i>	0–20	–
	bush muhly	MUPO2	<i>Muhlenbergia porteri</i>	0–20	–
2	Cool season grasses			10–70	
	desert needlegrass	ACSP12	<i>Achnatherum speciosum</i>	10–50	–
	squirreltail	ELEL5	<i>Elymus elymoides</i>	0–10	–
	needle and thread	HECO26	<i>Hesperostipa comata</i>	0–10	–
	New Mexico feathergrass	HENE5	<i>Hesperostipa neomexicana</i>	0–10	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	0–10	–
3	Misc. perennial grasses			10–100	
	big galleta	PLRI3	<i>Pleuraphis rigida</i>	0–25	–
	hairy grama	BOHI2	<i>Bouteloua hirsuta</i>	0–20	–
	curly-mesquite	HIBE	<i>Hilaria belangeri</i>	0–10	–
	tobosagrass	PLMU3	<i>Pleuraphis mutica</i>	0–10	–
	southwestern bristlegrass	SESC2	<i>Setaria scheelei</i>	0–5	–
	plains bristlegrass	SEVU2	<i>Setaria vulpiseta</i>	0–5	–
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	0–5	–
	slim tridens	TRMUE	<i>Tridens muticus var. elongatus</i>	0–5	–
	fall witchgrass	DICO6	<i>Digitaria cognata</i>	0–5	–
	red grama	BOTR2	<i>Bouteloua trifida</i>	0–5	–
	Hall's panicgrass	PAHA	<i>Panicum hallii</i>	0–5	–
	Fendler threeawn	ARPUL	<i>Aristida purpurea var. longiseta</i>	0–5	–
	blue threeawn	ARPUN	<i>Aristida purpurea var. nealleyi</i>	0–5	–
	spidergrass	ARTEG	<i>Aristida ternipes var. gentilis</i>	0–5	–
	low woollygrass	DAPU7	<i>Dasyochloa pulchella</i>	0–1	–
	nineawn pappusgrass	ENDE	<i>Enneapogon desvauxii</i>	0–1	–
	plains lovegrass	ERIN	<i>Eragrostis intermedia</i>	0–1	–
	green sprangletop	LEDI1	<i>Lentochloa dubia</i>	0–1	–

	green sprangletop	LEBO	<i>Leptochloa dubia</i>	0-1	-
	vine mesquite	PAOB	<i>Panicum obtusum</i>	0-1	-
4	Annual grasses			5-150	
	sixweeks threeawn	ARAD	<i>Aristida adscensionis</i>	1-50	-
	mucronate sprangletop	LEPAB	<i>Leptochloa panicea ssp. brachiata</i>	1-25	-
	small fescue	VUMI	<i>Vulpia microstachys</i>	1-20	-
	Eastwood fescue	VUMIC	<i>Vulpia microstachys var. ciliata</i>	1-20	-
	sixweeks fescue	VUOC	<i>Vulpia octoflora</i>	1-20	-
	Mexican panicgrass	PAH15	<i>Panicum hirticaule</i>	0-10	-
	Arizona signalgrass	URAR	<i>Urochloa arizonica</i>	0-10	-
	prairie threeawn	AROL	<i>Aristida oligantha</i>	0-10	-
	needle grama	BOAR	<i>Bouteloua aristidoides</i>	0-5	-
	witchgrass	PACA6	<i>Panicum capillare</i>	0-5	-
	delicate muhly	MUFR	<i>Muhlenbergia fragilis</i>	0-2	-
	littleseed muhly	MUMI	<i>Muhlenbergia microsperma</i>	0-2	-
	Rothrock's grama	BORO2	<i>Bouteloua rothrockii</i>	0-2	-
	Arizona brome	BRAR4	<i>Bromus arizonicus</i>	0-2	-
	feather fingergrass	CHV14	<i>Chloris virgata</i>	0-1	-
	canyon cupgrass	ERLE7	<i>Eriochloa lemmonii</i>	0-1	-
	tufted lovegrass	ERPE	<i>Eragrostis pectinacea</i>	0-1	-
	desert lovegrass	ERPEM	<i>Eragrostis pectinacea var. miserrima</i>	0-1	-
	little barley	HOPU	<i>Hordeum pusillum</i>	0-1	-
	Mexican sprangletop	LEFUU	<i>Leptochloa fusca ssp. uninervia</i>	0-1	-
	sixweeks grama	BOBA2	<i>Bouteloua barbata</i>	0-1	-
	Bigelow's bluegrass	POBI	<i>Poa bigelovii</i>	0-1	-
Forb					
5	Perennial forbs			20-100	
	spikemoss	SELAG	<i>Selaginella</i>	1-40	-
	desert globemallow	SPAM2	<i>Sphaeralcea ambigua</i>	1-15	-
	brownplume wirelettuce	STPA4	<i>Stephanomeria pauciflora</i>	1-10	-
	slender janusia	JAGR	<i>Janusia gracilis</i>	0-10	-
	lacy tansyaster	MAPI	<i>Machaeranthera pinnatifida</i>	1-10	-
	plains blackfoot	MELE2	<i>Melampodium leucanthum</i>	0-5	-
	wishbone-bush	MILAV	<i>Mirabilis laevis var. villosa</i>	0-5	-
	Arizona wrightwort	CAAR7	<i>Carlwrightia arizonica</i>	0-5	-
	bluedicks	DICA14	<i>Dichelostemma capitatum</i>	1-5	-
	weakleaf bur ragweed	AMCO3	<i>Ambrosia confertiflora</i>	1-5	-
	perennial rockcress	ARPE2	<i>Arabis perennans</i>	1-5	-
	white sagebrush	ARLUM2	<i>Artemisia ludoviciana ssp. mexicana</i>	1-5	-
	climbing wartclub	BOSC	<i>Boerhavia scandens</i>	0-5	-
	lipfern	CHEIL	<i>Cheilanthes</i>	0-5	-
	trailing windmills	ALIN	<i>Allionia incarnata</i>	0-5	-
	shrubby deervetch	LORI3	<i>Lotus rigidus</i>	0-5	-

cliffbrake	PELLA	<i>Pellaea</i>	0-5	-
Coues' cassia	SECO10	<i>Senna covesii</i>	1-5	-
slender poreleaf	POGR5	<i>Porophyllum gracile</i>	0-5	-
Parry's false prairie-clover	MAPA7	<i>Marina parryi</i>	0-2	-
Parry's beardtongue	PEPA24	<i>Penstemon parryi</i>	0-2	-
sego lily	CANU3	<i>Calochortus nuttallii</i>	0-2	-
desert trumpet	ERIN4	<i>Eriogonum inflatum</i>	0-2	-
Mojave spurge	EUSC6	<i>Euphorbia schizoloba</i>	0-2	-
southwestern mock vervain	GLGO	<i>Glandularia gooddingii</i>	0-2	-
desert rosemallow	HICO	<i>Hibiscus coulteri</i>	0-2	-
paleface	HIDE	<i>Hibiscus denudatus</i>	0-1	-
Indian rushpea	HOGL2	<i>Hoffmannseggia glauca</i>	0-1	-
fleabane	ERIGE2	<i>Erigeron</i>	0-1	-
desert tobacco	NIOB	<i>Nicotiana obtusifolia</i>	0-1	-
cloak fern	NOTHO	<i>Notholaena</i>	0-1	-
New Mexico groundsel	PANE7	<i>Packera neomexicana</i>	0-1	-
Oak Creek ragwort	PAQU8	<i>Packera quercetorum</i>	0-1	-
toadflax penstemon	PELI2	<i>Penstemon linarioides</i>	0-1	-
bigseed alfalfa dodder	CUIN	<i>Cuscuta indecora</i>	0-1	-
Cooley's bundleflower	DECO2	<i>Desmanthus cooleyi</i>	0-1	-
desert larkspur	DEPA	<i>Delphinium parishii</i>	0-1	-
tall mountain larkspur	DESC	<i>Delphinium scaposum</i>	0-1	-
wavyleaf Indian paintbrush	CAAPM	<i>Castilleja applegatei</i> ssp. <i>martinii</i>	0-1	-
New Mexico silverbush	ARNE2	<i>Argythamnia neomexicana</i>	0-1	-
dense ayenia	AYMI	<i>Ayenia microphylla</i>	0-1	-
desert marigold	BAMU	<i>Baileya multiradiata</i>	0-1	-
scarlet spiderling	BOCO	<i>Boerhavia coccinea</i>	0-1	-
tuber anemone	ANTU	<i>Anemone tuberosa</i>	0-1	-
narrowleaf silverbush	ARLA12	<i>Argythamnia lanceolata</i>	0-1	-
white sagebrush	ARLU	<i>Artemisia ludoviciana</i>	0-1	-
largeflower onion	ALMA4	<i>Allium macropetalum</i>	0-1	-
dwarf desertpeony	ACNA2	<i>Acourtia nana</i>	0-1	-
brownfoot	ACWR5	<i>Acourtia wrightii</i>	0-1	-
San Felipe dogweed	ADPO	<i>Adenophyllum porophylloides</i>	0-1	-
desert penstemon	PEPS	<i>Penstemon pseudospectabilis</i>	0-1	-
orange fameflower	PHAU13	<i>Phemeranthus aurantiacus</i>	0-1	-
Wright's deervetch	LOWR	<i>Lotus wrightii</i>	0-1	-
ragged nettlespurge	JAMA	<i>Jatropha macrorhiza</i>	0-1	-
longflower tube tongue	JULO3	<i>Justicia longii</i>	0-1	-
Lemmon's ragwort	SELE8	<i>Senecio lemmonii</i>	0-1	-
silverleaf nightshade	SOEL	<i>Solanum elaeagnifolium</i>	0-1	-
glandleaf milkwort	POMA7	<i>Polygala macradenia</i>	0-1	-
scurfpea	PSORA2	<i>Psoralidium</i>	0-1	-
canigre dock	RIIHY	<i>Rumex hymenocarpus</i>	0-1	-

	Common name	Code	Scientific name	Height	Notes
	twinleaf senna	SEBA3	<i>Senna bauhinioides</i>	0-1	-
	turpentinebroom	THMO	<i>Thamnosma montana</i>	0-1	-
	branched noseburn	TRRA5	<i>Tragia ramosa</i>	0-1	-
	Louisiana vetch	VILUL2	<i>Vicia ludoviciana ssp. ludoviciana</i>	0-1	-
6	Annual forbs			5-150	
	Coulter's lupine	LUSP2	<i>Lupinus sparsiflorus</i>	0-30	-
	California poppy	ESCAM	<i>Eschscholzia californica ssp. mexicana</i>	0-25	-
	bristly fiddleneck	AMTE3	<i>Amsinckia tessellata</i>	0-25	-
	phacelia	PHACE	<i>Phacelia</i>	0-25	-
	western tansymustard	DEPI	<i>Descurainia pinnata</i>	0-20	-
	Coulter's spiderling	BOCO2	<i>Boerhavia coulteri</i>	0-15	-
	exserted Indian paintbrush	CAEXE	<i>Castilleja exserta ssp. exserta</i>	0-15	-
	pitseed goosefoot	CHBE4	<i>Chenopodium berlandieri</i>	0-10	-
	slender goldenweed	MAGR10	<i>Machaeranthera gracilis</i>	0-10	-
	tanseyleaf tansyaster	MATA2	<i>Machaeranthera tanacetifolia</i>	0-10	-
	combseed	PECTO	<i>Pectocarya</i>	0-5	-
	shaggyfruit pepperweed	LELA	<i>Lepidium lasiocarpum</i>	0-5	-
	foothill deervetch	LOHU2	<i>Lotus humistratus</i>	0-5	-
	desertparsley	LOMAT	<i>Lomatium</i>	0-5	-
	coastal bird's-foot trefoil	LOSA	<i>Lotus salsuginosus</i>	0-5	-
	Arizona lupine	LUAR4	<i>Lupinus arizonicus</i>	0-5	-
	American wild carrot	DAPU3	<i>Daucus pusillus</i>	0-5	-
	flatcrown buckwheat	ERDE6	<i>Eriogonum deflexum</i>	0-5	-
	sorrel buckwheat	ERPO4	<i>Eriogonum polycladon</i>	0-5	-
	Arizona poppy	KAGR	<i>Kallstroemia grandiflora</i>	0-5	-
	cryptantha	CRYPT	<i>Cryptantha</i>	0-5	-
	fivewing spiderling	BOIN	<i>Boerhavia intermedia</i>	0-5	-
	milkvetch	ASTRA	<i>Astragalus</i>	0-5	-
	Arizona popcornflower	PLAR	<i>Plagiobothrys arizonicus</i>	0-5	-
	sleepy silene	SIAN2	<i>Silene antirrhina</i>	0-5	-
	lyreleaf jewelflower	STCA5	<i>Streptanthus carinatus</i>	0-5	-
	moth combseed	PESE	<i>Pectocarya setosa</i>	0-5	-
	thelypod	THELY	<i>Thelypodium</i>	0-5	-
	woolly tidestromia	TILA2	<i>Tidestromia lanuginosa</i>	0-5	-
	desert Indianwheat	PLOV	<i>Plantago ovata</i>	1-5	-
	woolly plantain	PLPA2	<i>Plantago patagonica</i>	0-5	-
	New Mexico plumeseed	RANE	<i>Rafinesquia neomexicana</i>	0-5	-
	purslane	PORTU	<i>Portulaca</i>	0-2	-
	sand fringe-pod	THCU	<i>Thysanocarpus curvipes</i>	0-2	-
	creamcups	PLCA5	<i>Platystemon californicus</i>	0-2	-
	annual agoseris	AGHE2	<i>Agoseris heterophylla</i>	0-2	-
	carelessweed	AMPA	<i>Amaranthus palmeri</i>	0-2	-
	New Mexico thistle	CINE	<i>Cirsium neomexicanum</i>	0-2	-

NEW MEXICO NAME	SYMBOL	SCIENTIFIC NAME	HEIGHT	FRUIT
miner's lettuce	CLPEP	<i>Claytonia perfoliata</i> ssp. <i>perfoliata</i>	0–2	–
crestrub morning-glory	IPCO2	<i>Ipomoea costellata</i>	0–2	–
spurge	EUPHO	<i>Euphorbia</i>	0–2	–
miniature woollystar	ERDI2	<i>Eriastrum diffusum</i>	0–2	–
spreading fleabane	ERDI4	<i>Erigeron divergens</i>	0–2	–
wedgeleaf draba	DRCU	<i>Draba cuneifolia</i>	0–2	–
miniature lupine	LUBI	<i>Lupinus bicolor</i>	0–2	–
Thurber's pepperweed	LETH2	<i>Lepidium thurberi</i>	0–2	–
hollowleaf annual lupine	LUSU3	<i>Lupinus succulentus</i>	0–2	–
Gordon's bladderpod	LEGO	<i>Lesquerella gordonii</i>	0–2	–
Florida pellitory	PAFL3	<i>Parietaria floridana</i>	0–2	–
manybristle chinchweed	PEPA2	<i>Pectis papposa</i>	0–1	–
whitestem blazingstar	MEAL6	<i>Mentzelia albicaulis</i>	0–1	–
green carpetweed	MOVE	<i>Mollugo verticillata</i>	0–1	–
desert evening primrose	OEPR	<i>Oenothera primiveris</i>	0–1	–
Fendler's desertydandelion	MAFE	<i>Malacothrix fendleri</i>	0–1	–
sacred thorn-apple	DAWR2	<i>Datura wrightii</i>	0–1	–
Texas stork's bill	ERTE13	<i>Erodium texanum</i>	0–1	–
star gilia	GIST	<i>Gilia stellata</i>	0–1	–
longleaf false goldeneye	HELOA2	<i>Heliomeris longifolia</i> var. <i>annua</i>	0–1	–
redstar	IPCO3	<i>Ipomoea coccinea</i>	0–1	–
ivyleaf morning-glory	IPHE	<i>Ipomoea hederacea</i>	0–1	–
California goldfields	LACA7	<i>Lasthenia californica</i>	0–1	–
scrambled eggs	COAU2	<i>Corydalis aurea</i>	0–1	–
hairy prairie clover	DAMO	<i>Dalea mollis</i>	0–1	–
brittle spineflower	CHBR	<i>Chorizanthe brevicornu</i>	0–1	–
Esteve's pincushion	CHST	<i>Chaenactis stevioides</i>	0–1	–
yellow tackstem	CAPA7	<i>Calycoseris parryi</i>	0–1	–
white tackstem	CAWR	<i>Calycoseris wrightii</i>	0–1	–
hoary bowlesia	BOIN3	<i>Bowlesia incana</i>	0–1	–
woollyhead neststraw	STMI2	<i>Stylocline micropoides</i>	0–1	–
desert unicorn-plant	PRAL4	<i>Proboscidea althaeifolia</i>	0–1	–
doubleclaw	PRPA2	<i>Proboscidea parviflora</i>	0–1	–
chia	SACO6	<i>Salvia columbariae</i>	0–1	–
sawtooth sage	SASU7	<i>Salvia subincisa</i>	0–1	–
ragwort	SENEC	<i>Senecio</i>	0–1	–
spreading fanpetals	SIAB	<i>Sida abutilifolia</i>	0–1	–

Shrub/Vine

7	Evergreen shrubs		150–300		
	jojoba	SICH	<i>Simmondsia chinensis</i>	100–250	–
	Sonoran scrub oak	QUTU2	<i>Quercus turbinella</i>	0–100	–
	desert ceanothus	CEGR	<i>Ceanothus greggii</i>	0–10	–
	snapdragon penstemon	KEANM	<i>Keckiella antirrhinoides</i> ssp. <i>microphylla</i>	0–10	–

			<i>microphylla</i>		
	algerita	MATR3	<i>Mahonia trifoliolata</i>	0–10	–
	red barberry	MAHA4	<i>Mahonia haematocarpa</i>	0–5	–
	redberry buckthorn	RHCR	<i>Rhamnus crocea</i>	0–5	–
	sugar sumac	RHOV	<i>Rhus ovata</i>	0–5	–
8	Large shrubs			15–60	
	ocotillo	FOSP2	<i>Fouquieria splendens</i>	1–20	–
	catclaw acacia	ACGR	<i>Acacia greggii</i>	1–10	–
	Wright's beebrush	ALWR	<i>Aloysia wrightii</i>	0–10	–
	Berlandier's wolfberry	LYBE	<i>Lycium berlandieri</i>	1–5	–
	Arizona desert-thorn	LYEX	<i>Lycium exsertum</i>	0–5	–
	whitethorn acacia	ACCO2	<i>Acacia constricta</i>	0–5	–
	spiny hackberry	CEEH	<i>Celtis ehrenbergiana</i>	0–2	–
	desert sweet	CHMI2	<i>Chamaebatiaria millefolium</i>	0–1	–
	Warnock's snakewood	COWA	<i>Condalia warnockii</i>	0–1	–
	winterfat	KRLA2	<i>Krascheninnikovia lanata</i>	0–1	–
	creosote bush	LATR2	<i>Larrea tridentata</i>	0–1	–
	ambrosia leaf bur ragweed	AMAM2	<i>Ambrosia ambrosioides</i>	0–1	–
	Thurber's desert honeysuckle	ANTH2	<i>Anisacanthus thurberi</i>	0–1	–
	fourwing saltbush	ATCA2	<i>Atriplex canescens</i>	0–1	–
	desertbroom	BASA2	<i>Baccharis sarothroides</i>	0–1	–
	pale desert-thorn	LYPA	<i>Lycium pallidum</i>	0–1	–
	catclaw mimosa	MIACB	<i>Mimosa aculeaticarpa</i> var. <i>biuncifera</i>	0–1	–
	mariola	PAIN2	<i>Parthenium incanum</i>	0–1	–
	littleleaf sumac	RHMI3	<i>Rhus microphylla</i>	0–1	–
	skunkbush sumac	RHTR	<i>Rhus trilobata</i>	0–1	–
	Mexican bladdersage	SAME	<i>Salazaria mexicana</i>	0–1	–
	lotebush	ZIOBC	<i>Ziziphus obtusifolia</i> var. <i>canescens</i>	0–1	–
9	Dominant half shrubs			100–250	
	Eastern Mojave buckwheat	ERFA2	<i>Eriogonum fasciculatum</i>	50–200	–
	bastardsage	ERWR	<i>Eriogonum wrightii</i>	5–50	–
	fairyduster	CAER	<i>Calliandra eriophylla</i>	5–25	–
	littleleaf ratany	KRER	<i>Krameria erecta</i>	1–15	–
	rough menodora	MESC	<i>Menodora scabra</i>	0–10	–
	Coulter's brickellbush	BRCO	<i>Brickellia coulteri</i>	0–5	–
	American threefold	TRCA8	<i>Trixis californica</i>	1–5	–
	Parish's goldeneye	VIPA14	<i>Viguiera parishii</i>	0–5	–
	longleaf phlox	PHLO2	<i>Phlox longifolia</i>	0–2	–
	starry bedstraw	GAST	<i>Galium stellatum</i>	0–1	–
	ragged rockflower	CRBI2	<i>Crossosoma bigelovii</i>	0–1	–
	desert zinnia	ZIAC	<i>Zinnia acerosa</i>	0–1	–
	shortleaf baccharis	BABR	<i>Baccharis brachyphylla</i>	0–1	–
	yerba de pasmo	BAPT	<i>Baccharis pteronioides</i>	0–1	–

	sweetbush	BEJU	<i>Bebbia juncea</i>	0–1	–
10	Succulents			5–70	
	buck-horn cholla	CYAC8	<i>Cylindropuntia acanthocarpa</i>	1–15	–
	cactus apple	OPEN3	<i>Opuntia engelmannii</i>	1–15	–
	tulip pricklypear	OPPH	<i>Opuntia phaeacantha</i>	0–10	–
	banana yucca	YUBA	<i>Yucca baccata</i>	0–10	–
	walkingstick cactus	CYSP8	<i>Cylindropuntia spinosior</i>	0–5	–
	saguaro	CAGI10	<i>Carnegiea gigantea</i>	0–5	–
	dollarjoint pricklypear	OPCH	<i>Opuntia chlorotica</i>	0–2	–
	candy barrelcactus	FEWI	<i>Ferocactus wislizeni</i>	0–2	–
	pinkflower hedgehog cactus	ECFA	<i>Echinocereus fasciculatus</i>	0–2	–
	common sotol	DAWH2	<i>Dasyliion wheeleri</i>	0–2	–
	teddybear cholla	CYBI9	<i>Cylindropuntia bigelovii</i>	0–2	–
	jumping cholla	CYFU10	<i>Cylindropuntia fulgida</i>	0–2	–
	goldenflower century plant	AGCH2	<i>Agave chrysantha</i>	0–2	–
	Palmer's century plant	AGPA3	<i>Agave palmeri</i>	0–1	–
	Schott's century plant	AGSC3	<i>Agave schottii</i>	0–1	–
	Christmas cactus	CYLE8	<i>Cylindropuntia leptocaulis</i>	0–1	–
	Arizona hedgehog cactus	ECCOA	<i>Echinocereus coccineus var. arizonicus</i>	0–1	–
	Engelmann's hedgehog cactus	ECEN	<i>Echinocereus engelmannii</i>	0–1	–
	redspine fishhook cactus	ECER2	<i>Echinomastus erectocentrus</i>	0–1	–
	spinystar	ESVI2	<i>Escobaria vivipara</i>	0–1	–
	Whipple cholla	CYWH	<i>Cylindropuntia whipplei</i>	0–1	–
	soaptree yucca	YUEL	<i>Yucca elata</i>	0–1	–
	Graham's nipple cactus	MAGR9	<i>Mammillaria grahamii</i>	0–1	–
	sacahuista	NOMI	<i>Nolina microcarpa</i>	0–1	–
11	Increaser half-shrubs			1–20	
	broom snakeweed	GUSA2	<i>Gutierrezia sarothrae</i>	1–10	–
	brittlebush	ENFA	<i>Encelia farinosa</i>	0–5	–
	button brittlebush	ENFR	<i>Encelia frutescens</i>	0–5	–
	turpentine bush	ERLA12	<i>Ericameria laricifolia</i>	0–1	–
	narrowleaf goldenbush	ERLI6	<i>Ericameria linearifolia</i>	0–1	–
	threadleaf snakeweed	GUMI	<i>Gutierrezia microcephala</i>	0–1	–
	burweed	ISTE2	<i>Isocoma tenuisecta</i>	0–1	–
Tree					
12	Trees			5–100	
	yellow paloverde	PAMI5	<i>Parkinsonia microphylla</i>	5–35	–
	crucifixion thorn	CAHO3	<i>Canotia holacantha</i>	0–20	–
	redberry juniper	JUCO11	<i>Juniperus coahuilensis</i>	0–10	–
	oneseed juniper	JUMO	<i>Juniperus monosperma</i>	0–10	–
	blue paloverde	PAFL6	<i>Parkinsonia florida</i>	0–5	–
	western honey mesquite	PRGLT	<i>Prosopis glandulosa var. torreyana</i>	0–5	–

Animal community

This site is suitable for grazing year round, but is not easily traversed by livestock. Livestock grazing use is concentrated on south slopes, canyon bottoms and ridge-tops. North slopes may be little used. Slopes greater than 50% and areas with very cobbly surfaces limit grazing use by cattle. Areas of rock outcrop can form barriers to livestock movement. The site is susceptible to erosion in overgrazed areas like bed-grounds, livestock trails and lower slopes adjacent to water.

The site has good habitat diversity for a great variety of desert wildlife species. Water developments are very important to both livestock and wildlife on this site.

Hydrological functions

This site has smooth surfaces with high covers of gravels and channers. When soils are dry, it produces little runoff. It produces significant runoff only when heavy rain falls on snow or moist soils.

Recreational uses

Hunting, camping, horseback riding, backpacking, rock hounding, photography.

Wood products

Limited harvest of fuel-wood, fence posts and stays from mesquite, juniper and saguaro.

Other products

There is some native harvest of food plants like grass nuts, thistle, prickly pear tunas and jjoba nuts.

Type locality

Location 1: Pinal County, AZ	
General legal description	Boyce Thompson Arboretum.
Location 2: Pinal County, AZ	
Township/Range/Section	T7S R17E S13
General legal description	Miller Dry Camp Ranch, below Table mountain

Contributors

Dan Robinett
Larry D. Ellicott

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	

Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

1. **Number and extent of rills:**

2. **Presence of water flow patterns:**

3. **Number and height of erosional pedestals or terracettes:**

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**

5. **Number of gullies and erosion associated with gullies:**

6. **Extent of wind scoured, blowouts and/or depositional areas:**

7. **Amount of litter movement (describe size and distance expected to travel):**

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**
-

17. **Perennial plant reproductive capability:**
-