


## **Ecological site R038XA135AZ Diabase Hills 12-16" p.z.**

Accessed: 05/04/2024

### General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



**Figure 1. Mapped extent**

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

### MLRA notes

Major Land Resource Area (MLRA): 038X–Mogollon Transition South

AZ 38.1 – Lower Mogollon Transition

Elevations range from 3000 to 4500 feet and precipitation averages 12 to 16 inches per year. Vegetation includes canotia, one-seed juniper, mesquite, catclaw acacia, jojoba, turbinella oak, ratany, shrubby buckwheat, algerita, skunkbush, tobosa, vine mesquite, bottlebrush squirreltail, grama species, curly mesquite, desert needlegrass and New Mexico feathergrass. The soil temperature regime is thermic and the soil moisture regime is ustic aridic. This unit occurs within the Transition Zone Physiographic Province and is characterized by canyons and structural troughs or valleys. Igneous, metamorphic and sedimentary rock classes occur on rough mountainous terrain in association with less extensive sediment filled valleys exhibiting little integrated drainage.

### Associated sites

R038XA104AZ	<b>Granitic Hills 12-16" p.z.</b>
R038XA105AZ	<b>Limestone Hills 12-16" p.z.</b>
R038XA117AZ	<b>Volcanic Hills 12-16" p.z. Clayey</b>

## Similar sites

R040XA105AZ	Shallow Hills 10"-13" p.z.
R040XA119AZ	Schist Hills 10"-13" p.z.

Table 1. Dominant plant species

Tree	(1) <i>Parkinsonia</i>
Shrub	(1) <i>acacia greggii</i> (2) <i>erigonum wrightii</i>
Herbaceous	(1) <i>bouteloua eriopoda</i> (2) <i>muhlenbergia porteri</i>

## Physiographic features

This site occurs at the lowest elevations of the interior chaparral zone in the Mogollon Transition area. This site occurs in an upland position. It occurs on hill-slopes, ridge-tops and mountains.

Table 2. Representative physiographic features

Landforms	(1) Hill (2) Mountain slope (3) Ridge
Flooding frequency	None
Elevation	945–1,402 m
Slope	15–75%
Aspect	N, E, S

## Climatic features

Precipitation in this common resource area averages 12 to 16 inches annually. The winter-summer rainfall ratio ranges from about 60/40% in the northwest part of the area to 50/50% in the southeast part. Summer rains fall July through September; are from high-intensity, convective thunderstorms. This moisture originates primarily from the Gulf of Mexico, but can come from the remnants of Pacific hurricanes in September. Winter moisture is frontal, originates in the north Pacific, and falls as rain or snow in widespread storms of low intensity and long duration. Snowfall ranges from a trace to 10 inches per year and can occur from November through March. Snow seldom persists for more than a day except on north aspects. May and June are the driest months of the year. Humidity is generally low all year. Average annual air temperatures range from 59 to 70 degrees F (thermic temperature regime). Daytime temperatures in the summer are commonly in the high 90's. Freezing temperatures are common from October through April, usually during the night or early morning hours. The actual precipitation, available moisture and temperature vary, depending on, region, elevation, rain shadow effect and aspect.

Table 3. Representative climatic features

Frost-free period (average)	230 days
Freeze-free period (average)	285 days
Precipitation total (average)	406 mm

## Influencing water features

There are no water features associated with this site.

## Soil features

These soils are shallow (10 to 20 inches) and dark colored. They are loamy sand to coarse sandy loam in texture, calcareous in places and well drained. They have formed in residuum and colluvium from diabase and granodiorite parent materials. Bedrock has carbonates in the fractures. Soil surfaces are not well covered by gravels, cobbles and/or stones. Gravel size is inadequate to prevent soil erosion. The effective rooting depth is limited by slightly weathered bedrock at 10 to 20 inches. Runoff is moderate to high on moist soils. Rock outcrop and can be as high as 10%.

Soils mapped to date on this site include Schifflin.

**Table 4. Representative soil features**

Parent material	(1) Residuum–diabase (2) Colluvium–granodiorite
Surface texture	(1) Gravelly loamy sand (2) Very gravelly loamy sand (3) Very gravelly sandy loam
Family particle size	(1) Sandy
Drainage class	Well drained
Permeability class	Rapid to moderately rapid
Soil depth	25–51 cm
Surface fragment cover <=3"	5–30%
Surface fragment cover >3"	5–35%
Available water capacity (0-101.6cm)	1.52–3.81 cm
Calcium carbonate equivalent (0-101.6cm)	0–10%
Electrical conductivity (0-101.6cm)	0–2 mmhos/cm
Sodium adsorption ratio (0-101.6cm)	0–2
Soil reaction (1:1 water) (0-101.6cm)	7.4–8.4
Subsurface fragment volume <=3" (Depth not specified)	5–30%
Subsurface fragment volume >3" (Depth not specified)	0–15%

## Ecological dynamics

The historic native plant community is a diverse mixture of desert trees, shrubs, succulents, forbs and grasses. This includes a diverse flora of native annual grasses and forbs of both the winter and summer seasons. Periodic wildfires occurred at moderate intervals (15 to 30 years) and helped maintain a balance between herbs and shrubs. In the absence of fire for longer periods, shrubby species and cacti can become dominant. The interactions of drought, fire and continuous livestock grazing can, over time, result in the loss of palatable grasses, half shrubs and suffrutescent forbs. In some situations non-native annuals can dominate the site. These species can, over time, diminish the soil seed-bank of native annual species. Non-native annuals can act to increase the fire frequency of areas of the site near roads and urban areas, where the incidence of man-made fires is high.

## State and transition model

## **Ecosystem states**

1. Mixed Shrub-  
Herbaceous State

2. Shrub Dominated  
State

3. Shrub Dominated  
and Eroded State

4. Exotic Invaded State

## **State 1 submodel, plant communities**

1.1. Historic Native  
Plant Community

## **State 2 submodel, plant communities**

2.1. Shrub Dominated  
Plant Community

## **State 3 submodel, plant communities**

3.1. Shrub Dominated  
and Eroded Plant  
Community

## **State 4 submodel, plant communities**

4.1. Exotic Forb and  
Grass Plant  
Community

## **State 1**

### **Mixed Shrub-Herbaceous State**

#### **Community 1.1**

#### **Historic Native Plant Community**



Figure 4. Diabase Hills 12-16" pz. HCPC

The historic native plant community is a diverse mixture of perennial grasses, suffrutescent forbs, shrubs, succulents and desert trees. A rich flora of native annual forbs and grasses, of both the winter and summer seasons, exist in the plant community. Periodic, naturally occurring, wildfires were important in maintaining the potential plant community. North slopes have more evergreen shrubs like juniper and turbinella oak. Southern exposures will have a higher percentage of desert shrubs, trees and succulents in the plant community. More xeric grasses will dominate southern exposures (aristida, black grama, bush muhly). Grasses on cooler aspects include desert stipa and sideoats grama.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	112	280	628
Shrub/Vine	112	280	516
Forb	28	84	280
Tree	17	28	112
<b>Total</b>	<b>269</b>	<b>672</b>	<b>1536</b>

Table 6. Soil surface cover

Tree basal cover	0-1%
Shrub/vine/liana basal cover	1-2%
Grass/grasslike basal cover	2-5%
Forb basal cover	1-2%
Non-vascular plants	0%
Biological crusts	0-1%
Litter	20-50%
Surface fragments >0.25" and <=3"	5-30%
Surface fragments >3"	5-35%
Bedrock	1-10%
Water	0%
Bare ground	15-65%

Table 7. Canopy structure (% cover)

Height Above Ground (M)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.15	–	1-5%	0-10%	0-10%
>0.15 <= 0.3	–	5-10%	1-10%	1-5%
>0.3 <= 0.6	–	1-5%	1-5%	0-5%
>0.6 <= 1.4	–	5-15%	0-2%	0-1%
>1.4 <= 4	0-5%	1-5%	–	–
>4 <= 12	0-2%	–	–	–
>12 <= 24	–	–	–	–
>24 <= 37	–	–	–	–
>37	–	–	–	–

Figure 6. Plant community growth curve (percent production by month). AZ3811, 38.1 12-16" p.z. all sites. Growth begins in the spring, most growth occurs in the summer..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	1	7	15	20	22	20	10	5	0	0

## State 2 Shrub Dominated State

### Community 2.1 Shrub Dominated Plant Community



Figure 7. Diabase Hills 12-16" p.z. shrubby

Perennial grass canopy cover is reduced due to the interactions of drought, grazing and fire. Desert shrubs and cacti dominate the plant community. Shrub cover exceeds 40%. Annuals, both native and non-native, dominate the under-story. Fire frequency is reduced but the site can still burn, especially after "El Nino" years produce heavy fuel loads of annual grasses and forbs.

## State 3 Shrub Dominated and Eroded State

### Community 3.1 Shrub Dominated and Eroded Plant Community



Figure 8. Diabase Hills 12-16" pz. eroded

Shrubs like catclaw and whitethorn acacia; trees like mesquite, juniper, ocotillo, blue paloverde, and succulents like prickly pear, cholla and banana yucca can increase to dominate the site in the absence of fire for very long periods of time. Native and non-native annual forbs and grasses dominate the under-story. In "El Nino" years, herbaceous fuels can be sufficient to carry fire through the heavy canopy of shrubs. The major woody shrubs are, however, fire resistant once established. Such fires would remove less tolerant species like cacti and leave intact the sprouting woody plants to become more and more dominant. Extreme rainfall events coupled with; the fire, drought and grazing interaction, can lead to rilling of steep slopes. Compaction of soils can occur with heavy trailing from continuous livestock use. Loss of plant cover after repeated fire can lead to accelerated rill erosion under these circumstances. Small gravel sizes (1/4 inch) are inadequate to prevent soil movement on steep slopes when the plant cover has been depleted.

**State 4  
Exotic Invaded State**

**Community 4.1  
Exotic Forb and Grass Plant Community**

Non-native annual grasses and forbs like; red brome, cheatgrass, and wild oats, can invade and dominate areas of the site. These species can, over time, reduce the seed-bank of native annual grasses and forbs. Their presence can increase the fire frequency (of man made fires) especially where roads and urban areas are adjacent to areas of the site. Repeated fires tend to remove fire sensitive species like paloverde, cacti and juniper, and leave fire tolerant species like turbinella oak, mesquite, whitethorn, catclaw and mimosa.

**Additional community tables**

Table 8. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass/Grasslike					

## Grass/Grasslike

1	<b>Dominant perennial grasses</b>			106–336	
	black grama	BOER4	<i>Bouteloua eriopoda</i>	56–224	–
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	22–78	–
	purple threeawn	ARPU9	<i>Aristida purpurea</i>	6–56	–
	Parish's threeawn	ARPUP5	<i>Aristida purpurea</i> var. <i>parishii</i>	6–56	–
	bush muhly	MUPO2	<i>Muhlenbergia porteri</i>	11–56	–
	Arizona cottontop	DICA8	<i>Digitaria californica</i>	0–22	–
	tanglehead	HECO10	<i>Heteropogon contortus</i>	0–22	–
	spidergrass	ARTE3	<i>Aristida ternipes</i>	1–22	–
	spidergrass	ARTEG	<i>Aristida ternipes</i> var. <i>gentilis</i>	0–22	–
	cane bluestem	BOBA3	<i>Bothriochloa barbinodis</i>	0–11	–
2	<b>Cool season grasses</b>			1–56	
	desert needlegrass	ACSP12	<i>Achnatherum speciosum</i>	1–56	–
	squirreltail	ELEL5	<i>Elymus elymoides</i>	0–11	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	0–11	–
	Indian ricegrass	ACHY	<i>Achnatherum hymenoides</i>	0–6	–
3	<b>Misc. perennial grasses</b>			1–67	
	slender grama	BORE2	<i>Bouteloua repens</i>	0–34	–
	big galleta	PLRI3	<i>Pleuraphis rigida</i>	0–28	–
	hairy grama	BOHI2	<i>Bouteloua hirsuta</i>	0–22	–
	curly-mesquite	HIBE	<i>Hilaria belangeri</i>	0–11	–
	Hall's panicgrass	PAHA	<i>Panicum hallii</i>	0–11	–
	red grama	BOTR2	<i>Bouteloua trifida</i>	0–6	–
	low woollygrass	DAPU7	<i>Dasyochloa pulchella</i>	0–6	–
	southwestern bristlegrass	SESC2	<i>Setaria scheelei</i>	0–6	–
	plains bristlegrass	SEVU2	<i>Setaria vulpisetia</i>	0–6	–
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	0–6	–
	Fendler threeawn	ARPUL	<i>Aristida purpurea</i> var. <i>longiseta</i>	0–6	–
	blue threeawn	ARPUN	<i>Aristida purpurea</i> var. <i>nealleyi</i>	0–6	–
	nineawn pappusgrass	ENDE	<i>Enneapogon desvauxii</i>	0–2	–
	plains lovegrass	ERIN	<i>Eragrostis intermedia</i>	0–1	–
	vine mesquite	PAOB	<i>Panicum obtusum</i>	0–1	–
	tobosagrass	PLMU3	<i>Pleuraphis mutica</i>	0–1	–
	green sprangletop	LEDU	<i>Leptochloa dubia</i>	0–1	–
4	<b>Annual grasses</b>			6–168	
	sixweeks threeawn	ARAD	<i>Aristida adscensionis</i>	1–34	–
	mucronate sprangletop	LEPAB	<i>Leptochloa panicea</i> ssp. <i>brachiata</i>	1–28	–
	small fescue	VUMI	<i>Vulpia microstachys</i>	1–22	–
	Eastwood fescue	VUMIC	<i>Vulpia microstachys</i> var. <i>ciliata</i>	1–22	–
	sixweeks fescue	VUOC	<i>Vulpia octoflora</i>	1–22	–
	Mexican panicgrass	PAHI5	<i>Panicum hirticaule</i>	0–22	–
	prairie threeawn	AROL	<i>Aristida oligantha</i>	1–22	–
	Arizona signalgrass	URAR	<i>Urochloa arizonica</i>	0–17	–



	witchgrass	PACA6	<i>Panicum capillare</i>	0-6	-
	Arizona brome	BRAR4	<i>Bromus arizonicus</i>	0-2	-
	needle grama	BOAR	<i>Bouteloua aristidoides</i>	0-2	-
	sixweeks grama	BOBA2	<i>Bouteloua barbata</i>	0-1	-
	Rothrock's grama	BORO2	<i>Bouteloua rothrockii</i>	0-1	-
	feather fingergrass	CHVI4	<i>Chloris virgata</i>	0-1	-
	canyon cupgrass	ERLE7	<i>Eriochloa lemmonii</i>	0-1	-
	tufted lovegrass	ERPE	<i>Eragrostis pectinacea</i>	0-1	-
	desert lovegrass	ERPEM	<i>Eragrostis pectinacea</i> var. <i>miserrima</i>	0-1	-
	little barley	HOPU	<i>Hordeum pusillum</i>	0-1	-
	Mexican sprangletop	LEFUU	<i>Leptochloa fusca</i> ssp. <i>uninervia</i>	0-1	-
	delicate muhly	MUFR	<i>Muhlenbergia fragilis</i>	0-1	-
	littleseed muhly	MUMI	<i>Muhlenbergia microsperma</i>	0-1	-
	Bigelow's bluegrass	POBI	<i>Poa bigelovii</i>	0-1	-
<b>Forb</b>					
5	<b>Perennial forbs</b>			22-112	
	shrubby deervetch	LORI3	<i>Lotus rigidus</i>	1-22	-
	longflower tube tongue	JULO3	<i>Justicia longii</i>	0-11	-
	slender janusia	JAGR	<i>Janusia gracilis</i>	1-11	-
	lacy tansyaster	MAPI	<i>Machaeranthera pinnatifida</i>	0-11	-
	desert marigold	BAMU	<i>Baileya multiradiata</i>	1-11	-
	desert globemallow	SPAM2	<i>Sphaeralcea ambigua</i>	1-11	-
	brownplume wirelettuce	STPA4	<i>Stephanomeria pauciflora</i>	1-11	-
	Parry's false prairie-clover	MAPA7	<i>Marina parryi</i>	0-6	-
	perennial rockcress	ARPE2	<i>Arabis perennans</i>	0-6	-
	trailing windmills	ALIN	<i>Allionia incarnata</i>	1-6	-
	white sagebrush	ARLUM2	<i>Artemisia ludoviciana</i> ssp. <i>mexicana</i>	1-6	-
	wealeaf bur ragweed	AMCO3	<i>Ambrosia confertiflora</i>	1-6	-
	plains blackfoot	MELE2	<i>Melampodium leucanthum</i>	0-6	-
	wishbone-bush	MILAV	<i>Mirabilis laevis</i> var. <i>villosa</i>	1-6	-
	cliffbrake	PELLA	<i>Pellaea</i>	0-6	-
	Parry's beardtongue	PEPA24	<i>Penstemon parryi</i>	0-6	-
	desert penstemon	PEPS	<i>Penstemon pseudospectabilis</i>	0-6	-
	slender poreleaf	POGR5	<i>Porophyllum gracile</i>	0-6	-
	Coues' cassia	SECO10	<i>Senna covesii</i>	1-6	-
	climbing wartclub	BOSC	<i>Boerhavia scandens</i>	0-6	-
	lipfern	CHEIL	<i>Cheilanthes</i>	0-6	-
	desert trumpet	ERIN4	<i>Eriogonum inflatum</i>	1-6	-
	Mojave spurge	EUSC6	<i>Euphorbia schizoloba</i>	0-2	-
	southwestern mock vervain	GLGO	<i>Glandularia gooddingii</i>	0-2	-
	desert rosemallow	HICO	<i>Hibiscus coulteri</i>	0-2	-
	sego lily	CANU3	<i>Calochortus nuttallii</i>	0-2	-
	New Mexico groundsel	DANE7	<i>Desmodium neomexicanum</i>	0-2	-

	new mexican groundsel	PAQU8	<i>Packera neomexicana</i>	0-2	-
	Oak Creek ragwort	PAQU8	<i>Packera quercetorum</i>	0-1	-
	toadflax penstemon	PELI2	<i>Penstemon linarioides</i>	0-1	-
	turpentinebroom	THMO	<i>Thamnosma montana</i>	0-1	-
	branched noseburn	TRRA5	<i>Tragia ramosa</i>	0-1	-
	Louisiana vetch	VILUL2	<i>Vicia ludoviciana ssp. ludoviciana</i>	0-1	-
	Wright's deervetch	LOWR	<i>Lotus wrightii</i>	0-1	-
	wavyleaf Indian paintbrush	CAAPM	<i>Castilleja applegatei ssp. martinii</i>	0-1	-
	Arizona wrightwort	CAAR7	<i>Carlowrightia arizonica</i>	0-1	-
	ragged nettlespurge	JAMA	<i>Jatropha macrorhiza</i>	0-1	-
	paleface	HIDE	<i>Hibiscus denudatus</i>	0-1	-
	Indian rushpea	HOGL2	<i>Hoffmannseggia glauca</i>	0-1	-
	Cooley's bundleflower	DECO2	<i>Desmanthus cooleyi</i>	0-1	-
	desert larkspur	DEPA	<i>Delphinium parishii</i>	0-1	-
	tall mountain larkspur	DESC	<i>Delphinium scaposum</i>	0-1	-
	bluedicks	DICA14	<i>Dichelostemma capitatum</i>	0-1	-
	fleabane	ERIGE2	<i>Erigeron</i>	0-1	-
	spikemoss	SELAG	<i>Selaginella</i>	0-1	-
	Lemmon's ragwort	SELE8	<i>Senecio lemmonii</i>	0-1	-
	silverleaf nightshade	SOEL	<i>Solanum elaeagnifolium</i>	0-1	-
	glandleaf milkwort	POMA7	<i>Polygala macradenia</i>	0-1	-
	scurfpea	PSORA2	<i>Psoraleidium</i>	0-1	-
	canaigre dock	RUHY	<i>Rumex hymenosepalus</i>	0-1	-
	twinleaf senna	SEBA3	<i>Senna bauhinioides</i>	0-1	-
	orange fameflower	PHAU13	<i>Phemeranthus aurantiacus</i>	0-1	-
	desert tobacco	NIOB	<i>Nicotiana obtusifolia</i>	0-1	-
	tuber anemone	ANTU	<i>Anemone tuberosa</i>	0-1	-
	narrowleaf silverbush	ARLA12	<i>Argythamnia lanceolata</i>	0-1	-
	white sagebrush	ARLU	<i>Artemisia ludoviciana</i>	0-1	-
	dwarf desertpeony	ACNA2	<i>Acourtia nana</i>	0-1	-
	brownfoot	ACWR5	<i>Acourtia wrightii</i>	0-1	-
	San Felipe dogweed	ADPO	<i>Adenophyllum porophylloides</i>	0-1	-
	New Mexico silverbush	ARNE2	<i>Argythamnia neomexicana</i>	0-1	-
	largeflower onion	ALMA4	<i>Allium macropetalum</i>	0-1	-
	dense ayenia	AYMI	<i>Ayenia microphylla</i>	0-1	-
	scarlet spiderling	BOCO	<i>Boerhavia coccinea</i>	0-1	-
6	<b>Annual forbs</b>			6-168	
	bristly fiddleneck	AMTE3	<i>Amsinckia tessellata</i>	0-28	-
	Coulter's spiderling	BOCO2	<i>Boerhavia coulteri</i>	0-28	-
	fivewing spiderling	BOIN	<i>Boerhavia intermedia</i>	0-28	-
	California poppy	ESCAM	<i>Eschscholzia californica ssp. mexicana</i>	0-28	-
	western tansymustard	DEPI	<i>Descurainia pinnata</i>	0-22	-
	Coulter's lupine	LUSP2	<i>Lupinus sparsiflorus</i>	0-22	-
	rhacelia	PHACE	<i>Rhacelia</i>	0-17	-

	FRACE	FRACE	FRACE	FRACE	FRACE
	FRACE	FRACE	FRACE	FRACE	FRACE
exserted Indian paintbrush	CAEXE		<i>Castilleja exserta ssp. exserta</i>	0-17	-
pitseed goosefoot	CHBE4		<i>Chenopodium berlandieri</i>	0-11	-
slender goldenweed	MAGR10		<i>Machaeranthera gracilis</i>	0-11	-
tanseyleaf tansyaster	MATA2		<i>Machaeranthera tanacetifolia</i>	0-11	-
shaggyfruit pepperweed	LELA		<i>Lepidium lasiocarpum</i>	0-6	-
Arizona poppy	KAGR		<i>Kallstroemia grandiflora</i>	0-6	-
foothill deervetch	LOHU2		<i>Lotus humistratus</i>	0-6	-
desertparsley	LOMAT		<i>Lomatium</i>	0-6	-
coastal bird's-foot trefoil	LOSA		<i>Lotus salsuginosus</i>	0-6	-
Arizona lupine	LUAR4		<i>Lupinus arizonicus</i>	0-6	-
Arizona popcornflower	PLAR		<i>Plagiobothrys arizonicus</i>	0-6	-
sleepy silene	SIAN2		<i>Silene antirrhina</i>	0-6	-
lyreleaf jewelflower	STCA5		<i>Streptanthus carinatus</i>	0-6	-
New Mexico plumeseed	RANE		<i>Rafinesquia neomexicana</i>	0-6	-
combseed	PECTO		<i>Pectocarya</i>	0-6	-
desert Indianwheat	PLOV		<i>Plantago ovata</i>	1-6	-
woolly plantain	PLPA2		<i>Plantago patagonica</i>	0-6	-
milkvetch	ASTRA		<i>Astragalus</i>	0-6	-
carelessweed	AMPA		<i>Amaranthus palmeri</i>	0-6	-
American wild carrot	DAPU3		<i>Daucus pusillus</i>	0-6	-
cryptantha	CRYPT		<i>Cryptantha</i>	0-6	-
flatcrown buckwheat	ERDE6		<i>Eriogonum deflexum</i>	0-6	-
sorrel buckwheat	ERPO4		<i>Eriogonum polycladon</i>	0-6	-
thelypody	THELY		<i>Thelypodium</i>	0-6	-
woolly tidestromia	TILA2		<i>Tidestromia lanuginosa</i>	0-6	-
sand fringe-pod	THCU		<i>Thysanocarpus curvipes</i>	0-2	-
miniature woollystar	ERDI2		<i>Eriastrum diffusum</i>	0-2	-
spreading fleabane	ERDI4		<i>Erigeron divergens</i>	0-2	-
spurge	EUPHO		<i>Euphorbia</i>	0-2	-
New Mexico thistle	CINE		<i>Cirsium neomexicanum</i>	0-2	-
miner's lettuce	CLPEP		<i>Claytonia perfoliata ssp. perfoliata</i>	0-2	-
wedgeleaf draba	DRCU		<i>Draba cuneifolia</i>	0-2	-
annual agoseris	AGHE2		<i>Agoseris heterophylla</i>	0-2	-
purslane	PORTU		<i>Portulaca</i>	0-2	-
Gordon's bladderpod	LEGO		<i>Lesquerella gordonii</i>	0-2	-
Florida pellitory	PAFL3		<i>Parietaria floridana</i>	0-2	-
creamcups	PLCA5		<i>Platystemon californicus</i>	0-2	-
miniature lupine	LUBI		<i>Lupinus bicolor</i>	0-2	-
crestrub morning-glory	IPCO2		<i>Ipomoea costellata</i>	0-2	-
Thurber's pepperweed	LETH2		<i>Lepidium thurberi</i>	0-2	-
hollowleaf annual lupine	LUSU3		<i>Lupinus succulentus</i>	0-2	-
Fendler's desertyellow	MAFE		<i>Malacothrix fendleri</i>	0-1	-
whitestem blazingstar	MEAL6		<i>Mentzelia albicaulis</i>	0-1	-

	green carpetweed	MOVE	<i>Mollugo verticillata</i>	0-1	-
	desert evening primrose	OEPR	<i>Oenothera primiveris</i>	0-1	-
	redstar	IPCO3	<i>Ipomoea coccinea</i>	0-1	-
	ivyleaf morning-glory	IPHE	<i>Ipomoea hederacea</i>	0-1	-
	California goldfields	LACA7	<i>Lasthenia californica</i>	0-1	-
	woollyhead neststraw	STMI2	<i>Stylocline micropoides</i>	0-1	-
	chia	SACO6	<i>Salvia columbariae</i>	0-1	-
	sawtooth sage	SASU7	<i>Salvia subincisa</i>	0-1	-
	ragwort	SENEC	<i>Senecio</i>	0-1	-
	spreading fanpetals	SIAB	<i>Sida abutifolia</i>	0-1	-
	manybristle chinchweed	PEPA2	<i>Pectis papposa</i>	0-1	-
	desert unicorn-plant	PRAL4	<i>Proboscidea althaeifolia</i>	0-1	-
	doubleclaw	PRPA2	<i>Proboscidea parviflora</i>	0-1	-
	hoary bowlesia	BOIN3	<i>Bowlesia incana</i>	0-1	-
	brittle spineflower	CHBR	<i>Chorizanthe brevicornu</i>	0-1	-
	Esteve's pincushion	CHST	<i>Chaenactis stevioides</i>	0-1	-
	yellow tackstem	CAPA7	<i>Calycoseris parryi</i>	0-1	-
	white tackstem	CAWR	<i>Calycoseris wrightii</i>	0-1	-
	sacred thorn-apple	DAWR2	<i>Datura wrightii</i>	0-1	-
	scrambled eggs	COAU2	<i>Corydalis aurea</i>	0-1	-
	hairy prairie clover	DAMO	<i>Dalea mollis</i>	0-1	-
	star gilia	GIST	<i>Gilia stellata</i>	0-1	-
	longleaf false goldeneye	HELOA2	<i>Heliomeris longifolia var. annua</i>	0-1	-
	Texas stork's bill	ERTE13	<i>Erodium texanum</i>	0-1	-

#### Shrub/Vine

7	<b>Evergreen shrubs</b>			6-90	
	Sonoran scrub oak	QUTU2	<i>Quercus turbinella</i>	0-56	-
	narrowleaf yerba santa	ERAN2	<i>Eriodictyon angustifolium</i>	0-28	-
	desert ceanothus	CEGR	<i>Ceanothus greggii</i>	0-22	-
	sugar sumac	RHOV	<i>Rhus ovata</i>	0-6	-
	algerita	MATR3	<i>Mahonia trifoliolata</i>	0-6	-
	snapdragon penstemon	KEANM	<i>Keckiella antirrhinoides ssp. microphylla</i>	0-2	-
8	<b>Large shrubs</b>			34-168	
	catclaw acacia	ACGR	<i>Acacia greggii</i>	11-90	-
	Wright's beebrush	ALWR	<i>Aloysia wrightii</i>	6-56	-
	desert lavender	HYEM	<i>Hyptis emoryi</i>	1-50	-
	ocotillo	FOSP2	<i>Fouquieria splendens</i>	6-22	-
	whitethorn acacia	ACCO2	<i>Acacia constricta</i>	0-11	-
	winterfat	KRLA2	<i>Krascheninnikovia lanata</i>	0-6	-
	Berlandier's wolfberry	LYBE	<i>Lycium berlandieri</i>	0-6	-
	Arizona desert-thorn	LYEX	<i>Lycium exsertum</i>	0-6	-
	ambrosia leaf bur ragweed	AMAM2	<i>Ambrosia ambrosioides</i>	0-6	-
	catclaw mimosa	MIACB	<i>Mimosa aculeaticarpa var. hirsutifera</i>	0-6	-

			<i>Dunaliella</i>		
	Mexican bladdersage	SAME	<i>Salazaria mexicana</i>	0–2	–
	Arizona necklacepod	SOAR3	<i>Sophora arizonica</i>	0–2	–
	creosote bush	LATR2	<i>Larrea tridentata</i>	0–2	–
	pale desert-thorn	LYPA	<i>Lycium pallidum</i>	0–1	–
	Thurber's desert honeysuckle	ANTH2	<i>Anisacanthus thurberi</i>	0–1	–
	fourwing saltbush	ATCA2	<i>Atriplex canescens</i>	0–1	–
	desertbroom	BASA2	<i>Baccharis sarothroides</i>	0–1	–
	spiny hackberry	CEEH	<i>Celtis ehrenbergiana</i>	0–1	–
	desert sweet	CHMI2	<i>Chamaebatiaria millefolium</i>	0–1	–
	Warnock's snakewood	COWA	<i>Condalia warnockii</i>	0–1	–
	Florida hopbush	DOVI	<i>Dodonaea viscosa</i>	0–1	–
	lotebush	ZIOBC	<i>Ziziphus obtusifolia</i> var. <i>canescens</i>	0–1	–
	jojoba	SICH	<i>Simmondsia chinensis</i>	0–1	–
	mariola	PAIN2	<i>Parthenium incanum</i>	0–1	–
	skunkbush sumac	RHTR	<i>Rhus trilobata</i>	0–1	–
9	<b>Dominant half shrubs</b>			39–135	
	fairyduster	CAER	<i>Calliandra eriophylla</i>	22–78	–
	bastardsage	ERWR	<i>Eriogonum wrightii</i>	6–28	–
	littleleaf ratany	KRER	<i>Krameria erecta</i>	1–17	–
	rough menodora	MESC	<i>Menodora scabra</i>	0–11	–
	Parish's goldeneye	VIPA14	<i>Viguiera parishii</i>	0–11	–
	Eastern Mojave buckwheat	ERFA2	<i>Eriogonum fasciculatum</i>	0–11	–
	longleaf phlox	PHLO2	<i>Phlox longifolia</i>	0–6	–
	American threefold	TRCA8	<i>Trixis californica</i>	0–6	–
	Coulter's brickellbush	BRCO	<i>Brickellia coulteri</i>	0–6	–
	desert zinnia	ZIAC	<i>Zinnia acerosa</i>	0–2	–
	starry bedstraw	GAST	<i>Galium stellatum</i>	0–2	–
	ragged rockflower	CRBI2	<i>Crossosoma bigelovii</i>	0–1	–
	shortleaf baccharis	BABR	<i>Baccharis brachyphylla</i>	0–1	–
	yerba de pasmo	BAPT	<i>Baccharis pteronioides</i>	0–1	–
	sweetbush	BEJU	<i>Bebbia juncea</i>	0–1	–
10	<b>Succulents</b>			22–90	
	cactus apple	OPEN3	<i>Opuntia engelmannii</i>	6–45	–
	banana yucca	YUBA	<i>Yucca baccata</i>	1–22	–
	tulip pricklypear	OPPH	<i>Opuntia phaeacantha</i>	1–11	–
	common sotol	DAWH2	<i>Dasyllirion wheeleri</i>	1–11	–
	walkingstick cactus	CYSP8	<i>Cylindropuntia spinosior</i>	0–6	–
	saguaro	CAGI10	<i>Carnegiea gigantea</i>	0–6	–
	buck-horn cholla	CYAC8	<i>Cylindropuntia acanthocarpa</i>	0–6	–
	candy barrelcactus	FEWI	<i>Ferocactus wislizeni</i>	0–2	–
	Graham's nipple cactus	MAGR9	<i>Mammillaria grahamii</i>	0–1	–
	sacahuista	NOMI	<i>Nolina microcarpa</i>	0–1	–
	dollarplant pricklypear	OPCH	<i>Opuntia oblongifolia</i>	0–1	–

	collar joint pricklypear	OPCH	<i>Opuntia chiroloa</i>	0-1	-
	teddybear cholla	CYBI9	<i>Cylindropuntia bigelovii</i>	0-1	-
	jumping cholla	CYFU10	<i>Cylindropuntia fulgida</i>	0-1	-
	Christmas cactus	CYLE8	<i>Cylindropuntia leptocaulis</i>	0-1	-
	goldenflower century plant	AGCH2	<i>Agave chrysantha</i>	0-1	-
	Schott's century plant	AGSC3	<i>Agave schottii</i>	0-1	-
	Whipple cholla	CYWH	<i>Cylindropuntia whipplei</i>	0-1	-
	pinkflower hedgehog cactus	ECBO2	<i>Echinocereus bonkeriae</i>	0-1	-
	Arizona hedgehog cactus	ECCOA	<i>Echinocereus coccineus var. arizonicus</i>	0-1	-
	Engelmann's hedgehog cactus	ECEN	<i>Echinocereus engelmannii</i>	0-1	-
	redspine fishhook cactus	ECER2	<i>Echinomastus erectocentrus</i>	0-1	-
	pinkflower hedgehog cactus	ECFA	<i>Echinocereus fasciculatus</i>	0-1	-
	spiny star	ESVI2	<i>Escobaria vivipara</i>	0-1	-
11	<b>Increaser half-shrubs</b>			6-56	
	broom snakeweed	GUSA2	<i>Gutierrezia sarothrae</i>	1-34	-
	brittlebush	ENFA	<i>Encelia farinosa</i>	0-17	-
	button brittlebush	ENFR	<i>Encelia frutescens</i>	0-17	-
	turpentine bush	ERLA12	<i>Ericameria laricifolia</i>	0-6	-
	narrowleaf goldenbush	ERLI6	<i>Ericameria linearifolia</i>	0-6	-
	threadleaf snakeweed	GUMI	<i>Gutierrezia microcephala</i>	0-1	-
	burroweed	ISTE2	<i>Isocoma tenuisecta</i>	0-1	-
<b>Tree</b>					
12	<b>Trees</b>			17-112	
	blue paloverde	PAFL6	<i>Parkinsonia florida</i>	11-56	-
	crucifixion thorn	CAHO3	<i>Canotia holacantha</i>	0-45	-
	yellow paloverde	PAMI5	<i>Parkinsonia microphylla</i>	1-17	-
	western honey mesquite	PRGLT	<i>Prosopis glandulosa var. torreyana</i>	0-6	-
	velvet mesquite	PRVE	<i>Prosopis velutina</i>	0-6	-
	redberry juniper	JUCO11	<i>Juniperus coahuilensis</i>	0-6	-
	oneseed juniper	JUMO	<i>Juniperus monosperma</i>	0-6	-

## Animal community

This site is suitable for grazing year round, but is not easily traversed by livestock. Livestock grazing use is concentrated on south slopes, canyon bottoms and ridge-tops. North slopes may be little used. Slopes greater than 50% and areas with very cobbly surfaces limit grazing use by cattle. Areas of rock outcrop can form barriers to livestock movement. The site is very susceptible to erosion in overgrazed areas like bed-grounds, livestock trails and lower slopes adjacent to water, because gravel size is inadequate in preventing soil loss.

The site has good habitat diversity for a great variety of desert wildlife species. Water developments are very important to both livestock and wildlife on this site.

## Hydrological functions

This site has smooth surfaces, due to a high cover of fine gravels with some cobbles and stone. When the soils are dry, it produces limited runoff. It produces significant runoff when heavy rain falls on snow or moist soils.

## Recreational uses

Hunting, camping, horseback riding, backpacking, rock hounding, photography.

## Wood products

Limited harvest of fuel-wood, fence posts and stays from mesquite, juniper and saguaro.

## Other products

There is some native harvest of food plants like thistle, prickly pear tunas and mescal. There is limited harvest of herbs like herbaceous sage, yerba santa and mormon tea.

## Type locality

Location 1: Pinal County, AZ	
Township/Range/Section	T7S R18E S31
General legal description	Miller Dry Camp Allotment, below Table Mountain

## Contributors

Dan Robinett  
Larry D. Ellicott

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

1. **Number and extent of rills:**

---

2. **Presence of water flow patterns:**

---

3. **Number and height of erosional pedestals or terracettes:**

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):**
- 
5. **Number of gullies and erosion associated with gullies:**
- 
6. **Extent of wind scoured, blowouts and/or depositional areas:**
- 
7. **Amount of litter movement (describe size and distance expected to travel):**
- 
8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**
- 
9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**
- 
10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**
- 
11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**
- 
12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**
- Dominant:
- Sub-dominant:
- Other:
- Additional:
- 
13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**
- 
14. **Average percent litter cover (%) and depth ( in):**
- 
15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**



---

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:**

---

17. **Perennial plant reproductive capability:**

---