

Ecological site R038XB107NM Clayey

Accessed: 05/19/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

The topography of this site is level to moderately sloping and may include slightly depressed or swale-type positions which receive runoff from adjacent sites. Slopes range from 0 to 10 percent, but average less than 5 percent. Elevations range from just under 5,000 feet to just over 6,800 feet above sea level.

Landforms	(1) Depression(2) Swale
Flooding duration	Very brief (4 to 48 hours)
Flooding frequency	Rare to frequent
Elevation	1,524–2,073 m

Table 2. Representative physiographic features

Slope	0–10%
Aspect	Aspect is not a significant factor

Climatic features

Average annual precipitation varies from about 12 inches to just over 16 inches. Substantial fluctuations from year to year are common, ranging from a low of about 6 inches to a high of over 30 inches. Approximately one-half of the annual precipitation comes in the form of rainfall during the months of July, August, and September, although wintertime precipitation in the form of snow, sleet, or rain is sometimes significant. Spring and late fall months are normally dry.

The average frost-free period ranges from about 165 to 190 days and extends from approximately the third or fourth week in April to mid October. Average annual air temperatures are about 56 degrees F. Summer maximums can exceed 100 degrees F and winter minimums on occasion go below zero. Monthly mean temperatures generally exceed 70 degrees F for the period of June through August.

Growing conditions favor warm-season perennial vegetation, although late winter and late summer precipitation is adequate to foster a significant cool-season component in the potential plant community. Occasional wet springs also create good conditions for annual forb production, but frequent winds from the west and southwest are common during this time of year and tend to deplete soil moisture at a critical time for the growth of these plants.

Climate data was obtained from http://www.wrcc.sage.dri.edu/summary/climsmnm.html web site using 50% probability for freeze-free and frost-free seasons using 28.5 degrees F and 32.5 degrees F respectively.

Table 3. Representative climatic features

Frost-free period (average)	187 days
Freeze-free period (average)	211 days
Precipitation total (average)	406 mm

Influencing water features

This site is not influenced by water from a wetland or stream.

Soil features

Typical soils are moderately deep to deep well drained. The surface layer is fine or moderately fine. Sometimes it has a very thin layer that is medium textured and are slowly permeable. Water intake rates are moderate to slow. Soil cracking following dry periods provides opportunity for occasional deep wetting upon finally receiving moisture. Runoff in the absence of good vegetative cover can be high, and water-holding capacity is moderate to high.

-	
Surface texture	(1) Gravelly clay loam (2) Loam
Family particle size	(1) Clayey
Drainage class	Well drained
Permeability class	Slow
Soil depth	51–183 cm
Surface fragment cover <=3"	15–35%
Available water capacity (0-101.6cm)	7.62–30.48 cm

Table 4. Representative soil features

Electrical conductivity (0-101.6cm)	0–16 mmhos/cm
Soil reaction (1:1 water) (0-101.6cm)	6.6–9
Subsurface fragment volume <=3" (Depth not specified)	0–57%

Ecological dynamics

Retrogression in the plant community is characterized by increases in such plants as tobosa, threeawns, burrograss, and mesquite. Broom snakeweed and an abundance of annuals may also be symptomatic of site deterioration, especially if there are numerous areas of bare ground and an obvious absence or near absence of the better climax species such as the gramas. Following severe deterioration, the site does not recover rapidly through improved grazing management alone.

State and transition model



State 1 Historic Climax Plant Community

Community 1.1 Historic Climax Plant Community

This is a grassland site characterized by short and mid-grasses. Blue grama and sideoats grama are the dominant grasses. Such species as black grama, alkali sacaton, tobosa, and bottlebrush squirreltail are also characteristic

although alkali sacaton may not be present in all cases. Fourwing saltbush, winterfat, and yucca are common woody species, with broom snakeweed coming and going cyclically and as condition deteriorates. Forbs such as perennial buckwheat and globemallow also occur commonly on the site but not often in substantial amounts. NOTE: Where site occurs in depressed areas or shallow swale positions, additional overflow may cause the site to support as much as 15 percent western wheatgrass, 20 percent tobosa or galleta, and 10 percent vine-mesquite. Under these conditions, blue grama normally will not exceed 20 percent.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Grass/Grasslike	286	560	834
Forb	27	53	78
Total	313	613	912

Table 6. Ground cover

Tree foliar cover	0%
Shrub/vine/liana foliar cover	4%
Grass/grasslike foliar cover	0%
Forb foliar cover	0%
Non-vascular plants	0%
Biological crusts	0%
Litter	12%
Litter Surface fragments >0.25" and <=3"	12% 0%
Litter Surface fragments >0.25" and <=3" Surface fragments >3"	12% 0% 0%
Litter Surface fragments >0.25" and <=3" Surface fragments >3" Bedrock	12% 0% 0% 0%
Litter Surface fragments >0.25" and <=3" Surface fragments >3" Bedrock Water	12% 0% 0% 0%

Figure 5. Plant community growth curve (percent production by month). NM0607, R038XB107NM Clayey HCPC. R038XB107NM Clayey HCPC Mixed short/mid-grassland with shrub and forb component..

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	5	7	10	15	25	25	8	5	0	0

Additional community tables

 Table 7. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
Grass	/Grasslike				
1				197–263	
	blue grama	BOGR2	Bouteloua gracilis	197–263	-
2				66–99	
	sideoats grama	BOCU	Bouteloua curtipendula	66–99	-
3				99–132	
	James' galleta	PLJA	Pleuraphis jamesii	99–132	-
	tobosagrass	PLMU3	Pleuraphis mutica	99–132	-
	alkali sacaton	SPAI	Sporobolus airoides	99–132	_

4				7–33	
	black grama	BOER4	Bouteloua eriopoda	7–33	-
5			•	7–33	
	vine mesquite	PAOB	Panicum obtusum	7–33	_
6		•	•	20–33	
	squirreltail	ELEL5	Elymus elymoides	20–33	_
7		•	•	7–33	
	cane bluestem	BOBA3	Bothriochloa barbinodis	7–33	_
	Arizona cottontop	DICA8	Digitaria californica	7–33	-
	plains lovegrass	ERIN	Eragrostis intermedia	7–33	-
	plains bristlegrass	SEVU2	Setaria vulpiseta	7–33	-
8		-		7–20	
	low woollygrass	DAPU7	Dasyochloa pulchella	7–20	-
	curly-mesquite	HIBE	Hilaria belangeri	7–20	-
	creeping muhly	MURE	Muhlenbergia repens	7–20	-
	mat muhly	MURI	Muhlenbergia richardsonis	7–20	-
9		-		20–53	
	threeawn	ARIST	Aristida	20–53	-
	burrograss	SCBR2	Scleropogon brevifolius	20–53	-
10				7–20	
	Graminoid (grass or grass- like)	2GRAM	Graminoid (grass or grass- like)	7–20	_
Forb	•	•	•	•	
11				7–20	
	croton	CROTO	Croton	7–20	-
	buckwheat	ERIOG	Eriogonum	7–20	_
	globemallow	SPHAE	Sphaeralcea	7–20	_
12		•	•	7–20	
	Forb, annual	2FA	Forb, annual	7–20	_
13			·	7–33	
	Forb, perennial	2FP	Forb, perennial	7–33	-
Shrub	/Vine				
14				7–20	
	fourwing saltbush	ATCA2	Atriplex canescens	7–20	-
15				7–20	
	longleaf jointfir	EPTR	Ephedra trifurca	7–20	-
	pale desert-thorn	LYPA	Lycium pallidum	7–20	-
	уисса	YUCCA	Yucca	7–20	-
16				7–20	
	winterfat	KRLA2	Krascheninnikovia lanata	7–20	_
17				7–20	
	broom snakeweed	GUSA2	Gutierrezia sarothrae	7–20	

Animal community

This ecological site provides habitat which can support a resident animal community characterized by desert cottontail, Botta's pocket gopher, bannertail kangaroo rat, southern plains woodrat, coyote, burrowing owl, meadowlark, scaled quail, lesser earless lizard, leopard lizard, and short-horned lizard.

Where shrubs are present, mourning dove nest.

Hydrological functions

The runoff curve numbers are determined by field investigations using hydrologic cover conditions and hydrologic soil groups.

Hydrologic Interpretations Soil Series------Hydrologic Group Carnero-----C Denver-----C Manzano------B Navajo------D Ruidoso-----C White House-----C

Recreational uses

This site offers limited recreation potential for hiking, horseback riding, picnicking, camping, nature observation, photography, and hunting for scaled quail and mourning dove. When favorable spring moisture conditions occur, a colorful display of wildflowers may be seen.

Wood products

This site has no significant value for wood products.

Other products

Grazing: This site, at its potential, is suitable for grazing in all seasons of the year, since most of its production is in the form of perennial grasses which persist as dry-standing herbage well after their growing season has ended. It is adapted for cattle, sheep, horses, and to some extent goats, generally without regard to class of livestock. Retrogression in the plant community is characterized by increases in such plants as tobosa, threeawns, burrograss, and mesquite. Broom snakeweed and an abundance of annuals may also be symptomatic of site deterioration, especially if there are numerous areas of bare ground and an obvious absence or near absence of the better climax species such as the gramas. Following severe deterioration, the site does not recover rapidly through improved grazing management alone.

Other information

Guide to Suggested Initial Stocking Rate Acres per Animal Unit Month

Similarity Index	Ac/AUM
100 - 76	4.0 – 5.0
75 – 51	4.8 – 7.2
50 – 26	7.0 – 13.0
25 – 0	13.0+

Type locality

Location 1: Grant County, NM Location 2: Catron County, NM Location 3: Hidalgo County, NM Location 4: Sierra County, NM

Location 5: Socorro County, NM

Other references

Data collection for this site was done in conjunction with the progressive soil surveys within the New Mexico and Arizona Plateaus and Mesas 36 Major Land Resource Area of New Mexico. This site has been mapped and correlated with soils in the following soil surveys: Socorro, Sierra, Grant, Catron.

Characteristic Soils Are: Denver, Ruidoso

Other Soils included are: Carnero, Lonti, Manzano, Navajo, White House

Contributors

Brandon Bestelmeyer Don Sylvester Elizabeth Wright John Tunberg

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- 1. Number and extent of rills:
- 2. Presence of water flow patterns:
- 3. Number and height of erosional pedestals or terracettes:

4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):

- 5. Number of gullies and erosion associated with gullies:
- 6. Extent of wind scoured, blowouts and/or depositional areas:
- 7. Amount of litter movement (describe size and distance expected to travel):
- 8. Soil surface (top few mm) resistance to erosion (stability values are averages most sites will show a range of values):
- 9. Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):
- 10. Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:
- 11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):
- 12. Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):

Dominant:

Sub-dominant:

Other:

Additional:

- 13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):
- 14. Average percent litter cover (%) and depth (in):
- 15. Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annualproduction):
- 16. Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize

degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:

17. Perennial plant reproductive capability: