

Ecological site R038XB209AZ **Loamy Upland 16-20" p.z.**

Last updated: 5/07/2020
 Accessed: 02/17/2025

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

MLRA notes

Major Land Resource Area (MLRA): 038X–Mogollon Transition South

AZ 38-2 Middle Mogollon Transition

Elevations range from 4200 to 5800 feet and precipitation averages 16 to 20 inches per year in this Land Resource Unit. Vegetation includes turbinella oak, Wright silktassel, hollyleaf buckthorn, desert buckbrush, one-seed juniper, alligator juniper, pinyon, algerita, sugar sumac, prairie junegrass, blue grama, curly mesquite, bottlebrush squirreltail, muttongrass, cane beardgrass, plains lovegrass and bullgrass. The soil temperature regime ranges from thermic to mesic and the soil moisture regime is aridic ustic. This unit occurs within the Transition Zone Physiographic Province and is characterized by canyons and structural troughs or valleys. Igneous, metamorphic and sedimentary rock classes occur on rough mountainous terrain in association with less extensive sediment filled valleys exhibiting little integrated drainage.

Associated sites

R038XB202AZ	Clayey Upland 16-20" p.z. Clayey Upland 16-20" p.z.
R038XB203AZ	Clay Loam Upland 16-20" p.z. Clayloam Upland 16-20" p.z.
R038XB215AZ	Clayey Hills 16-20" p.z. Clayey Hills 16-20" p.z.

Similar sites

R038XC303AZ	Clay Loam Upland 20-24" p.z. Clayloam Upland 20+" p.z. will have trace amounts of oneseed, Utah, and redberry juniper but the savanna tree species is alligator juniper and when the site has been converted to woodland through juniper invasion alligator juniper is the dominant tree.
-------------	---

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

This site occurs in the mid to upper elevations of the Mogollon Transition zone south of the Mogollon Rim in central Arizona.

This site is in an upland position. It neither benefits significantly from run-in moisture nor suffers from excess runoff. Slopes range from 0 to 15 percent on valley fill plains, basalt flows and mesa tops.

Table 2. Representative physiographic features

Landforms	(1) Alluvial flat (2) Mesa
Flooding frequency	None
Ponding frequency	None
Elevation	4,200–5,800 ft
Slope	0–15%
Aspect	N, S, W

Climatic features

Precipitation in this Land Resource Unit averages 16 to 20 inches annually. The winter-summer rainfall ratio ranges from about 60/40% in the western part of the area to 45/55% in the eastern part. Summer rains fall July through September; and are from high-intensity, convective, thunderstorms. This moisture originates primarily from the Gulf of Mexico, but can come from the remnants of Pacific hurricanes in September. Winter moisture is frontal, originates in the north Pacific, and falls as rain or snow in widespread storms of low intensity and long duration. Snowfall ranges from 5 to 35 inches per year and can occur from November through April. Snow seldom persists for more than a week. May and June are the driest months of the year. Humidity is moderate to low all year. Average annual air temperatures range from 51 to 60 degrees F (thermic temperature regime). Daytime temperatures in the summer are commonly in the low 90's. Freezing temperatures are common from October through April. The actual precipitation, available moisture and temperature vary, depending on, region, elevation, rain shadow effect and aspect.

Table 3. Representative climatic features

Frost-free period (average)	180 days
Freeze-free period (average)	240 days
Precipitation total (average)	20 in




Figure 1. Monthly precipitation range




Figure 2. Monthly average minimum and maximum temperature

Influencing water features

There are no water features associated with this site.

Soil features

Soils mapped under this site include: SSA675 San Carlos IR area MU's 8 Stanford and Lanque, 11 Biplane family, 26 & 81 Terrarossa.

Table 4. Representative soil features

Parent material	(1) Alluvium–andesite
Surface texture	(1) Gravelly loam (2) Gravelly sandy loam (3) Loam
Family particle size	(1) Clayey
Drainage class	Well drained to moderately well drained
Permeability class	Very slow to moderately rapid
Soil depth	30–60 in
Surface fragment cover <=3"	0–25%
Surface fragment cover >3"	0–15%
Available water capacity (0-40in)	2.6–7.9 in
Calcium carbonate equivalent (0-40in)	0–10%
Electrical conductivity (0-40in)	0–2 mmhos/cm
Sodium adsorption ratio (0-40in)	0–2
Soil reaction (1:1 water) (0-40in)	6.6–8.4
Subsurface fragment volume <=3" (Depth not specified)	0–30%
Subsurface fragment volume >3" (Depth not specified)	0–15%

Ecological dynamics


The plant communities found on an ecological site are naturally variable. Composition and production will vary with yearly conditions, location, aspect, and the natural variability of the soils. The Reference Plant Community

represents the natural potential plant communities found on relict or relatively undisturbed sites. Other plant communities described here represent plant communities that are known to occur when the site is disturbed by factors such as grazing, fire, absence of fire, or drought.

Production data provided in this site description is standardized to air-dry weight at the end of the summer growing season. The plant communities described in this site description are based on near normal rainfall years.

NRCS uses a Similarity Index to compare existing plant communities to the plant communities described here. Similarity Index is determined by comparing the production and composition of a plant community to the production and composition of a plant community described in this site description. To determine Similarity Index, compare the production (air-dry weight) of each species to that shown in the plant community description. For each species, count no more than the maximum amount shown for the species, and for each group, count no more than the maximum shown for the group. Divide the resulting total by the total normal year production shown in the plant community description. If rainfall has been significantly above or below normal, use the total production shown for above or below normal years. If field data is not collected at the end of the summer growing season, then the field data must be corrected to the end of the year production before comparing it to the site description. The growth curve can be used as a guide for estimating production at the end of the summer growing season.

State and transition model



State 1

Reference Plant Community

The Midgrass State is the Reference Plant Community. It is a warm season dominated grassland (canopy cover of 55 to 65%) with an important component of cool season grasses that fluctuate with climate. Cool and warm season annual grasses and forbs are well represented in the flora. Half shrubs and perennial forbs are an important group also. Fire return intervals are difficult to determine in woody plant communities in this LRU. However, periodic wildfires likely occurred every 10 to 15 years from June through August, and controlled shrubs and succulents invading from adjacent, shallow soil, areas. In the absence of fire for long periods of time; shrubs, trees and cacti can dominate the site. The interactions of drought, fire and grazing can result in a loss of perennial grass cover. In these situations annuals, both native and non-native, can dominate the plant community. Non-native annuals can, over time, diminish the soil seed-bank of native annual species.

Community 1.1

Reference Plant Community



The Reference Plant Community is a grassland dominated by sideoats grama and blue grama. Prairie junegrass and bottlebrush squirreltail are important in the plant community, but can diminish to low levels after severe winter - spring drought. Squirreltail appears to be slightly less susceptible to drought than junegrass. Shrubby buckwheat is an important half-shrub in the plant community. A flora of native annual forbs and grasses, of both the winter and summer seasons, exist in the plant community. Periodic, naturally occurring, wildfires were important in maintaining the potential plant community.

Table 5. Annual production by plant type

Plant Type	Low (Lb/Acre)	Representative Value (Lb/Acre)	High (Lb/Acre)
Grass/Grasslike	700	1500	1900
Forb	20	150	400
Shrub/Vine	25	50	75
Tree	0	0	25
Total	745	1700	2400

Table 6. Soil surface cover

Tree basal cover	0-1%
Shrub/vine/liana basal cover	0-1%
Grass/grasslike basal cover	10-15%
Forb basal cover	0-1%
Non-vascular plants	0%

Biological crusts	0%
Litter	60-80%
Surface fragments >0.25" and <=3"	0-25%
Surface fragments >3"	0-15%
Bedrock	0%
Water	0%
Bare ground	0-15%

Table 7. Canopy structure (% cover)

Height Above Ground (Ft)	Tree	Shrub/Vine	Grass/ Grasslike	Forb
<0.5	—	0-1%	5-10%	0-5%
>0.5 <= 1	—	0-1%	10-20%	0-15%
>1 <= 2	—	0-1%	30-50%	—
>2 <= 4.5	—	0-1%	—	—
>4.5 <= 13	—	0-1%	—	—
>13 <= 40	—	—	—	—
>40 <= 80	—	—	—	—
>80 <= 120	—	—	—	—
>120	—	—	—	—




Figure 4. Plant community growth curve (percent production by month). AZ3812, 38.2 16-20" p.z. all sites. Growth begins in the spring and continues into the summer and fall..

State 2 Short grass State

The Short grass state has a plant community dominated by short grasses.

Community 2.1 Blue grama high basal Community



The Blue Grama high basal community is dominated by blue grama that has basal area ranging from 10-15%. Proper livestock grazing in the form of periodic summer deferment or low intensity to no livestock grazing maintains high basal area.

Community 2.2

Blue grama medium basal Community

The Blue Grama medium basal Community is dominated by blue grama that has basal area ranging from 5-10%. Continuous livestock grazing with little to no summer deferment has reduced the vigor of blue grama to a point where its basal area begins to decline.

Pathway 2.1A

Community 2.1 to 2.2

Continuous heavy livestock grazing with no summer deferment.

Pathway 2.2a

Community 2.2 to 2.1

Application of prescribed grazing in the form of summer deferment.

Conservation practices

Prescribed Grazing

State 3

Short grass / subshrub State

The Short grass / subshrub State is a community where high levels of snakeweed have established and are co-dominant with blue grama.

Community 3.1

Blue grama medium basal / subshrub Community



The Blue grama medium basal Community of this state is a community where high levels of snakeweed have established and are now co-dominant with blue grama. Continuous livestock grazing with no summer deferment has reduced the vigor of blue grama to approximately 5-10% basal area and opened the community up to invasion by high densities of snakeweed. Snakeweed canopy is approximately 30-40%.

State 4 Eroded State

The eroded state is a plant community where blue grama abundance has been substantially reduced.

Community 4.1 Blue grama low basal Community

Continuous heavy livestock grazing and drought have substantially reduced the vigor of blue grama. Basal area of blue grama is reduced to less than 5% basal area. Plants are pedestalled with approximately 0.5-1" of the root exposed.

State 5 Juniper State

Juniper has increased in the absence of fire to dominate the site.

Community 5.1 Juniper Community



Juniper has established in the absence of fire to canopy levels of 15-50%. Herbaceous production ranges from 0-400 lbs/ac. The dominant juniper species is oneseed or redberry juniper. This is likely a stable state because trees are of sufficient size that very high temperatures and wind speeds are required to burn the site; conditions where

practitioners are unlikely to apply prescribed fire. Mechanical brush management can be applied followed by burning of skeletons if sufficient fine fuels are on site. Perennial grass canopy in the interspaces of trees is > 50% in some areas with existing plants likely providing an adequate seed source for restoration following brush management or fires. Range planting should likely only be considered where perennial grass canopy in the interspaces of trees is less than 25%; however this threshold needs additional investigation.

Transition T1a

State 1 to 2

Continuous livestock grazing with no summer deferrment.

Restoration pathway R2a

State 2 to 1

Unknown. Contour ripping to break up the dominance of blue grama and allow other species to come in has been successful in blue grama dominated plant communities in central New Mexico (Pat Shaver, pers. comm.) and squirreltail is only observed in areas where pipelines have been ripped through blue grama communities north of Springerville, AZ. Contour ripping followed by range planting.

Transition T2a

State 2 to 3

Continuous heavy livestock grazing with no summer deferrment.

Transition T2b

State 2 to 5

Juniper establishment and growth to maturity in the absence of fire.

Restoration pathway R3a

State 3 to 2

Unknown. Contour ripping, range planting, prescribed grazing in the form of summer deferrment.

Transition T3a

State 3 to 4

Continued yearlong heavy livestock grazing with no summer deferrment.

Restoration pathway R4a

State 4 to 3

Long term application of prescribed grazing in the form of summer deferrment. This restoration pathway likely only possible where soil erosion is less than 1/2 inch and very long term application of summer deferrment.

Restoration pathway R5a

State 5 to 2

Mechanical brush management with sufficient remnant perennial grass in the interspaces between trees. Reseeding may be needed where perennial grass is less than 25% canopy cover in the interspaces between trees. However, this needs additional investigation.

Additional community tables

Table 8. Community 1.1 plant community composition

Group	Common Name	Symbol	Scientific Name	Annual Production (Lb/Acre)	Foliar Cover (%)
-------	-------------	--------	-----------------	-----------------------------	------------------

Grass/Grasslike					
1	mid grasses			400–700	
	sideoats grama	BOCU	<i>Bouteloua curtipendula</i>	400–700	–
	cane bluestem	BOBA3	<i>Bothriochloa barbinodis</i>	0–100	–
	plains lovegrass	ERIN	<i>Eragrostis intermedia</i>	0–50	–
	green sprangletop	LEDU	<i>Leptochloa dubia</i>	0–50	–
2	short grasses			100–400	
	blue grama	BOGR2	<i>Bouteloua gracilis</i>	100–300	–
	hairy grama	BOHI2	<i>Bouteloua hirsuta</i>	0–100	–
	sprucetop grama	BOCH	<i>Bouteloua chondrosioides</i>	0–100	–
	black grama	BOER4	<i>Bouteloua eriopoda</i>	0–100	–
3	cool season grasses			10–200	
	squirreltail	ELEL5	<i>Elymus elymoides</i>	10–200	–
	prairie Junegrass	KOMA	<i>Koeleria macrantha</i>	10–200	–
	muttongrass	POFE	<i>Poa fendleriana</i>	0–10	–
4	miscellaneous perennial grasses			10–50	
	spidergrass	ARTEG	<i>Aristida ternipes</i> var. <i>gentilis</i>	0–50	–
	threeawn	ARIST	<i>Aristida</i>	0–25	–
	Fendler threeawn	ARPUL	<i>Aristida purpurea</i> var. <i>longiseta</i>	0–25	–
	spidergrass	ARTE3	<i>Aristida ternipes</i>	0–25	–
	sand dropseed	SPCR	<i>Sporobolus cryptandrus</i>	0–25	–
	common wolfstail	LYPH	<i>Lycurus phleoides</i>	0–15	–
	Arizona threeawn	ARAR6	<i>Aristida arizonica</i>	0–15	–
	poverty threeawn	ARDI5	<i>Aristida divaricata</i>	0–10	–
	creeping muhly	MURE	<i>Muhlenbergia repens</i>	0–10	–
	slender muhly	MUTE4	<i>Muhlenbergia tenuifolia</i>	0–10	–
	curly-mesquite	HIBE	<i>Hilaria belangeri</i>	0–10	–
5	annual grasses			25–300	
	Mexican sprangletop	LEFUU	<i>Leptochloa fusca</i> ssp. <i>uninervia</i>	10–200	–
	mucronate sprangletop	LEPAB	<i>Leptochloa panicea</i> ssp. <i>brachiata</i>	10–200	–
	Mexican panicgrass	PAHI5	<i>Panicum hirticaule</i>	5–200	–
	prairie threeawn	AROL	<i>Aristida oligantha</i>	5–100	–
	needle grama	BOAR	<i>Bouteloua aristidoides</i>	0–50	–
	sixweeks threeawn	ARAD	<i>Aristida adscensionis</i>	0–50	–
	feather fingergrass	CHVI4	<i>Chloris virgata</i>	0–50	–
	witchgrass	PACA6	<i>Panicum capillare</i>	0–50	–
	tufted lovegrass	ERPE	<i>Eragrostis pectinacea</i>	0–50	–
	desert lovegrass	ERPEM	<i>Eragrostis pectinacea</i> var. <i>miserrima</i>	0–50	–
	sticky sprangletop	LEVI5	<i>Leptochloa viscida</i>	0–50	–
	small fescue	VUMI	<i>Vulpia microstachys</i>	0–50	–
	Eastwood fescue	VUMIC	<i>Vulpia microstachys</i> var. <i>ciliata</i>	0–50	–
	sixweeks fescue	VUOC	<i>Vulpia octoflora</i>	5–50	–
	Bigelow's bluegrass	POBI	<i>Poa bigelovii</i>	0–25	–
	delicate muhly	MUFR	<i>Muhlenbergia fragilis</i>	0–25	–

	littleseed muhly	MUMI	<i>Muhlenbergia microsperma</i>	0–25	–
	little barley	HOPU	<i>Hordeum pusillum</i>	0–25	–
	canyon cupgrass	ERLE7	<i>Eriochloa lemmonii</i>	0–25	–
	sixweeks grama	BOBA2	<i>Bouteloua barbata</i>	0–25	–
	Arizona brome	BRAR4	<i>Bromus arizonicus</i>	0–25	–
	Arizona signalgrass	URAR	<i>Urochloa arizonica</i>	0–15	–
Forb					
6	perennial forbs			15–70	
	desert globemallow	SPAM2	<i>Sphaeralcea ambigua</i>	1–15	–
	copper globemallow	SPAN3	<i>Sphaeralcea angustifolia</i>	0–15	–
	Lewis flax	LILE3	<i>Linum lewisii</i>	0–10	–
	bluedicks	DICA14	<i>Dichelostemma capitatum</i>	0–10	–
	Forb, perennial	2FP	<i>Forb, perennial</i>	0–10	–
	weakleaf bur ragweed	AMCO3	<i>Ambrosia confertiflora</i>	1–10	–
	brownfoot	ACWR5	<i>Acourtia wrightii</i>	0–5	–
	largeflower onion	ALMA4	<i>Allium macropetalum</i>	1–5	–
	scarlet spiderling	BOCO	<i>Boerhavia coccinea</i>	0–5	–
	leastdaisy	CHAET2	<i>Chaetopappa</i>	0–5	–
	rose heath	CHER2	<i>Chaetopappa ericoides</i>	0–5	–
	sego lily	CANU3	<i>Calochortus nuttallii</i>	0–2	–
	onion	ALLIU	<i>Allium</i>	0–2	–
	Greene's bird's-foot trefoil	LOGR4	<i>Lotus greenei</i>	0–2	–
	Wright's deervetch	LOWR	<i>Lotus wrightii</i>	0–2	–
	Parry's beardtongue	PEPA24	<i>Penstemon parryi</i>	0–2	–
	orange fameflower	PHAU13	<i>Phemeranthus aurantiacus</i>	0–2	–
	twinleaf senna	SEBA3	<i>Senna bauhinoides</i>	0–2	–
	ragwort	SENEC	<i>Senecio</i>	0–2	–
	brownplume wirelettuce	STPA4	<i>Stephanomeria pauciflora</i>	0–2	–
	southwestern mock vervain	GLGO	<i>Glandularia gooddingii</i>	0–2	–
	vetch	VICIA	<i>Vicia</i>	0–1	–
	beeblossom	GAURA	<i>Gaura</i>	0–1	–
	tuber anemone	ANTU	<i>Anemone tuberosa</i>	0–1	–
	perennial rockcress	ARPE2	<i>Arabis perennans</i>	0–1	–
	Watson's dutchman's pipe	ARWA	<i>Aristolochia watsonii</i>	0–1	–
	Texas bindweed	COEQ	<i>Convolvulus equitans</i>	0–1	–
	desert larkspur	DEPA	<i>Delphinium parishii</i>	0–1	–
7	annual forbs			0–150	
	longleaf false goldeneye	HELO6	<i>Heliomeris longifolia</i>	1–100	–
	Forb, annual	2FA	<i>Forb, annual</i>	0–25	–
	western tansymustard	DEPI	<i>Descurainia pinnata</i>	0–10	–
	Arizona popcornflower	PLAR	<i>Plagiobothrys arizonicus</i>	1–10	–
	creamcups	PLCA5	<i>Platystemon californicus</i>	0–10	–

spreading fanpetals	SIAB	<i>Sida abutifolia</i>	0-5	-
Arizona lupine	LUAR4	<i>Lupinus arizonicus</i>	0-5	-
miniature lupine	LUBI	<i>Lupinus bicolor</i>	0-5	-
Coulter's lupine	LUSP2	<i>Lupinus sparsiflorus</i>	0-5	-
hollowleaf annual lupine	LUSU3	<i>Lupinus succulentus</i>	0-5	-
California poppy	ESCAM	<i>Eschscholzia californica</i> ssp. <i>mexicana</i>	0-5	-
common sunflower	HEAN3	<i>Helianthus annuus</i>	0-5	-
Coulter's spiderling	BOCO2	<i>Boerhavia coulteri</i>	0-5	-
spreading fleabane	ERDI4	<i>Erigeron divergens</i>	0-5	-
bristly fiddleneck	AMTE3	<i>Amsinckia tessellata</i>	0-5	-
aster	ASTER	<i>Aster</i>	0-5	-
New Mexico thistle	CINE	<i>Cirsium neomexicanum</i>	0-5	-
milkvetch	ASTRA	<i>Astragalus</i>	0-2	-
sorrel buckwheat	ERPO4	<i>Eriogonum polycladon</i>	0-2	-
lambsquarters	CHAL7	<i>Chenopodium album</i>	0-2	-
spurge	EUPHO	<i>Euphorbia</i>	0-2	-
cudweed	GNAPH	<i>Gnaphalium</i>	0-2	-
tanseyleaf tansyaster	MATA2	<i>Machaeranthera tanacetifolia</i>	0-2	-
desert Indianwheat	PLOV	<i>Plantago ovata</i>	0-2	-
woolly plantain	PLPA2	<i>Plantago patagonica</i>	0-2	-
sleepy silene	SIAN2	<i>Silene antirrhina</i>	0-1	-
sand fringe pod	THCU	<i>Thysanocarpus curvipes</i>	0-1	-
goldeneye	VIGUI	<i>Viguiera</i>	0-1	-
green carpetweed	MOVE	<i>Mollugo verticillata</i>	0-1	-
evening primrose	OENOT	<i>Oenothera</i>	0-1	-
lacy tansyaster	MAPI	<i>Machaeranthera pinnatifida</i>	0-1	-
aridland goosefoot	CHDE	<i>Chenopodium desiccatum</i>	0-1	-
sanddune wallflower	ERCA14	<i>Erysimum capitatum</i>	0-1	-
miniature woollystar	ERDI2	<i>Eriastrum diffusum</i>	0-1	-
New Mexico fleabane	ERNE3	<i>Erigeron neomexicanus</i>	0-1	-
crestrub morning-glory	IPCO2	<i>Ipomoea costellata</i>	1	-
California goldfields	LACA7	<i>Lasthenia californica</i>	0-1	-
grassleaf lettuce	LAGRA	<i>Lactuca graminifolia</i> var. <i>arizonica</i>	0-1	-
Goodding's bladderpod	LEGO2	<i>Lesquerella gooddingii</i>	0-1	-
shaggyfruit pepperweed	LELA	<i>Lepidium lasiocarpum</i>	0-1	-
Thurber's pepperweed	LETH2	<i>Lepidium thurberi</i>	0-1	-
foothill deervetch	LOHU2	<i>Lotus humistratus</i>	0-1	-
coastal bird's-foot trefoil	LOSA	<i>Lotus salsuginosus</i>	0-1	-
scrambled eggs	COAU2	<i>Corydalis aurea</i>	0-1	-
croton	CROTO	<i>Croton</i>	0-1	-
cryptantha	CRYPT	<i>Cryptantha</i>	0-1	-
American wild carrot	DAPU3	<i>Daucus pusillus</i>	0-1	-
annual agoseris	AGHE2	<i>Agoseris heterophylla</i>	0-1	-

Shrub/Vine					
8	shrubs			0–10	
	catclaw acacia	ACGR	<i>Acacia greggii</i>	0–5	–
	velvet mesquite	PRVE	<i>Prosopis velutina</i>	0–5	–
	fourwing saltbush	ATCA2	<i>Atriplex canescens</i>	0–2	–
	pale desert-thorn	LYPA	<i>Lycium pallidum</i>	0–1	–
	catclaw mimosa	MIACB	<i>Mimosa aculeaticarpa var. biuncifera</i>	0–1	–
	Sonoran scrub oak	QUTU2	<i>Quercus turbinella</i>	0–1	–
	currant	RIBES	<i>Ribes</i>	0–1	–
9	half shrubs			10–40	
	broom snakeweed	GUSA2	<i>Gutierrezia sarothrae</i>	1–25	–
	bastardsage	ERWR	<i>Eriogonum wrightii</i>	5–20	–
	prairie acacia	ACAN	<i>Acacia angustissima</i>	1–5	–
	yerba de pasmo	BAPT	<i>Baccharis pteronioides</i>	1–5	–
	Cooley's bundleflower	DECO2	<i>Desmanthus cooleyi</i>	1–5	–
10	succulents			0–100	
	sacahuista	NOMI	<i>Nolina microcarpa</i>	0–100	–
	cactus apple	OPEN3	<i>Opuntia engelmannii</i>	0–5	–
	soaptree yucca	YUEL	<i>Yucca elata</i>	0–5	–
	walkingstick cactus	CYSP8	<i>Cylindropuntia spinosior</i>	0–1	–
	Whipple cholla	CYWH	<i>Cylindropuntia whipplei</i>	0–1	–
Tree					
12	trees			0–25	
	redberry juniper	JUCO11	<i>Juniperus coahuilensis</i>	0–25	–
	oneseed juniper	JUMO	<i>Juniperus monosperma</i>	0–25	–
	twoneedle pinyon	PIED	<i>Pinus edulis</i>	0–25	–
	singleleaf pinyon	PIMO	<i>Pinus monophylla</i>	0–25	–
	Utah juniper	JUOS	<i>Juniperus osteosperma</i>	0–15	–

Animal community

This site is suitable for grazing year round and is easily traversed by all classes of livestock. The site is very susceptible to erosion in overgrazed areas, old roads, cattle trails and concentration areas like bed grounds, water-lots and salt grounds.

The site has good habitat diversity for grassland wildlife species. Where it is adjacent to hill sites, with tree species or chaparral, it is a foraging area for elk.

Hydrological functions

This site produces runoff when soils are moist. Surfaces can be easily compacted by traffic and high densities of livestock when soils are moist. Normal depth of soil freezing in the winter is 5 to 6 inches. This will not break up compacted layers deeper than that. Compacted surfaces will produce much more runoff than surfaces with good tilth and structure.

Recreational uses

Hunting, camping, hiking, horseback riding, and backpacking.

Wood products

None.

Other products

There is some native harvest of foods like wild onion, sunflower and thistle.

Type locality

Location 1: Yavapai County, AZ	
Township/Range/Section	T23 R1E S15N
UTM zone	N
UTM northing	3837140
UTM easting	387689
General legal description	Between Prescott Valley and Jerome. Approximately 1700 feet north of Hwy 89A and 1.8 miles west of the base of Hickey Mountain. 5,400 feet elevation. Minor component among more dominant Clayey Upland 38-2.

Contributors

Dave Womack
Larry D. Ellicott
Steve Barker

Approval

Scott Woodall, 5/07/2020

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Dave Womack, Dan Robinett, Emilio Carrillo
Contact for lead author	USDA NRCS, Tucson Area Office
Date	04/29/2011
Approved by	Scott Woodall
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- Number and extent of rills:** None present on the site. Perennial bunch grass plants are approximately 1-2 foot apart, promote high infiltration, and make the site not conducive to rill formation.

- Presence of water flow patterns:** Flow patterns are indiscrete and highly sinuous. Water flows no more than 1-2 feet

before encountering base of perennial bunch grass plant.

3. **Number and height of erosional pedestals or terracettes:** None present on the site. Bunch grass community not conducive to forming terracettes.

4. **Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 5-15%.

5. **Number of gullies and erosion associated with gullies:** None present on the site. Very dense bunchgrass community promotes high infiltration and very stable soils.

6. **Extent of wind scoured, blowouts and/or depositional areas:** None present on the site. Perennial bunchgrass canopy cover is 55-65%.

7. **Amount of litter movement (describe size and distance expected to travel):** Herbaceous litter is the dominant litter on the site. Litter moves no more than 1-2 feet before being intercepted by plant bases.

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Soil surface is highly resistant to erosion. Expect soil stability values of 4-6 across the site.

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Strong very fine granular structure. Soil organic matter content is 1-3%. A horizon is 4-8" thick with 7.5YR4/3 dry color and 7.5YR3/3 moist color.

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Very high density bunch grass dominated plant community promotes very high infiltration. Perennial grass densities are approximately 3-5 plants per square yard.

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None present on the site. Dry argillic horizon at 4-8 inches can be mistaken for compaction layer.

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: Midgrasses >> shortgrasses

Sub-dominant: Cool season grass > miscellaneous grasses > Perennial forbs = subshrubs > annual forbs = annual grass
(Note: In El Nino years annual grasses = annual forbs > misc. grass > perennial forbs).

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Approximately 2-5% mortality of perennial grass plants.
-

14. **Average percent litter cover (%) and depth (in):**
-

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** Below average year = 745 lb/ac, normal year = 1,700 lbs/ac, above average year 2,400 lbs/ac.
-

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** With continuous heavy livestock grazing blue grama will become dominant. Deterioration of blue grama cover will open community up to spiny tansyaster, snakeweed, annual goldeneye, and tumble mustard. Absence of fire can result in increase in beargrass and juniper.
-

17. **Perennial plant reproductive capability:** Not affected after several years of regional drought.
-