

Ecological site F039XA110AZ Limestone Hills 17-22" p.z. (PIPO, JUDE2)

Last updated: 9/05/2019 Accessed: 05/21/2024

General information

Provisional. A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.



Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

MLRA notes

Major Land Resource Area (MLRA): 039X–Mogollon Transition North

AZ 39.1 Mogollon Plateau Coniferous Forests

Elevations range from 7000 to 12,500 feet and precipitation averages 20 to 35 inches per year. Vegetation includes ponderosa pine, Gambel oak, Arizona walnut, sycamore, Douglas fir, blue spruce, Arizona fescue, sheep fescue, mountain muhly, muttongrass, junegrass, pine dropseed, and dryland sedges. The soil temperature regime ranges from mesic to frigid and the soil moisture regime ranges from typic ustic to udic ustic. This unit occurs within the Colorado Plateau Physiographic Province and is characterized by a sequence of flat to gently dipping sedimentary rocks eroded into plateaus, valleys and deep canyons. Sedimentary rock classes dominate the plateau with volcanic fields occurring for the most part near its margin.

Table 1. Dominant plant species

Tree	(1) Pinus ponderosa(2) Juniperus deppeana
Shrub	(1) Robinia neomexicana
Herbaceous	Not specified

Physiographic features

This site is mapped on hills and mountains.

Table 2. Representative physiographic features

Landforms	(1) Hill (2) Mountain
Elevation	1,829–2,563 m
Slope	25–70%
Aspect	Aspect is not a significant factor

Climatic features

About 40% of the moisture in this Common Resource Area (CRA), or Land Resource Unit (LRU) comes as rain from June to September. The remainder comes from October to May as snow or light rain. Extreme temperatures of 97 and -37 degrees Fahrenheit have been recorded. Some moisture is usually received every month.

Table 3. Representative climatic features

Frost-free period (average)	168 days
Freeze-free period (average)	120 days
Precipitation total (average)	559 mm

Influencing water features

Soil features

Soils mapped on this site include: from SSA-695 Navajo Depot MU 6-Huachuca family.

Table 4. Representative soil features

Surface texture	(1) Very stony loam
Drainage class	Moderately well drained to well drained
Permeability class	Moderately slow to moderate
Soil depth	8–20 cm
Surface fragment cover <=3"	0–10%
Surface fragment cover >3"	10–35%
Calcium carbonate equivalent (0-101.6cm)	0–5%

Ecological dynamics

The plant communities found on an ecological site are naturally variable. Composition and production will vary with yearly conditions, location, aspect, and the natural variability of the soils. The historical climax plant community represents the natural potential plant communities found on relict or relatively undisturbed sites. Other plant communities described here represent plant communities that are known to occur when the site is disturbed by factors such as grazing, fire, or drought.


Production data provided in this site description is standardized to air-dry weight at the end of the summer growing season. The plant communities described in this site description are based on near normal rainfall years.

NRCS uses a Similarity Index to compare existing plant communities to the plant communities described here.

Similarity Index is determined by comparing the production and composition of a plant community to the production and composition of a plant community described in this site description. To determine Similarity Index, compare the production (air-dry weight) of each species to that shown in the plant community description. For each species, count no more than the maximum amount shown for the species, and for each group, count no more than the maximum shown for the group. Divide the resulting total by the total normal year production shown in the plant community description. If rainfall has been significantly above or below normal, use the total production shown for above or below normal years. If field data is not collected at the end of the summer growing season, then the field data must be corrected to the end of the year production before comparing it to the site description. The growth curve can be used as a guide for estimating production at the end of the summer growing season.

State and transition model

Ecosystem states



State 1 Ponderosa Pine, mature Alligator Juniper

Mature Ponderosa Pine, mature alligator juniper, shrubs, and grasses.

State 2 Immature alligator juniper

Immature alligator juniper

Transition T1 State 1 to 2

Disturbance such as sever fire will cause a thick stand of immature alligator juniper.

Restoration pathway R1 State 2 to 1

A setback to alligator juniper along with re-colonization of ponderosa pine.

Contributors

Karlynn Huling Larry D. Ellicott

Approval

Scott Woodall, 9/05/2019

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

,	author(s)/participant(s)				
Со	Contact for lead author				
Da	Pate				
Аp	pproved by				
Аp	pproval date				
Со	Composition (Indicators 10 and 12) based on Annua	al Production			
	dicators . Number and extent of rills:				
	Presence of water flow patterns:				
	. Number and height of erosional pedestals or terracettes:				
	Bare ground from Ecological Site Description bare ground):	or other stud	ies (rock, litte	r, lichen, moss,	, plant canopy are not
5.	Number of gullies and erosion associated with	ı gullies:			
6.	Extent of wind scoured, blowouts and/or depo	sitional areas	s :		
7.	. Amount of litter movement (describe size and	distance expe	ected to trave	l):	
	Soil surface (top few mm) resistance to erosio values):	n (stability va	alues are aver	ages - most site	es will show a range of
9.	Soil surface structure and SOM content (include	de type of str	ucture and A-	horizon color a	nd thickness):
	Effect of community phase composition (relating distribution on infiltration and runoff:	ive proportior	n of different f	unctional grou	ps) and spatial
	Presence and thickness of compaction layer (mistaken for compaction on this site):	usually none;	describe soil	profile features	s which may be

Author(s)/participant(s)

12.	Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):		
	Dominant:		
	Sub-dominant:		
	Other:		
	Additional:		
13.	Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):		
14.	Average percent litter cover (%) and depth (in):		
15.	Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):		
16.	Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:		
17.	Perennial plant reproductive capability:		