

Ecological site R040XA121AZ Granitic Upland 10"-13" p.z.

Accessed: 12/06/2024

Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	Dave Womack, Dan Robinett, Emilio Carrillo
Contact for lead author	NRCS Tucson Area Office
Date	03/08/2005
Approved by	S Cassidy
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

Indicators

- Number and extent of rills:** Rills are present on this site but follow joints, bedding planes, and fractures in the bedrock parent materials.

- Presence of water flow patterns:** Discontinuous, 10-15 feet in length.

- Number and height of erosional pedestals or terracettes:** No pedestals on plants, rock or gravel fragments and no terracettes are present.

- Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not bare ground):** 10-60% (low values due to high rock and gravel cover).

- Number of gullies and erosion associated with gullies:** None

- Extent of wind scoured, blowouts and/or depositional areas:** None

- Amount of litter movement (describe size and distance expected to travel):** Most litter size classes stay in place.

Herbaceous litter moves 10-15 feet in flow paths.

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** Expect ratings of 1-3 in perennial plant interspaces, 4-5 under shrub canopies.

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):** Weak thin platy to granular; color is 7.5-10YR5/4 dry, 7.5-10YR4/4 moist; thickness to 2 inches.

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** Canopy 15-20%; 50-65% shrubs, 15-20 trees, 5-10% succulents, 2-5% perennial grasses. Cove is well dispersed throughout site.

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant: sub shrubs > trees > winter annuals > shrubs > perennial forbs > perennial grasses > summer annuals > succulents

Sub-dominant:

Other:

Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** 20-50% tree & shrub canopy mortality; 75-90% perennial grass mortality

14. **Average percent litter cover (%) and depth (in):**

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):** 122 lbs/ac unfavorable precipitation; 400 lbs/ac normal precipitation; 780 lbs/ac favorable precipitation.

16. **Potential invasive (including noxious) species (native and non-native). List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:** triangle bursage, turpentine bush, snakeweed, cacti, bufflegass

17. **Perennial plant reproductive capability:** Not impaired for shrubs, drought impaired for perennial grasses and forbs.
