

Ecological site R042BB003NM

Vegetated Gypsum Dunes, Desert Shrub

Accessed: 05/19/2024

General information

**Provisional.** A provisional ecological site description has undergone quality control and quality assurance review. It contains a working state and transition model and enough information to identify the ecological site.

Figure 1. Mapped extent

Areas shown in blue indicate the maximum mapped extent of this ecological site. Other ecological sites likely occur within the highlighted areas. It is also possible for this ecological site to occur outside of highlighted areas if detailed soil survey has not been completed or recently updated.

Associated sites

R042BB005NM	<b>Gyp Interdune (Dry), Desert Shrub</b> Vegetated Gypsum Dunes site occurs in association with the the Gyp Interdune(dry)site.
R042BB002NM	<b>Gyp Duneland Barren, Desert Shrub</b> This site occurs at the outer edges of the Gyp Duneland Barren site.

Table 1. Dominant plant species

Tree	Not specified
Shrub	Not specified
Herbaceous	Not specified

Physiographic features

This site occurs on gypsum dunefields. Typically the dune type is parabolic dunes, where the arms of the dune point upwind, having become stabilized and anchored by vegetation. Slope ranges from 5 to 60 percent. Elevations range from 3800 to 4500 feet above sea level.

Table 2. Representative physiographic features

Landforms	(1) Parabolic dune (2) Dune
Flooding frequency	None
Ponding frequency	None
Elevation	1,158–1,372 m
Slope	5–35%
Aspect	Aspect is not a significant factor

Climatic features

Annual average precipitation ranges from 7 to 12 inches. Wide fluctuations from year to year are common. At least one-half of the annual precipitation comes in the form of rainfall during July, August, and September. Precipitation in the form of snow or sleet averages less than 4 inches annually. The average annual air temperature is about 60 degree F. Summer maximums can exceed 100 degrees F. and winter minimums can go below zero. The average

frost-free season exceeds 200 days and extends from April 1 to November 1. Both the temperature regime and rainfall distribution favor warm-season perennial plants on this site. Spring moisture conditions are only occasionally adequate to cause significant growth during this period of year. High winds from the west and southwest are common from March to June, which further tends to create poor soil moisture conditions in the springtime

Climate data was obtained from  
<http://www.wrcc.dri.edu/summary/climsmnm.html>

**Table 3. Representative climatic features**

Frost-free period (average)	205 days
Freeze-free period (average)	227 days
Precipitation total (average)	305 mm

## Influencing water features

### Soil features

The soils of this site occur as parabolic gypsum dunes. Gypsum is present in high amounts (60-90%) throughout the soil profile. Surface and underlying layers are white gypsiferous sand. Cover of biological crusts ranges from 5 to 30 percent, with most occurring on the lower dune skirts where the soils are more stable. Soils on these dune types are only typically only slightly saline.

NM-688 MU-421 Duneland-Lark Assoc. 5-60% slopes

NM-688 MU-422 Firebee-Lark Assoc. 0-35% slopes

**Table 4. Representative soil features**

Surface texture	(1) Gypsiferous sand
Family particle size	(1) Sandy
Drainage class	Excessively drained
Permeability class	Very rapid
Soil depth	152 cm
Surface fragment cover <=3"	0%
Surface fragment cover >3"	0%
Available water capacity (0-101.6cm)	5.08–7.62 cm
Electrical conductivity (0-101.6cm)	1–3 mmhos/cm
Subsurface fragment volume <=3" (Depth not specified)	0%
Subsurface fragment volume >3" (Depth not specified)	0%

## Ecological dynamics


Overview:

The Vegetated Gypsum Dunes site occurs primarily on parabolic dunes. The parabolic dunes are usually found near the outer edges of the barchan dune fields.

Several theories exist as to why the parabolic dunes are stabilized by vegetation. One idea is that the parabolic dunes are situated topographically higher than the Barchanoid dunes and a perched freshwater lens sits above the saline water table of the barchanoid dune types. This fresh water lens promotes the establishment of vegetation that might be excluded by more saline conditions. Others attribute the establishment and subsequent stabilization by

vegetation to decreased gypsum sand supply at the periphery of the dune fields. This decreased sand supply and associated changes in the rate of sand transport makes the dunes inherently less mobile and more stable. The predominant historic community for this site is believed to be a shrubland usually dominated by frosted mint. Varying amounts of gyp grama and or sandhill muhly occur as sub dominants scattered within the shrub interspaces or in isolated patches. In some instances typically on the taller dunes a community dominated by cottonwood may exist. Each of these communities is susceptible to the invasion and in some instances domination by saltcedar.

## State and transition model



### State 1 Shrubland

This state is dominated by shrubs, but can have a fair component of mixed grasses scattered across the site.

### Community 1.1 Shrub-Mixed grass



Figure 4. Shrub/mixed grass community 1.1

This is the more common community type within the vegetated dune system. Shrubs are dominant and are scattered amid the dunes usually occupying the summits of the dune. Frosted mint is typically the dominant species. Other shrubs can include skunkbush sumac, soap tree yucca, ephedra, fourwing saltbush, broom dalea, and rubber rabbitbrush. Grasses are usually the subordinate component scattered between shrubs. Sandhill muhly or gyp grama is the dominant grass species. Sandhill muhly seems to be more prevalent on the more active dunes and on dunes with coarser textured surface sands. Gypsum grama usually dominates on smaller dunes and is most prolific near the interface between the dune and the interdune. Other grasses can include Indian ricegrass, New Mexico bluestem, and mesa dropseed. In some areas rubber rabbitbrush may be the dominant shrub with sparse amounts of alkali sacaton present.

Table 5. Annual production by plant type

Plant Type	Low (Kg/Hectare)	Representative Value (Kg/Hectare)	High (Kg/Hectare)
Shrub/Vine	123	252	331
Grass/Grasslike	50	112	219
Forb	11	28	45
Total	184	392	595

State 2  
Cottonwood Grove

This state is characterized by the dominance of cottonwoods.

Community 2.1  
Cottonwood-Mixed Shrubs



Figure 6. Cottonwood/Mixed shrubs community 2.1

The plant community of this state is dominated by Rio Grande cottonwoods. Other species include frosted mint, rabbitbrush, skunkbush sumac, alkali sacaton and Indian ricegrass. It occurs infrequently in isolated small groves. It tends to inhabit the higher dunes amid the parabolic dunefields and occasionally on barchanoid type dunes. One theory as to why cottonwoods are able to survive in the midst of the dunes is they are located along a perched freshwater lens above the more saline groundwater. The trunks of the cottonwood can elongate relatively fast, and they can also generate adventitious shoots. These aid in oxygen supply and distribution which assists the attempt to stay ahead of dune sand burial rates.

### **State 3**

#### **Exotic-Invaded**

This state is characterized by the presence of saltcedar. The occurrence of a relatively shallow water table promotes conditions that could allow for the establishment of saltcedar. The Exotic Invaded and the Exotic Dominated state currently tend to occur where vegetated dunes are located near gyp playa sites. It is not clear if the transition is more easily facilitated at these locations due to hydrologic differences or the proximity to a ready seed source. It is presumed however that this transition could occur throughout the vegetated dunes ecosite.

### **Community 3.1**

#### **Shrubs/Grass/Saltcedar**



Figure 7. Shrubs/Grass/Saltcedar community 3.1

This community is characterized by the presence of saltcedar amid mixed shrubs and scattered grasses. Shrubs typically include rabbitbrush, frosted mint, and skunkbush sumac. Grass species for this community typically include alkali sacaton, sandhill muhly, and Indian ricegrass.

### **Community 3.2**

#### **Cottonwood/Saltcedar**

This community is dominated by Rio Grande cottonwoods, but has occasional saltcedar present. Grasses include alkali sacaton, Indian ricegrass, and sandhill muhly. Shrubs typically include frosted mint, rabbitbrush, and skunkbush sumac.

### **State 4**

#### **Exotic-Dominated**

This state is characterized by the predominance of saltcedar.

### **Community 4.1**

#### **Saltcedar**



**Figure 8. Saltcedar community 4.1**

High seed production and the ability of saltcedar to exploit suitable habitat makes the site susceptible to the establishment of dense localized stands among the dunes.

**Additional community tables**

**Table 6. Community 1.1 plant community composition**

Group	Common Name	Symbol	Scientific Name	Annual Production (Kg/Hectare)	Foliar Cover (%)
<b>Grass/Grasslike</b>					
1	<b>Warm season short grasses</b>			22–112	
	gypsum grama	BOBR	<i>Bouteloua breviseta</i>	22–112	–
2	<b>Cool season short grasses</b>			11–17	
	Hopi tea greenthread	THME	<i>Thelesperma megapotamicum</i>	6–17	–
3	<b>warm season mid grasses</b>			17–90	
	sandhill muhly	MUPU2	<i>Muhlenbergia pungens</i>	17–84	–
	mesa dropseed	SPFL2	<i>Sporobolus flexuosus</i>	0–34	–
	alkali sacaton	SPAI	<i>Sporobolus airoides</i>	0–17	–
	spike dropseed	SPCO4	<i>Sporobolus contractus</i>	0–6	–
<b>Forb</b>					
4	<b>Forbs perennial</b>			11–34	
	greenthread	THELE	<i>Thelesperma</i>	11–17	–
	purple sand verbena	ABAN	<i>Abronia angustifolia</i>	0–11	–
	Hartweg's sundrops	CAHA14	<i>Calylophus hartwegii</i>	0–6	–
	White Sands fanmustard	NELI	<i>Nerisyrenia linearifolia</i>	0–6	–
	pale evening primrose	OEPA	<i>Oenothera pallida</i>	0–3	–
	sand milkweed	ASAR	<i>Asclepias arenaria</i>	0–3	–
5	<b>Forbs annual</b>			0–6	
	Forb, annual	2FA	<i>Forb, annual</i>	0–6	–
	blazingstar	MENTZ	<i>Mentzelia</i>	0–6	–
<b>Shrub/Vine</b>					
6	<b>Shrubs</b>			90–252	
	frosted mint	POIN3	<i>Poliomintha incana</i>	90–168	–
	skunkbush sumac	RHTR	<i>Rhus trilobata</i>	0–56	–
	Torrey's jointfir	EPTO	<i>Ephedra torreyana</i>	0–22	–
	fourwing saltbush	ATCA2	<i>Atriplex canescens</i>	0–6	–
7	<b>Sub-shrubs</b>			0–11	
	broom snakeweed	GUSA2	<i>Gutierrezia sarothrae</i>	0–11	–
	hairy crinklemat	TIHI	<i>Tiquilia hispidissima</i>	0–11	–
8	<b>Succulents</b>			34–56	
	soaptree yucca	YUEL	<i>Yucca elata</i>	34–56	–

## Animal community

Characteristic wildlife includes little striped whiptail, White Sands prairie lizard, bleached earless lizard, spadefoot toad, Apache pocket mouse, White sands woodrat, spotted ground squirrel, camel cricket, tiger beetle, lycosid spider, and scorpions. Ord's kangaroo rat and spotted ground squirrel.

## Recreational uses

This site has potential for camping, hiking, and picnicking. Photography and bird watching for numerous birds, raptors and others can be fair to good, especially during migration seasons. Most small animals of the site are nocturnal and secretive, seen only at night, early morning or evening. Scenic beauty is greatest during spring and sometimes summer months when flowering of forbs, shrubs, and yucca occurs.



## Wood products

This site provides no wood products other than very limited groves of cottonwood trees.

## Other products

Gypsum

## Other references

Fryberger, S.G., 2000. Geological overview of White Sands National Monument.  
<http://www.nature.nps.gov/geology/parks/whsa/geows/index.htm>

Langford, R.P., Rose, J.M., White, D.E., 2009. Groundwater salinity as a control on development of eolian landscape: An example from White Sands of New Mexico. *Geomorphology* 105(2009) 39-49.

Kocurek, G., Carr, M., Ewing, R., Havholm, K.G., Nagar, Y.C., Singhvi, a.K. 2007. White Sands Dune Field, New Mexico: Age, dune dynamics and recent accumulations. *Sedimentary Geology* 197 (2007) 313-331

## Contributors

David Trujillo

## Rangeland health reference sheet

Interpreting Indicators of Rangeland Health is a qualitative assessment protocol used to determine ecosystem condition based on benchmark characteristics described in the Reference Sheet. A suite of 17 (or more) indicators are typically considered in an assessment. The ecological site(s) representative of an assessment location must be known prior to applying the protocol and must be verified based on soils and climate. Current plant community cannot be used to identify the ecological site.

Author(s)/participant(s)	
Contact for lead author	
Date	
Approved by	
Approval date	
Composition (Indicators 10 and 12) based on	Annual Production

## Indicators

### 1. Number and extent of rills:

---

### 2. Presence of water flow patterns:

---

### 3. Number and height of erosional pedestals or terracettes:

---

### 4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are not



bare ground):

---

5. **Number of gullies and erosion associated with gullies:**

---

6. **Extent of wind scoured, blowouts and/or depositional areas:**

---

7. **Amount of litter movement (describe size and distance expected to travel):**

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):**

---

9. **Soil surface structure and SOM content (include type of structure and A-horizon color and thickness):**

---

10. **Effect of community phase composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:**

---

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):**

---

12. **Functional/Structural Groups (list in order of descending dominance by above-ground annual-production or live foliar cover using symbols: >>, >, = to indicate much greater than, greater than, and equal to):**

Dominant:

Sub-dominant:

Other:

Additional:

---

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):**

---

14. **Average percent litter cover (%) and depth ( in):**

---

15. **Expected annual annual-production (this is TOTAL above-ground annual-production, not just forage annual-production):**

- 
16. **Potential invasive (including noxious) species (native and non-native).** List species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicators, we are describing what is NOT expected in the reference state for the ecological site:
- 

17. **Perennial plant reproductive capability:**
-